arXiv:1208.4334v1 [math.NT] 21 Aug 2012

REAL ALGEBRAIC NUMBER THEORY II.
IDEOLOGICAL ARITHMETIC

T.M. GENDRON

Abstract. The arithmetic of diophantine approximation groups ($\mathbb{Z}$-integral, K-integral, matrix and polynomial) is defined in a way which extends the ideal-theoretic arithmetic of algebraic number theory, using the structure of an *ideology*: a bifiltration by subgroups defined by denominator growth and error decay.

Contents

Introduction 1
1. Tropical Growth-Decay Semi-Ring 7
2. Growth-Decay Filtration 10
3. Nonvanishing Spectra 13
4. Flat Spectra 18
5. Ideological Arithmetic 21
6. Flat Arithmetic 25
7. Symmetric Diophantine Approximations and Lorentzian Groupologies 29
8. Ideological Arithmetic of $\mathcal{O}$-Approximation Groups 38
9. Matrix Ideological Arithmetic 44
10. Polynomial Ideology and Rational Maps 50
11. Polynomial Nonvanishing Spectra and Mahler's Classification 56
12. Resultant Arithmetic 63
13. Polynomial Ideological Arithmetic I: Finite Degree 66
14. Polynomial Growth-Decay Arithmetic II: Infinite Degree 70
15. Epilogue: Ideological Classes 76
References 79

Introduction

In the first paper in this series [11], we introduced, for $\theta \in \mathbb{R}$, the diophantine approximation group ${}^*\mathbb{Z}(\theta)$, as well as variants in which $\mathbb{Z}$ is replaced by the ring of integers $\mathcal{O}$ of a finite extension $K/\mathbb{Q}$ or by the polynomial ring $\mathbb{Z}[X]$, or θ is replaced by a real matrix Θ. The focus of that study was the relationship between diophantine approximation groups, Kronecker foliations and linear/algebraic independence.

Date: 6 August 2012.

2000 *Mathematics Subject Classification.* Primary, 11J99, 11U10, 11R99.

Key words and phrases. Diophantine approximation groups, ideological arithmetic, Mahler classification, resultant arithmetic .

In this second installment we turn to the issue of arithmetic, motivated by a desire to answer the following fundamental question: let ${}^*m \in {}^*\mathbb{Z}(\theta)$, ${}^*n \in {}^*\mathbb{Z}(\eta)$ be diophantine approximations with ${}^*m\theta \simeq {}^*m^\perp$, ${}^*n\eta \simeq {}^*n^\perp$ where ${}^*m^\perp, {}^*n^\perp \in {}^*\mathbb{Z}$.

Question. *Under what conditions and with what meaning can the numerator-denominator pairs* $(m^\perp, {}^*m)$, $(n^\perp, {}^*n)$ *be manipulated as ring elements: that is, when do the pairs*

$$({}^*m^{\perp}\,{}^*n^{\perp}, {}^*m\,{}^*n), \quad ({}^*m\,{}^*n^{\perp} \pm {}^*m^{\perp}\,{}^*n, {}^*m\,{}^*n)$$

define diophantine approximations of $\theta\eta$, $\theta \pm \eta$*?*

The present paper consists of a detailed response to this question, articulated in such a way as to be compatible with the arithmetic of ideals in algebraic number theory. It is our hope that the approach developed here will provide a setting for a unified theory of algebraic and transcendental numbers.

One could set out to develop the arithmetic of diophantine approximation groups using the method of groupoids, in which each diophantine approximation is regarded as a morphism in a category, and its domain – the set of all elements composable with it – becomes the object of study. However this focus on individual elements is at odds with the communal spirit of the theory of ideals, in which pairs of collections of elements (ideals) are composed with a uniform result.

So instead, we look to find substructures of diophantine approximation groups which are defined by certain quantitative measures, then describe conditions which allow us to compose such substructures *en bloc*, rather than according to the vagaries of individual elements. This is not to call into question the groupoid method, but to distinguish it from the somewhat experimental approach adopted here.

There are two quantitative measures of a diophantine approximation $({}^*n^\perp, {}^*n)$ that have defined the field of Diophantine Approximation since the time of Dirichlet and Liouville:

1. The growth of the denominator ${}^*n \in {}^*\mathbb{Z}(\theta)$.
2. The decay of the error term $\varepsilon({}^*n) = \theta\,{}^*n - {}^*n^\perp \in {}^*\mathbb{R}_\varepsilon$.

We measure these in the following way. Let

$$\langle\cdot\rangle : {}^*\mathbb{R} \longrightarrow {}^{\star}\mathbb{PR} := {}^*\mathbb{R}/{}^*\mathbb{R}^\times_{\rm fin}$$

be the Krull valuation on ${}^*\mathbb{R}$ associated to the local ring ${}^*\mathbb{R}_{\rm fin} \subset {}^*\mathbb{R}$ of bounded nonstandard reals. The ordered group ${}^{\star}\mathbb{PR} = {}^*\mathbb{R}/{}^*\mathbb{R}^\times_{\rm fin}$ of this valuation is a tropical semi ring with respect to operations $\cdot, +$ induced from their counterparts on ${}^*\mathbb{R}$, the *growth-decay semi ring.* An important feature of ${}^{\star}\mathbb{PR}$ is the existence of a flow by tropical automorphisms: $\mu \mapsto \mu^{{}^*r}$, ${}^*r \in {}^*\mathbb{R}^\times_{\rm fin}$, the *Frobenius flow.* The quotient by the Frobenius flow gives rise to a second tropical semi ring ${}^{\star}\overline{\mathbb{PR}}$ called the *Frobenius growth-decay semi-ring.* See §1.

For ${}^*n \in {}^*\mathbb{Z}(\theta)$ we define its growth to be $\mu({}^*n) := \langle {}^*n^{-1}\rangle$ and its decay to be $\nu({}^*n) := \langle \varepsilon({}^*n)\rangle$. Then for each pair $\mu, \nu \in {}^{\star}\mathbb{PR}_\varepsilon$ = the infinitesimal part of ${}^{\star}\mathbb{PR}$,

$${}^*\mathbb{Z}^\mu_\nu(\theta) = \{{}^*n \in {}^*\mathbb{Z}(\theta) \mid \mu < \mu({}^*n),\ \nu({}^*n) \leq \nu\}$$

is a subgroup of ${}^*\mathbb{Z}(\theta)$. The bi-filtered group structure ${}^*\mathbb{Z}(\theta) = \{{}^*\mathbb{Z}^\mu_\nu(\theta)\}$ is referred to as an *ideology*; it is natural with respect to $\mathrm{PGL}_2(\mathbb{Z})$-equivalence of real numbers. See §2.

The concept of an ideology generalizes that of ideal in the following sense: if we consider the filtration ${}^*\mathbb{Z} = \{{}^*\mathbb{Z}^{\nu}\}$ where ${}^*\mathbb{Z}^{\nu} = \{{}^*n|\ \nu < \mu({}^*n)\}$ then for each ν,

$${}^*\mathbb{Z}^{\nu} \cdot {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \subset {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta).$$

By forgetting the indices one recovers the usual definition of an ideal. The notion of a partially defined product of algebraic structures deployed along a filtration of substructures, while unorthodox, seems nevertheless natural in this context.

Determining when the subgroup ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ is non trivial is the first problem which we address. We define the *nonvanishing spectrum* of θ to be

$$\mathrm{Spec}(\theta) = \{(\mu,\nu)|\ {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \neq 0\}.$$

For $\theta \notin \mathbb{Q}$, the nonvanishing spectrum is delimited by growth-decay pairs corresponding to *best denominator classes*: diophantine approximations ${}^*\widehat{q}$ built from the sequence of denominators of principal convergents of θ. In §3, we characterize the rational, badly approximable, (very) well approximable and Liouville numbers in terms of their nonvanishing spectra as well as the density of the set of *best growths*: the set of growths of best denominator classes. Among the irrational numbers, the badly approximable numbers have the smallest nonvanishing spectrum (lying above the line $\mu = \nu$) and the largest best growth set ($= {}^*\mathbb{PR}_{\varepsilon}$); the Liouville numbers have the largest nonvanishing spectra and the sparsest best growth set. This is discussed in §3. The intersection $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ of $\mathrm{Spec}(\theta)$ with the line $\mu = \nu$ represents a critical divide called the *flat spectrum.* In §4 we study the flat spectrum, proving that for $\theta \in \mathbb{R} - \mathbb{Q}$ it is a proper subset of ${}^*\mathbb{PR}_{\varepsilon}$ consisting of a union of intervals bounded by best growth-decay pairs having infinite partial quotient.

The Question posed above is answered in §5 in the ideological sense: there is a bilinear map

$$\text{(1)}\qquad {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \times {}^*\mathbb{Z}^{\nu}_{\mu}(\eta) \longrightarrow {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta\eta) \cap {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta+\eta) \cap {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta-\eta)$$

defined by the ordinary product in ${}^*\mathbb{Z}$. When $\theta = a/b$, $\eta = c/d \in \mathbb{Q}$, (1) reduces to the product of principal ideals ${}^*(c) \cdot {}^*(d) = {}^*(cd)$ generated by the denominators. That the image is contained in the groups associated to the product as well the sum and difference of θ and η is the key to understanding the relationship between *ideological arithmetic* and classical ideal theoretic arithmetic.

For $\mu \geq \nu$ we define the relation

$$\theta\,{}_{\mu}\!\oslash_{\nu}\,\eta$$

whenever the groups appearing in the product (1) are nontrivial i.e. for $(\mu,\nu) \in \mathrm{Spec}(\theta)$, $(\nu,\mu) \in \mathrm{Spec}(\eta)$. The remainder of §5 is devoted to analyzing the various classes of real numbers with respect to the relation ${}_{\mu}\!\oslash_{\nu}$. The badly approximable numbers $\mathfrak{B}$ are *antiprimes*: the relation ${}_{\mu}\!\oslash_{\nu}$ is empty on $\mathfrak{B}^2$ for all pairs (μ,ν). On other hand, the very well approximable numbers $\mathfrak{W}_{>1}$ are *omnidivisors*: for any $\theta \in \mathfrak{W}_{>1}$ and $\eta \in \mathbb{R}$, there exist $\mu \geq \nu$ such that $\theta\,{}_{\mu}\!\oslash_{\nu}\,\eta$.

Ideological arithmetic is somewhat more mysterious in the case of the flat product (which amounts to the consideration of the flat relation ${}_{\mu}\!\oslash_{\mu}$) since the exponent of $\theta \in \mathfrak{W}$ (= the well approximable numbers) tells us very little about its prospects for forming flat products. The relevant invariant here is the partial quotient representation $\theta = [a_1a_2\ldots]$. For example, when θ is *resolute* i.e. $\lim a_i = \infty$, then for any $\eta \in \mathfrak{W}$ and $\mu \in \mathrm{Spec}_{\mathrm{flat}}(\eta)$, $\theta\,{}_{\mu}\!\oslash_{\mu}\,\eta$. More generally, we show that if $\theta \in \mathfrak{W}$ is *abyssless* – having a maximally connected flat spectrum – then $\mathrm{Spec}_{\mathrm{flat}}(\theta) \supset \mathrm{Spec}_{\mathrm{flat}}(\eta)$

for all $\eta \in \mathfrak{W}$, and hence $\theta_{\mu} \text{\textcircled{$\flat$}}_{\mu}\, \eta$ for all $\mu \in \mathrm{Spec}_{\mathrm{flat}}(\eta)$. On the other hand, there exist $\theta, \eta \in \mathfrak{W}$ which are *abyssal* (not abyssless), for which the flat product is never defined. These and other results are discussed in §6.

In the field of Diophantine Approximation, one often restricts attention to the set

$$^{*}\mathbb{Z}(\theta|\psi) = \{^{*}n \in {}^{*}\mathbb{Z}(\theta) |\, |\varepsilon(^{*}n)| < |\psi(^{*}n)|\}$$

for some decay function $\psi : {}^{*}\mathbb{Z} \to {}^{*}\mathbb{R}$. When $\psi(x) = x^{-1}$, ${}^{*}\mathbb{Z}(\theta|x^{-1})$ is the set of elements of bounded θ-*norm*

$$|^{*}n|_{\theta} := (|^{*}n| \cdot |\varepsilon(^{*}n)|)^{1/2} \mod {}^{*}\mathbb{R}_{\varepsilon}.$$

In §7 we show that ${}^{*}\mathbb{Z}(\theta|x^{-1})$ has the structure of a *groupology*: with respect to the grading ${}^{*}\mathbb{Z}(\theta|x^{-1}) = \{{}^{*}\mathbb{Z}^{\mu}_{\nu}(\theta|x^{-1})\}$ there is a sum

$$^{*}\mathbb{Z}(\theta|x^{-1})^{\mu}_{\nu} + {}^{*}\mathbb{Z}(\theta|x^{-1})^{\nu}_{\mu} \subset {}^{*}\mathbb{Z}(\theta|x^{-1})^{\mu - \nu} \tag{2}$$

where $\mu - \nu = \min(\mu, \nu)$.

The θ-norm satisfies the triangle inequality with respect to all sums defined by (2), but only in a trivial sense since $|^{*}n|_{\theta} \equiv 0$ on ${}^{*}\mathbb{Z}(\theta|x^{-1})^{\mu}_{\nu}$ for $\mu \geq \nu$. This leads us to consider a refinement of the growth-decay groupology defined on the set of *symmetric diophantine approximations*

$$^{*}\mathbb{Z}^{\mathrm{sym}}(\theta) = \{^{*}n \in {}^{*}\mathbb{Z}(\theta) |\, 0 < |^{*}n|_{\theta} < \infty\} \subset {}^{*}\mathbb{Z}(\theta|x^{-1}).$$

We show that ${}^{*}\mathbb{Z}^{\mathrm{sym}}(\theta)$ is non trivial, containing the intermediate best classes as well as many other elements, and that it has the structure of a uni-indexed groupology, see Theorem 20 of §7.

In the case of $\theta = \varphi =$ the golden mean, we give an explicit description of the elements of ${}^{*}\mathbb{Z}^{\mathrm{sym}}(\varphi)$ using the Zeckendorf representations of natural numbers. The latter may be useful in the consideration of the Littlewood conjecture:

$$\liminf_{n} n\|n\theta\|\|n\eta\| = 0, \quad \theta, \eta \in \mathfrak{B},$$

which is clearly implied by the statement

$$^{*}\mathbb{Z}^{\mathrm{sym}}(\theta) \cap {}^{*}\mathbb{Z}(\eta) \neq \emptyset \quad \text{or} \quad {}^{*}\mathbb{Z}^{\mathrm{sym}}(\eta) \cap {}^{*}\mathbb{Z}(\theta) \neq \emptyset, \quad \theta, \eta \in \mathfrak{B}.$$

The restriction of $|\cdot|_{\theta}$ to ${}^{*}\mathbb{Z}^{\mathrm{sym}}(\theta)$ is not subadditive: rather, it satisfies the reverse triangle inequality, due to the fact that it most naturally arises from a Lorentzian bilinear pairing of signature $(1,1)$ on ${}^{*}\mathbb{Z}^{\mathrm{sym}}(\theta)$. If we view diophantine approximations as material particles departing from θ then $|^{*}n|_{\theta}$ is nothing more than the initial speed; for badly approximable numbers, we have the Heisenberg's uncertaintly principle

$$|^{*}n|_{\theta} > C_{\theta}$$

where C_{θ} is the corresponding element of the Lagrange spectrum. More generally, we consider a $\mathrm{PGL}_2(\mathbb{Z})$-natural family of "Lorentzian norms" $|\cdot|_{\theta, {}^{*}\kappa}$ for each non-standard exponent ${}^{*}\kappa > 1$ and we obtain a collection of groupologies $\{{}^{*}\mathbb{Z}(\theta|x^{-{}^{*}\kappa})\}$ called a *Frechet-Lorentzian groupology.*

✠

The remainder of the paper concerns extensions of the first part to:

A. Diophantine approximation of elements of the Minkowski space $\mathbb{K}$ of K by the ring $\mathcal{O}$ of K-integers, where $K/\mathbb{Q}$ is a finite field extension.

B. Diophantine approximation of real matrices (families of linear forms) by vectors.

C. Polynomial (nonlinear) diophantine approximation of real numbers.

We will not attempt to duplicate all of the results found in the previous sections, an endeavor falling outside of the intended scope of the present work. Rather, the imperative is to set up the basic apparatus of the previous sections in such a way that the theory in each of A.–C. may be seen to be a natural arithmetic extension of classical Diophantine Approximation. The third paper in this series will deal with the Galois theory of these extensions.

A. For $K/\mathbb{Q}$ a finite extension, growth and decay are measured in the tropical semi ring ${}^{\star}\mathbb{PK} \cong {}^{\star}\mathbb{PR}^{r+s}$ (where r, s are the number of real and complex-pair places). The diophantine approximation group of $\boldsymbol{z} \in \mathbb{K}$ has the structure of an ideology

$$ {}^{*}\mathcal{O}(\boldsymbol{z}) = \{ {}^{*}\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \}, $$

and these K-ideologies may be composed according to an obvious analogue of (1). If $K/\mathbb{Q}$ is Galois, then the action of $\mathrm{Gal}(K/\mathbb{Q})$ on $\mathbb{K}$ extends to an action on growth-decay indices so that the growth-decay product becomes Galois natural, c.f. Proposition 25. The K-nonvanishing spectrum $\mathrm{Spec}_{\mathcal{O}}(\boldsymbol{z})$ may be used to define the nontrivial classes of K-badly approximable, K-(very) well approximable and K-Liouville points in $\mathbb{K}$. In this connection one observes the phenomenon of antiprime splitting, where a $\mathbb{Q}$-badly approximable number θ loses its antiprime status upon diagonal inclusion in $\mathbb{K}$: this happens for quadratic Pisot-Vijayaraghavan numbers, see Theorem 26.

B. Given Θ a real $r \times s$ matrix (or in the classical language: a family of r linear forms in s variables), the matrix ideology

$$ {}^{*}\mathbb{Z}^{s}(\Theta) = \{ ({}^{*}\mathbb{Z}^{s})^{\mu}_{\nu}(\Theta) \} $$

is the subject of §9. Growth and decay are measured by the house norm and the ideological product gives rise to a fractional arithmetic on the set of all real matrices $\tilde{\mathcal{M}}(\mathbb{R})$ based on the Kronecker product, as well as an arithmetic based on the Kronecker sum of matrices on the subset $\mathcal{M}(\mathbb{R}) \subset \tilde{\mathcal{M}}(\mathbb{R})$ of square matrices. The classes of badly approximable, very well approximable and Liouville matrices are characterized (or defined) by the shape of their associated nonvanishing spectra. In the special case of a single form the dual groups give rise to an ideological arithmetic of *nonprincipal* ideologies. The projective linear group $\mathrm{PGL}_{r,s}(\mathbb{Z}) := \mathrm{PGL}_{r+s}(\mathbb{Z})$ (its elements partitioned into $r \times r, r \times s, s \times r$ and $s \times s$ blocks) acts on the set of $r \times s$ matrices $\tilde{\mathcal{M}}_{r,s}(\mathbb{R})$ inducing ideological isomorphisms that preserve the matrix ideological product.

C. The most elaborate version of ideological arithmetic that we encounter arises in the setting of polynomial diophantine approximation. Due to the role power laws play in Mahler's classification it is natural to introduce ideologies based on the Frobenius semi ring ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ in which elements of ${}^{\star}\mathbb{PR}_{\varepsilon}$ define subideologies. Denote ${}^{*}\tilde{\mathfrak{Z}} = {}^{*}(\mathbb{Z}[X])$ the ring of (possibly infinite degree) nonstandard polynomials and ${}^{*}\mathfrak{Z} := ({}^{*}\mathbb{Z})[X]$ the subring of finite degree polynomials. For each θ we define the (degree filtered) *Frobenius ideology*

$$ {}^{*}\tilde{\mathfrak{Z}}(\theta) = \{ {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}\breve{d}} \} $$

for $\bar{\mu}, \bar{\nu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ and ${}^{*}\breve{d}$ the Frobenius class of the degree ${}^{*}d$. See §10 for details. Each ${}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}\breve{d}}$ is in fact a subring of ${}^{*}\tilde{\mathfrak{Z}}(\theta)$, and since any Frobenius orbit $\bar{\mu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ is itself a tropical semi-ring we may define in turn the ideological refinement:

$$ {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}\breve{d}} = \{ {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu},\mu}_{\bar{\nu},\nu}(\theta)_{{}^{*}\breve{d},{}^{*}d} \} $$

for all $\mu \in \bar{\mu}, \nu \in \bar{\nu}, {}^{*}d \in {}^{*}\breve{d}$. The latter is called the *ordinary ideological structure.*

It turns out that the external sum of polynomials is not of arithmetic interest; rather it is the *Cauchy product* of polynomials that plays the role of the subordinate (acted upon) operation in polynomial arithmetic. In this connection, the monoid of integral rational maps $(\mathsf{Rat}_{\mathbb{Z}}, \circ)$ with the operation of composition provides the relevant notion of equivalence (taking over the role of $\mathrm{PGL}_2(\mathbb{Z})$ in linear diophantine approximation), acting naturally on the above ideologies viewed as families of Cauchy semigroups. Since the elements of $(\mathsf{Rat}_{\mathbb{Z}}, \circ)$ are not invertible, we introduce the relation $\theta \circeq \eta$ whenever there is a pair (R, S) of rational maps for which $R(\theta) = \eta$ and $S(\eta) = \theta$; when this happens we obtain an *ideological consensus* ${}^{*}\tilde{\mathfrak{Z}}(\theta) \rightleftharpoons {}^{*}\tilde{\mathfrak{Z}}(\eta)$: a pair of Cauchy monomorphisms respecting the Frobenius ideology and acting as a multiplicative shift on the ordinary ideology. See Theorem 37 of §10.

In §11 we study the Frobenius nonvanishing spectrum

$$ \overline{\mathrm{Spec}}[X](\theta) = \{ \overline{\mathrm{Spec}}[X](\theta)_{{}^{*}\breve{d}} \}, $$

which is viewed as a directed set, as well as the filtration of subsets defined by ${}^{*}d \in {}^{*}\breve{d}$. We characterize the Mahler classes and Mahler types according to the portraits of their nonvanishing spectra e.g. see Figure 3 and Theorems 38–41.

The definition of ideological arithmetic in the polynomial setting is dictated by the wholly reasonable demand that the inclusion ${}^{*}\mathbb{Z}(\theta) \hookrightarrow {}^{*}\tilde{\mathfrak{Z}}(\theta)$ be ideological. The nonlinear analogue of fractional arithmetic of linear equations is provided by the *resultant sum* $f \diamondplus g$ and *resultant product* $f \diamondtimes g$ of polynomials, which are essentially characterized by the property that their root sets are the sum and the product, respectively, of the root sets of f and g. It seems fair to assume that the resultant sum and product are known; nevertheless in §12 we develop their basic properties from scratch. Both $\diamondplus$ and $\diamondtimes$ distribute over the Cauchy product; if we let $\check{\mathfrak{Z}}$ be the Cauchy monoid of non-0 polynomials modulo multiplication by non-0 integers, then $\diamondplus$ and $\diamondplus$ define operations on $\check{\mathfrak{Z}}$ making the latter a double monoid, in which the inclusion $\mathbb{Q} \hookrightarrow \check{\mathfrak{Z}}$, $a/b \mapsto aX - b$, takes the operations $+$ to $\diamondplus$ and $\times$ to $\diamondtimes$. Thus we can view $\check{\mathfrak{Z}}$ as a generalized field extension of $\mathbb{Q}$; since $\diamondtimes$ does not distribute over $\diamondplus$, it is a spiritual relative of the nonlinear number field construction of [15].

Ideological arithmetic of the resultant product $\diamondtimes$ for finite degree polynomials is studied in §13; the infinite degree case presents particular complications and is deferred to §14. The most general version of the ordinary ideological product is given by Theorem 48 of §14:

$$ {}^{*}\tilde{\mathfrak{Z}}^{\mu}_{(\mu^{1-{}^{*}e^{-1}}\cdot\nu)^{{}^{*}e}}(\theta)_{{}^{*}d} \times {}^{*}\tilde{\mathfrak{Z}}^{\nu}_{(\mu\cdot\nu^{1-{}^{*}d^{-1}})^{{}^{*}d}}(\eta)_{{}^{*}e} \xrightarrow{\diamondtimes} {}^{*}\tilde{\mathfrak{Z}}^{\mu\cdot\nu}(\theta\eta)_{{}^{*}d{}^{*}e} $$

which agrees with (1) on the images of ${}^{*}\mathbb{Z}(\theta) \subset {}^{*}\mathfrak{Z}(\theta), {}^{*}\mathbb{Z}(\eta) \subset {}^{*}\mathfrak{Z}(\eta)$; assuming $\bar{\mu} \geq \bar{\nu}$ it reduces to

$$ {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}^{{}^{*}e}}(\theta)_{{}^{*}d} \times {}^{*}\tilde{\mathfrak{Z}}^{\bar{\nu}}_{\bar{\nu}^{{}^{*}d}}(\eta)_{{}^{*}e} \xrightarrow{\diamondtimes} {}^{*}\tilde{\mathfrak{Z}}^{\bar{\nu}}(\theta\eta)_{{}^{*}d{}^{*}e}. \tag{3} $$

upon passage to the Frobenius ideology.

The relation

$$\theta \; {}^{*}\!\underset{\bar{\upmu}}{\overset{d}{}} \rotatebox{45}{$\boxplus$} {}^{*}\!\underset{\bar{\nu}}{\overset{e}{}} \; \eta$$

is defined whenever the Cauchy semigroups appearing in (3) are nonempty; at the end of §14 we describe its satisfaction according to the Mahler classes of its arguments as well as the growth, decay and degree parameters. For θ an A-number (an algebraic number), its composability with other reals hinges upon whether the corresponding approximation semigroup is the ideal generated by the minimal polynomial or not. The former case is optimal, and θ imposes no restrictions on composability whatsoever, see Theorem 50; in the latter case, the conditions imposed by θ are severe, see Theorem 51. The S-numbers, modulo a technical condition on the index $\bar{\upmu}$, form an antiprime set, see again Theorem 51. The T and U numbers impose the fewest restrictions on composition: in particular, if both θ and η U-numbers, they satisfy infinitely many "flat relations" $\theta \, {}^{*}\!\underset{\bar{\upmu}}{\overset{d}{}} \rotatebox{45}{$\boxplus$} {}^{*}\!\underset{\bar{\upmu}}{\overset{d}{}} \, \eta$, where ${}^{*}d$ can be either finite or infinite, see Theorem 52.

The last section, §15, is offered as an epilogue, in which the sum/difference/product range of the K-ideological product ($K/\mathbb{Q}$ a finite degree extension) is explained in terms of the product of (nonprincipal) ideals. This is done through the *ideological class* ${}^{*}[\mathcal{O}](\boldsymbol{z})$ of the ideology ${}^{*}\mathcal{O}(\boldsymbol{z})$, defined as the K-projective class of the *decoupled diophantine approximation ideology*

$${}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}) := {}^{*}\mathcal{O}(\boldsymbol{z}) + {}^{*}\mathcal{O}(\boldsymbol{z})^{\perp}.$$

If $\mathfrak{a} = (\alpha, \beta) \subset \mathcal{O}$ is a classical ideal and $\gamma = \alpha/\beta$ then ${}^{*}[\mathcal{O}](\gamma)$ coincides with the ideal class ${}^{*}[\mathfrak{a}]$ of ${}^{*}\mathfrak{a}$. The quotient $\mathcal{C}\ell(\mathbb{K}) := \mathrm{PGL}_2(\mathcal{O})\backslash\mathbb{K}$ parametrizes ideology classes and contains the usual ideal class group $\mathcal{C}\ell(K) \approx \mathrm{PGL}_2(\mathcal{O})\backslash K$. If we define the $\circledast$-product of ideological classes as the projective class of

$$\text{(4)} \qquad {}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}) \circledast {}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{w}) := {}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{zw}) + {}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}+\boldsymbol{w}) + {}^{*}\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}-\boldsymbol{w})$$

then the restriction of $\circledast$ to $\mathcal{C}\ell(K)$ reduces to the usual product of ideal classes. Moreover, the range of an $\mathcal{O}$-ideological product embeds canonically in the $\circledast$-product of the associated decoupled ideologies. In general, there is no reason to believe that the $\circledast$-product (4) defines an element of $\mathcal{C}\ell(\mathbb{K})$. Its proper domain of definition should be a Stone space extension of $\mathbb{K}$, in which $\mathbb{K}$ is discrete. Its determination will be postponed to [14].

Acknowledgements. I would like to thank Gregor Weingart for suggesting the proof of Theorem 45. This paper was supported in part by the CONACyT grant 058537 as well as the PAPIIT grant IN103708.

1. Tropical Growth-Decay Semi-Ring

Let ${}^{*}\mathbb{Z} \subset {}^{*}\mathbb{R}$ be ultrapowers of $\mathbb{Z} \subset \mathbb{R}$ with respect to a fixed non-principal ultrafilter on $\mathbb{N}$. As usual ${}^{*}\mathbb{R}_{\mathrm{fin}} \subset {}^{*}\mathbb{R}$ denotes the subring of bounded sequence classes and ${}^{*}\mathbb{R}_{\varepsilon} \subset {}^{*}\mathbb{R}_{\mathrm{fin}}$ the maximal ideal of infinitesimals. The subscript "+" will refer to the positive elements of any of the above sets e.g. ${}^{*}\mathbb{Z}_{+} \subset {}^{*}\mathbb{Z}$ is the subset of positive elements in ${}^{*}\mathbb{Z}$. See §1 of [13] and references contained therein for more background.

Let $({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+} =$ the group of positive units in the ring ${}^{*}\mathbb{R}_{\mathrm{fin}}$; thus $({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$ is the multiplicative subgroup of noninfinitesimal, noninfinite elements in ${}^{*}\mathbb{R}_{+}$. Consider the multiplicative quotient group

$$^*\mathbb{PR} := {}^*\mathbb{R}_+/({}^*\mathbb{R}_{\mathrm{fin}})^\times_+,$$

whose elements will be written

$$\mu = {}^*x \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+.$$

We denote the product in $^*\mathbb{PR}$ by "$\cdot$".

Proposition 1. *Every element* $\mu \in {}^*\mathbb{PR}$ *may be written in the form*

$$^*n^\varepsilon \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$$

where $^*n \in {}^*\mathbb{Z}_+ - \mathbb{Z}_+$ *or* $^*n = 1$, *and* $\varepsilon = \pm 1$.

Proof. Every element of $^*\mathbb{PR}$ is the class of 1, the class of an infinite element or the class of an infinitesimal element. If μ is the class of $^*r$ infinite, then there exists $^*\bar{r} \in [0,1) = \{^*x |\, 0 \leq {}^*x < 1\}$ and $^*n \in {}^*\mathbb{Z}_+$ for which $^*r = {}^*n + {}^*\bar{r} = {}^*n \cdot (({}^*n + {}^*\bar{r})/{}^*n)$. But $({}^*n + {}^*\bar{r})/{}^*n = 1 + {}^*\bar{r}/{}^*n \in ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$, so $\mu = {}^*n \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$. A similar argument may be used to show that when μ represents an infinitesimal class, $\mu = {}^*n^{-1} \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$ for some $^*n \in {}^*\mathbb{Z}_+ - \mathbb{Z}_+$. ☐

Proposition 2. $^*\mathbb{PR}$ *is a densely ordered group.*

Proof. The order is defined by declaring that $\mu < \mu'$ in $^*\mathbb{PR}$ if for any pair of representatives $^*x \in \mu, {}^*x' \in \mu'$ we have $^*x < {}^*x'$, evidently a dense order without endpoints. The left-multiplication action of $^*\mathbb{R}_+$ on $^*\mathbb{PR}$ preserves this order, therefore so does the product: if $\mu < \nu$ then for all $\xi \in {}^*\mathbb{PR}$, $\xi \cdot \mu < \xi \cdot \nu$. ☐

We introduce the maximum of a pair of elements in $^*\mathbb{PR}$ as a formal binary operation:

$$\mu + \nu := \max(\mu, \nu).$$

The operation + is clearly commutative and associative. The following Proposition says that + is the quotient of the operation + of $^*\mathbb{R}^\times_+$.

Proposition 3. *Let* $\mu = {}^*x \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$, $\mu' = {}^*x' \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$. *Then*

$$({}^*x + {}^*x') \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+ = \mu + \mu'.$$

Proof. Note that $^*x + {}^*x' \in {}^*\mathbb{R}_+$ and $^*x + {}^*x' \in \max(\mu, \mu')$. Indeed, suppose first that $\mu \neq \mu'$, say $\mu < \mu'$. Then there exists $^*\varepsilon$ infinitesimal for which $^*x = {}^*\varepsilon {}^*x'$, and we have $^*x' + {}^*x = {}^*x'(1 + {}^*\varepsilon) \in \mu'$. If $\mu = \mu'$ then $^*x' = {}^*r {}^*x$ for $^*r \in {}^*\mathbb{R}^\times_+$ and $({}^*x + {}^*x') \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+ = {}^*x(1 + {}^*r) \cdot ({}^*\mathbb{R}_{\mathrm{fin}})^\times_+ = \mu = \mu + \mu$. ☐

Proposition 4. *Let* $^*r, {}^*s \in {}^*\mathbb{R}_+$ *and* $\mu, \nu, \nu' \in {}^*\mathbb{PR}$. *Then*

1. $\mu \cdot (\nu + \nu') = (\mu \cdot \nu) + (\mu \cdot \nu')$.
2. $^*r \cdot (\nu + \nu') = ({}^*r \cdot \nu) + ({}^*r \cdot \nu')$.
3. $({}^*r + {}^*s) \cdot \mu = ({}^*r \cdot \mu) + ({}^*s \cdot \mu)$.

Proof. 1. It is enough to check the equality in the case $\nu' > \nu$. Then $\mu \cdot (\nu + \nu') = \mu \cdot \nu'$. But the latter is equal to $(\mu \cdot \nu) + (\mu \cdot \nu')$ since the product preserves the order. The proof of 2. is identical, where we use the fact that the multiplicative action by $^*\mathbb{R}_+$ preserves the order. Item 3. is trivial. ☐

It will be convenient to add the class $-\infty$ of the element $0 \in {}^*\mathbb{R}$ to the space ${}^*\mathbb{PR}$: in other words, we will reconsider ${}^*\mathbb{PR}$ as the quotient $({}^*\mathbb{R}_+ \cup \{0\})/({}^*\mathbb{R}_{\rm fin})^\times_+$. Note that we have for all $\mu \in {}^*\mathbb{PR}$

$$-\infty + \mu = \mu, \quad -\infty \cdot \mu = -\infty.$$

In particular, $-\infty$ is the neutral element for the operation $+$. Thus, by Proposition 4:

Theorem 1. ${}^*\mathbb{PR}$ *is an abstract (multiplicative) tropical semi-ring: that is, a max-times semi ring.*

We will refer to ${}^*\mathbb{PR}$ as the **growth-decay semi-ring**. Let ${}^*\mathbb{PR}_\varepsilon \subset {}^*\mathbb{PR}$ be the image of the $({}^*\mathbb{R}_{\rm fin})^\times_+$-invariant multiplicatively closed set $({}^*\mathbb{R}_\varepsilon)_+$. With the operations $\cdot, +$, ${}^*\mathbb{PR}_\varepsilon$ is a sub tropical semi-ring: the **decay semi-ring**.

If we forget the tropical addition, considering ${}^*\mathbb{PR}$ as a linearly ordered multiplicative group, then the map

$$\langle \cdot \rangle : {}^*\mathbb{R} \to {}^*\mathbb{PR}, \quad \langle {}^*x \rangle = |{}^*x| \cdot ({}^*\mathbb{R}_{\rm fin})^\times_+,$$

is the Krull valuation associated to the local ring ${}^*\mathbb{R}_{\rm fin}$ (see for example [28]). The restriction of $\langle \cdot \rangle$ to $\mathbb{R}$ is just the trivial valuation, so that $\langle \cdot \rangle$ cannot be equivalent to the usual valuation $|\cdot|$ on ${}^*\mathbb{R}$ induced from the euclidean norm. Note also that $\langle \cdot \rangle$ is nonarchimedean. We refer to $\langle \cdot \rangle$ as the **growth-decay valuation**.

There is a natural "Frobenius" action of the multiplicative group $\mathbb{R}^\times_+$ on ${}^*\mathbb{PR}$: for $\mu \in {}^*\mathbb{PR}_\varepsilon$ and ${}^*x \in \mu$ define

$$\Phi_r(\mu) = \mu^r := {}^*x^r \cdot ({}^*\mathbb{R}_{\rm fin})^\times_+$$

for each $r \in \mathbb{R}^\times_+$. Note that this action does not depend on the choice of representative ${}^*x$. We may extend the Frobenius action to $({}^*\mathbb{R}_{\rm fin})^\times_+$ as follows. For ${}^*r = {}^*\{r_i\} \in ({}^*\mathbb{R}_{\rm fin})^\times_+$ and $\mu \in {}^*\mathbb{PR}$ represented by ${}^*x = {}^*\{x_i\} \in {}^*\mathbb{R}_+$ define

$$\Phi_{{}^*r}(\mu) = \mu^{{}^*r} := {}^*\{x_i^{r_i}\} \cdot ({}^*\mathbb{R}_{\rm fin})^\times_+,$$

which is again well-defined. Note that it is *not* the case that if ${}^*r \simeq r \in \mathbb{R}_+$ that $\mu^{{}^*r} = \mu^r$.

Theorem 2. *The map* $\Phi_{{}^*r} : {}^*\mathbb{PR} \to {}^*\mathbb{PR}$ *is a tropical automorphism for each* ${}^*r \in ({}^*\mathbb{R}_{\rm fin})^\times_+$ *and defines a faithful representation*

$$\Phi : ({}^*\mathbb{R}_{\rm fin})^\times_+ \longrightarrow \mathrm{Aut}({}^*\mathbb{PR}).$$

Proof. $\Phi_{{}^*r}$ is clearly multiplicative. Moreover: $(\mu+\nu)^{{}^*r} = (\max(\mu,\nu))^{{}^*r} = \mu^{{}^*r} + \nu^{{}^*r}$. □

We denote by $\bar{\mu}$ the orbit of μ by $({}^*\mathbb{R}_{\rm fin})^\times_+$ with respect to Φ. Note that by Theorem 2:

- $\bar{\mu}$ is a sub tropical semi-ring of ${}^*\mathbb{PR}$.
- The quotient of ${}^*\mathbb{PR}$ by Φ, denoted ${}^*\overline{\mathbb{PR}}$, is a tropical semi-ring.

We denote by $[{}^*x]$ the image in ${}^*\overline{\mathbb{PR}}$ of ${}^*x \in {}^*\mathbb{R}$. For all $\bar{\mu}, \bar{\nu} \in {}^*\overline{\mathbb{PR}}$, we write $\bar{\mu} < \bar{\nu}$ $\Leftrightarrow$ for all $\mu \in \bar{\mu}$, $\nu \in \bar{\nu}$, $\mu < \nu$.

Proposition 5. ${}^*\overline{\mathbb{PR}}$ *is a dense linear order.*

Proof. If $\bar{\mu} \nleq \bar{\nu}$ and $\bar{\mu} \ngeq \bar{\nu}$ then it follows that there exist representatives $\mu \in \bar{\mu}$, $\nu \in \bar{\nu}$ for which $\mu < \nu$ and $\mu^{{}^{*}r} > \nu$ for ${}^{*}r \in ({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$. We may assume without loss of generality that both μ, ν represent infinite classes so that ${}^{*}r > 1$. Representing ${}^{*}x = {}^{*}\{x_i\} \in \mu$ and ${}^{*}y = {}^{*}\{y_i\} \in \nu$, let ${}^{*}s = {}^{*}\{s_i\}$ where s_i is the unique positive real satisfying $x_i^{s_i} = y_i$. Then ${}^{*}s \in [1, {}^{*}r] \subset ({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$, $\mu^{{}^{*}s} = \nu$ and therefore $\bar{\mu} = \bar{\nu}$. Thus ${}^{\star}\overline{\mathbb{PR}}$ is a linear order. On the other hand, if $\bar{\mu} < \bar{\nu}$, then choosing representatives ${}^{*}x$, ${}^{*}y$ as above, we have ${}^{*}x^{{}^{*}s} = {}^{*}y$ for ${}^{*}s$ infinite. If we let μ' be the class of ${}^{*}x^{\sqrt{{}^{*}s}}$, then $\bar{\mu} < \bar{\mu}' < \bar{\nu}$. ☐

We call ${}^{\star}\overline{\mathbb{PR}}$ the **Frobenius growth-decay semi-ring**.

Note 1. In fact, we may extend Φ to an action of $({}^{*}\mathbb{R}_{\varepsilon})_{+}$ = the multiplicative monoid of positive infinitesimals on ${}^{\star}\mathbb{PR}$. That is, infinitesimal powers $\mu^{{}^{*}\varepsilon}$, ${}^{*}\varepsilon \in ({}^{*}\mathbb{R}_{\varepsilon})_{+}$, are well-defined, since ${}^{*}r^{{}^{*}\varepsilon} \in ({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$ for all ${}^{*}r \in ({}^{*}\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$. This induces an action of $({}^{*}\mathbb{R}_{\varepsilon})_{+}$ on ${}^{\star}\overline{\mathbb{PR}}$, $\bar{\mu} \mapsto \bar{\mu}^{{}^{*}\varepsilon}$ by tropical homomorphisms which either expand or contract the order according to whether $\bar{\mu}$ is the class of an infinitesimal or infinite element: e.g. $\bar{\mu} < \bar{\mu}^{{}^{*}\varepsilon}$ if $\bar{\mu}$ is an infinitesimal class. We will say that $\bar{\mu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ is a **${}^{*}\varepsilon$ root of unity** if $\bar{\mu}^{{}^{*}\varepsilon} = \bar{1}$. The set of ${}^{*}\varepsilon$ roots of unity clearly forms a sub tropical semi ring of ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$.

Define the **tropical subtraction** of classes $\mu, \nu \in {}^{\star}\mathbb{PR}$ by

$$\mu - \nu = \nu - \mu := \min(\mu, \nu).$$

Tropical subtraction satisfies $\mu - \nu \geq \mu \cdot \nu$, and descends to a well defined binary operation in ${}^{\star}\overline{\mathbb{PR}}$. Let ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ be the image of the decay semi-ring in ${}^{\star}\overline{\mathbb{PR}}$.

Proposition 6. *Tropical subtraction coincides with tropical product in* ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$*:* $\bar{\mu} \cdot \bar{\nu} = \bar{\mu} - \bar{\nu}$ *for all* $\bar{\mu}, \bar{\nu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$.

Proof. If $\bar{\mu} = \bar{\nu}$ then $\bar{\mu} \cdot \bar{\mu} = \bar{\mu} = \bar{\mu} - \bar{\mu}$. Now assume that $\bar{\mu} > \bar{\nu}$. Then there exists ${}^{*}\delta \in {}^{*}\mathbb{R}_{\varepsilon}$ such that $\bar{\mu} = \bar{\nu}^{{}^{*}\delta}$. Then $\bar{\mu} \cdot \bar{\nu} = \bar{\nu}^{1 + {}^{*}\delta} = \bar{\nu} = \bar{\mu} - \bar{\nu}$.

☐

2. GROWTH-DECAY FILTRATION

As in the previous section, ${}^{\star}\mathbb{PR}_{\varepsilon} \subset {}^{\star}\mathbb{PR}$ denotes the decay semi-ring. We will measure growth of an infinite element of ${}^{*}\mathbb{Z}$ in terms of the decay of its reciprocal: this has the advantage of allowing us to make the vital comparison of denominator growth with error decay of a diophantine approximation in a single, unambiguous setting.

For each ${}^{*}n \in {}^{*}\mathbb{Z}$ define its **growth** by

$$\mu({}^{*}n) := \langle |{}^{*}n^{-1}| \rangle \in {}^{\star}\mathbb{PR}_{\varepsilon} \cup \{1\};$$

note that $\mu({}^{*}n) = 1 \Leftrightarrow {}^{*}n = n \in \mathbb{Z}$. For each $\mu \in {}^{\star}\mathbb{PR}_{\varepsilon}$ denote by

$${}^{*}\mathbb{Z}^{\mu} = \{{}^{*}n \in {}^{*}\mathbb{Z} |\ \mu({}^{*}n) > \mu\} = \{{}^{*}n \in {}^{*}\mathbb{Z} |\ |{}^{*}n| \cdot \mu \in {}^{\star}\mathbb{PR}_{\varepsilon}\}.$$

Note that ${}^{*}\mathbb{Z}^{\mu}$ is a well-defined subgroup of ${}^{*}\mathbb{Z}$. If $\mu < \mu'$ then

$${}^{*}\mathbb{Z}^{\mu} \supset {}^{*}\mathbb{Z}^{\mu'}. \tag{5}$$

The collection $\{{}^{*}\mathbb{Z}^{\mu}\}$ forms an order-reversing filtration of ${}^{*}\mathbb{Z}$ by subgroups, called the **growth filtration**. Notice that

$${}^{*}\mathbb{Z}^{\mu} \cdot \mathbb{Z}^{\mu'} \subset \mathbb{Z}^{\mu \cdot \mu'}$$

so that ${}^*\mathbb{Z}$ has the structure of a filtered ring with respect to the growth filtration.

It will be useful to introduce the following subordinate filtration to the growth filtration. Fix $\mu \in {}^*\mathbb{PR}_\varepsilon$ and for each $\iota \in {}^*\mathbb{PR}_\varepsilon$ define

$${}^*\mathbb{Z}^{\mu[\iota]} := \{{}^*n |\, |{}^*n| \cdot \mu < \iota\} .$$

Then ${}^*\mathbb{Z}^{\mu[\iota]}$ is a group since by Proposition 4, item 3.,

$$({}^*m + {}^*n) \cdot \mu = {}^*m \cdot \mu + {}^*n \cdot \mu < \iota.$$

Note that if $\iota < \lambda$ then ${}^*\mathbb{Z}^{\mu[\iota]} \subset {}^*\mathbb{Z}^{\mu[\lambda]}$. We call this the **fine growth bi-filtration**. The fine growth bi-filtration makes of ${}^*\mathbb{Z}$ a bi-filtered ring:

$${}^*\mathbb{Z}^{\mu[\iota]} \cdot \mathbb{Z}^{\mu'[\iota']} \subset \mathbb{Z}^{\mu\cdot\mu'[\iota\cdot\iota']}.$$

For $\theta \in \mathbb{R}$, recall (see §2 of [13]) that by a **diophantine approximation** we mean an element ${}^*n \in {}^*\mathbb{Z}$ such that

$$\varepsilon({}^*n) := {}^*n\theta - {}^*n^{\perp} \in {}^*\mathbb{R}_\varepsilon$$

for some

$${}^*n^{\perp} = {}^*n^{\perp_\theta} \in {}^*\mathbb{Z},$$

called the **θ-dual** or simply the **dual** of ${}^*n$ if θ is understood. The **diophantine approximation group** is then

$$\text{(6)} \qquad {}^*\mathbb{Z}(\theta) = \{{}^*n \in {}^*\mathbb{Z} |\ {}^*n \text{ is a diophantine approximation of } \theta\} \subset {}^*\mathbb{Z}.$$

Write ${}^*\mathbb{Z}^{\mu}(\theta) = {}^*\mathbb{Z}^{\mu} \cap {}^*\mathbb{Z}(\theta)$ and ${}^*\mathbb{Z}^{\mu[\iota]}(\theta) = {}^*\mathbb{Z}^{\mu[\iota]} \cap {}^*\mathbb{Z}(\theta)$.

We now introduce a second filtration which is only available for the groups ${}^*\mathbb{Z}(\theta)$. Let $\nu \in {}^*\mathbb{PR}_\varepsilon$. For each ${}^*n \in {}^*\mathbb{Z}(\theta)$ write

$$\nu({}^*n) := \langle |\varepsilon({}^*n)| \rangle \in {}^*\mathbb{PR}_\varepsilon,$$

the **decay** of ${}^*n$. We define

$${}^*\mathbb{Z}_\nu(\theta) = \{{}^*n \in {}^*\mathbb{Z}(\theta) |\ \nu({}^*n) \leq \nu\}$$

which is a subgroup of ${}^*\mathbb{Z}(\theta)$: for ${}^*n, {}^*n' \in {}^*\mathbb{Z}_\nu(\theta)$, $|\varepsilon({}^*n + {}^*n')| \leq |\varepsilon({}^*n)| + |\varepsilon({}^*n)|$ and therefore $\nu({}^*n + {}^*n') \leq \nu$. Note that if $\nu < \nu'$,

$$\text{(7)} \qquad {}^*\mathbb{Z}_\nu(\theta) \subset {}^*\mathbb{Z}_{\nu'}(\theta)$$

which produces an order-preserving filtration of ${}^*\mathbb{Z}(\theta)$ called the **decay filtration**. Finally we denote the intersection

$${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) = {}^*\mathbb{Z}^{\mu}(\theta) \cap {}^*\mathbb{Z}_\nu(\theta),$$

which we refer to as the **growth-decay bi-filtration** of ${}^*\mathbb{Z}(\theta)$. In addition we have the **fine growth-decay tri-filtration** ${}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta)$.

Aside. The reader may wonder why we have chosen to use a *strict* inequality to define the growth filtration and yet a *non strict* inequality to define the decay filtration. The strict inequality in the growth filtration is needed in order to ensure a pleasing, concise formulation of the ideological product (see Theorem 11). The non strict inequality in the decay filtration is employed to take into account the fact that the strict inequality present in Dirichlet's Theorem may become non strict upon passage to the growth-decay semi ring ${}^*\mathbb{PR}$.

Proposition 7. *For all* $\mu, \nu, \iota \in {}^*\mathbb{PR}_\varepsilon$, $\nu \neq -\infty$, ${}^*\mathbb{Z}^{\mu[\iota]}(\theta)$ *and* ${}^*\mathbb{Z}_\nu(\theta)$ *are nontrivial and uncountable. For* $\nu = -\infty$, $\mathbb{Z}_{-\infty}(\theta)$ *is non trivial* $\Leftrightarrow$ $\theta \in \mathbb{Q}$.

Proof. Let μ, ν and ι be represented by sequences of positive real numbers $\{r_k\}$, $\{s_k\}$ and $\{i_k\}$ converging to 0. Let ${}^*n \in {}^*\mathbb{Z}(\theta)$ be represented by the sequence of integers $\{n_k\}$. We may choose ${}^*n$ so that $n_k r_k \to 0$; in fact, so that $|n_k r_k| < i_k$. The sequence $\{n_k\}$ may then be used to construct uncountably many elements of ${}^*\mathbb{Z}^{\mu[\iota]}(\theta)$. Similarly, we may find a class ${}^*m \in {}^*\mathbb{Z}(\theta)$ represented by $\{m_k\}$ so that $|m_k\theta - m_k^{\perp}| \leq s_k$, and the sequence $\{m_k\}$ may then be used to construct uncountably many elements of ${}^*\mathbb{Z}_{\nu}(\theta)$. The last claim in the statement of the Proposition follows from the fact that $\theta \in \mathbb{R}$ admits $0 \neq {}^*n \in {}^*\mathbb{Z}(\theta)$ with $\varepsilon({}^*n) = 0 \Leftrightarrow \theta \in \mathbb{Q}$.

□

The duality map ${}^*n \mapsto {}^*n^{\perp}$ defines an isomorphism

$$\perp : {}^*\mathbb{Z}(\theta) \longrightarrow {}^*\mathbb{Z}(\theta^{-1}) \tag{8}$$

for all $\theta \neq 0$.

Proposition 8. *Let $\theta \neq 0$. Then the duality isomorphism (8) respects the fine growth-decay tri-filtration:*

$${}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta)^{\perp} = {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta^{-1})$$

Proof. Note that

$${}^*n \cdot \mu < \iota \Leftrightarrow {}^*n^{\perp} \cdot \mu < \iota \tag{9}$$

which implies that duality respects the fine growth bi-filtration. On the other hand, $\varepsilon({}^*n^{\perp}) = -\theta^{-1}\varepsilon({}^*n)$ so the decay filtration is preserved as well. □

Recall [19] that θ is **projective linear equivalent** to η if there exists $A \in \mathrm{PGL}_2(\mathbb{Z})$ such that $A(\theta) = \eta$. The relation of projective linear equivalence is denoted here

$$\theta \mathrel{\overset{\frown}{\smile}} \eta.$$

Theorem 3. *If $\theta \mathrel{\overset{\frown}{\smile}} \eta$ by $A \in \mathrm{PGL}_2(\mathbb{Z})$, then A induces an isomorphism of ${}^*\mathbb{Z}(\theta)$ with ${}^*\mathbb{Z}(\eta)$ preserving the fine growth-decay tri-filtration.*

Proof. The isomorphism is induced by the matrix action of a linear representative $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ on pairs $({}^*n^{\perp}, {}^*n)$ where ${}^*n \in {}^*\mathbb{Z}(\theta)$ and $\theta \cdot {}^*n \simeq {}^*n^{\perp}$. That is, $A({}^*n) = c{}^*n^{\perp} + d{}^*n$ and $A({}^*n^{\perp}) = a{}^*n^{\perp} + b{}^*n$. By (9), ${}^*n \in {}^*\mathbb{Z}^{\mu[\iota]} \Leftrightarrow {}^*n^{\perp} \in {}^*\mathbb{Z}^{\mu[\iota]}$. It follows then that ${}^*n \in {}^*\mathbb{Z}^{\mu[\iota]} \Leftrightarrow A({}^*n) \in {}^*\mathbb{Z}^{\mu[\iota]}$. On the other hand,

$$\begin{aligned}
\eta \cdot A({}^*n) - A({}^*n^{\perp}) &= \frac{1}{c\theta + d}\Big\{(a\theta + b)\big(c{}^*n^{\perp} + d{}^*n\big) - (c\theta + d)\big(a{}^*n^{\perp} + b{}^*n\big)\Big\} \\
&= \frac{1}{c\theta + d}\big(\theta{}^*n - {}^*n^{\perp}\big) \\
&= \frac{\varepsilon({}^*n)}{c\theta + d}.
\end{aligned}$$

Therefore: ${}^*n \in {}^*\mathbb{Z}_{\nu}(\theta) \Leftrightarrow A({}^*n) \in {}^*\mathbb{Z}_{\nu}(\eta)$. □

3. Nonvanishing Spectra

The nontriviality of the group ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ for specific indices $\mu, \nu \in {}^*\mathbb{PR}_{\varepsilon}$ depends intimately on the type of θ. We define the **nonvanishing spectrum** to be the subset

$$\mathrm{Spec}(\theta) = \{(\mu,\nu) |\ {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \neq 0\} \subset {}^*\mathbb{PR}^2_{\varepsilon}.$$

In this section, we will characterize the spectra of a real number according to its "linear classification" (rational, badly approximable, well approximable, etc.). We begin with some very general results.

Proposition 9. *If* $\theta \Bumpeq \eta$ *then* $\mathrm{Spec}(\theta) = \mathrm{Spec}(\eta)$.

Proof. This follows immediately from Theorem 3. ☐

Theorem 4. *For all* $\theta \in \mathbb{R}$ *and* $\mu < \nu$, ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \neq 0$.

Proof. By Proposition 2, we may find ρ with $\mu < \rho < \nu$; and by Proposition 1, $\rho = \langle {}^*N^{-1} \rangle$ for some ${}^*N \in {}^*\mathbb{Z}_+ - \mathbb{Z}_+$. By the Uniform Dirichlet Theorem[1] there is ${}^*n \in {}^*\mathbb{Z}(\theta)$ such that $|\varepsilon({}^*n)| < {}^*N^{-1}$ where ${}^*n < {}^*N$. Therefore, $|\nu({}^*n)| \leq \nu$. On the other hand ${}^*n \cdot \mu \leq {}^*N \cdot \mu \in {}^*\mathbb{PR}_{\varepsilon}$ since $\mu < \rho$, so ${}^*n \in {}^*\mathbb{Z}^{\mu}(\theta)$ ☐

The set $\{(\mu,\nu) | \mu < \nu\} \subset \mathrm{Spec}(\theta)$ is called the **slow component**.

For $\theta \in \mathbb{R}$, denote by $\{a_i = a_i(\theta)\}$, $i = 0, 1, \ldots$, the sequence of its partial quotients [19]: an infinite sequence $\Leftrightarrow$ $\theta \notin \mathbb{Q}$. As is the custom, we write

$$\theta = [a_0 a_1 \ldots].$$

The sequence $\{q_i\}$ of **best denominators** of θ is defined recursively by the formula

$$q_{i+1} = a_{i+1}q_i + q_{i-1}, \quad q_0 = 1,\ q_1 = a_1.$$

Similarly, the sequence $\{p_i\}$ of **best numerators** is defined

$$p_{i+1} = a_{i+1}p_i + p_{i-1}, \quad p_0 = a_0,\ p_1 = a_1 a_0 + 1.$$

We have (e.g. see Theorem 5 of [19])

$$q_i|q_i\theta - p_i| < q_i^{-1}. \tag{10}$$

The sequence of quotients $\{p_i/q_i\}$ is called the sequence of **best approximations** (or **principal convergents**) of θ: by (10) they satisfy $p_i/q_i \to \theta$. See [6], [19], [22].

Consider now a sequence $\{q_{n_i}\}$ in which q_{n_i} is the n_ith best denominator of θ, where $n_i \leq n_{i+1}$ for all i and $n_i \to \infty$. By (10) the associated sequence class defines an element

$${}^*\widehat{q} := {}^*\{q_{n_i}\} \in {}^*\mathbb{Z}(\theta)$$

called a **best denominator class**, and the classes

$$\widehat{\mu} := \mu({}^*\widehat{q}), \quad \text{resp.} \quad \widehat{\nu} := \nu({}^*\widehat{q})$$

will be referred to as the associated **best growth** resp. **best decay** of ${}^*\widehat{q}$. We will denote by ${}^*\widehat{q}^{+}$ resp. ${}^*\widehat{q}^{-}$ the classes of the successor and predecessor sequences ${}^*\{q_{n_i+1}\}$ resp. ${}^*\{q_{n_i-1}\}$, with a similar notation employed for the associated best growth and best decay classes e.g. $\widehat{\mu}^{+}$ = the growth class of ${}^*\widehat{q}^{+}$. Note that $\widehat{\mu}^{+} \leq \mu$ and $\widehat{\nu}^{+} \leq \nu$. The above terminology applies without change to the corresponding sequence of best numerators $\{p_{n_i}\}$, yielding the associated best numerator class ${}^*\widehat{p}$, etc.

---

[1] For any real number $N > 1$, there exist $p, q \in \mathbb{Z}$ with $1 \leq q < N$ such that $|q\theta - p| < 1/N$. See [25].

Proposition 10. *Let* ${}^*\widehat{q}$ *be a best denominator class,* ${}^*\widehat{p}$ *the corresponding best numerator class. Then* ${}^*\widehat{q}^{\perp} = {}^*\widehat{p} \in {}^*\mathbb{Z}(\theta^{-1})$*. In particular, the best growth* $\widehat{\mu}$ *and best decay* $\widehat{\nu}$ *of* ${}^*\widehat{q}$ *are also the best growth and best decay of* ${}^*\widehat{p}$*.*

Proof. That ${}^*\widehat{q}^{\perp} = {}^*\widehat{p}$ follows from (10). Since ${}^*\widehat{q}\theta - {}^*\widehat{p} = \varepsilon({}^*\widehat{q})$, the best growth and best decay classes of ${}^*\widehat{p}$ coincide with those of ${}^*\widehat{q}$. □

Note 2. When $\theta \in \mathbb{Q}$, the sequence of best approximations is finite, so every best approximation class ${}^*\widehat{q}$ is standard and equal to one of the q_i. In this case, every best growth is $\widehat{\mu} = 1$ and $\widehat{\nu} = -\infty$ or 1.

For $\theta \in \mathbb{R} - \mathbb{Q}$, we denote by:

- ${}^*\mathbb{Z}_{\mathrm{b}}(\theta)$ the set of best denominator classes.
- ${}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta)$ (${}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bd}}(\theta)$) the set of best growths (best decays) of best denominator classes.

Proposition 11. *For* $\theta \in \mathbb{R} - \mathbb{Q}$*,* ${}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta)$ *is closed in the order topology.*

Proof. If ${}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta) = {}^*\mathbb{PR}_{\varepsilon}$ we are done, so suppose otherwise. Given $\mu \in {}^*\mathbb{PR}_{\varepsilon} - {}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta)$, we will construct an interval $(\mu', \mu'') \ni \mu$ containing no elements of ${}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta)$. Let ${}^*x \in \mu^{-1}$. Then there exists a largest ${}^*\widehat{q}$ for which ${}^*x > {}^*\widehat{q}$: indeed, if we choose $\{x_i\} \in {}^*x$ non-decreasing and let q_{n_i} be the largest member of $\{q_i\}$ which is less than x_i, then ${}^*\widehat{q} = {}^*\{q_{n_i}\}$ works. Since $\mu \notin {}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta)$, there exists ${}^*r$ infinite with ${}^*r \cdot {}^*\widehat{q} = {}^*x$. Now let ${}^*s \in {}^*\mathbb{R}_{+}$ be such that both ${}^*s$ and ${}^*r/{}^*s$ are infinite, and let ${}^*y = ({}^*r/{}^*s) \cdot {}^*\widehat{q}$. If we denote by μ' the class of ${}^*y^{-1}$ then $\widehat{\mu} > \mu' > \mu$ and $[\mu, \mu'] \cap {}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta) = \varnothing$. In the same way, we may produce $\mu'' < \mu$ with $[\mu'', \mu] \cap {}^*\mathbb{PR}_{\varepsilon}^{\mathrm{bg}}(\theta) = \varnothing$. Thus (μ'', μ') is the sought after interval. □

The following result is our first vanishing theorem: a straightforward reinterpretation of the quality of being a best denominator class in terms of the growth-decay bi-filtration.

Theorem 5. *Let* $\theta \in \mathbb{R} - \mathbb{Q}$ *and let* ${}^*\widehat{q}$ *be any best denominator class with associated growth and decay* $\widehat{\mu}, \widehat{\nu}$*. Then for all* $\mu \geq \widehat{\mu}$ *and* $\nu < \widehat{\nu}$*,* ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) = 0$*.*

Proof. For $\mu \geq \widehat{\mu}$ and $\nu < \widehat{\nu}$, suppose there exists a non-zero ${}^*n \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$, which we may assume is positive. Then ${}^*n \cdot \widehat{\mu} \leq {}^*n \cdot \mu \in {}^*\mathbb{PR}_{\varepsilon}$ implies that ${}^*n < {}^*\widehat{q}$. In turn, the latter implies, since ${}^*\widehat{q}$ is the class of a non decreasing sequence of best denominators of θ, that

$$|\varepsilon({}^*n)| = |\theta{}^*n - {}^*n^{\perp}| \geq |\theta{}^*\widehat{q} - {}^*\widehat{q}^{\perp}| = |\varepsilon({}^*\widehat{q})|.$$

From this we derive $\nu({}^*n) \geq \widehat{\nu} > \nu$, contradiction. □

In the (μ, ν)-plane the coordinates belonging to the right-infinite horizontal strip

$$\widehat{R} = \{(\mu, \nu) |\ \mu \geq \widehat{\mu},\ \nu < \widehat{\nu}\}$$

give parameters where the groups ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ vanish. We call $\widehat{R}$ a **vanishing strip**. See the graph labeled "generic irrational" in Figure 1.

We now characterize the spectra of some well-known classes of real numbers.

Proposition 12. $\theta \in \mathbb{Q} \Leftrightarrow \mathrm{Spec}(\theta) = {}^*\mathbb{PR}^2_{\varepsilon}$.

Proof. If $\theta \in \mathbb{Q}$ then for all ν, ${}^*\mathbb{Z}_{\nu}(\theta) = {}^*\mathbb{Z}_{-\infty}(\theta) = {}^*\mathbb{Z}(\theta)$, so ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) = {}^*\mathbb{Z}^{\mu}(\theta) \neq 0$ for all μ, ν. On the other hand, if $\theta \in \mathbb{R} - \mathbb{Q}$ then by Theorem 5, $\mathrm{Spec}(\theta) \subsetneq {}^*\mathbb{PR}^2_{\varepsilon}$. □



rational

generic irrational



badly approximable

well but not very well approximable



very well approximable (exponent κ)

Liouville

Figure 1. Portraits of Spectra. Shaded regions and heavy lines represent nonvanishing.

Recall that $\theta \in \mathbb{R} - \mathbb{Q}$ is **badly approximable** if

$$\lim_{n\to\infty} \inf \ n\|n\theta\| > 0,$$

where $\|\cdot\|$ is the distance-to-the-nearest-integer function. Or equivalently, if there exists a real number $C > 0$ such that for all $0 \neq {}^*n \in {}^*\mathbb{Z}(\theta)$,

$$|{}^*n| \cdot |\varepsilon({}^*n)| \geq C.$$

The set $\mathfrak{B}$ of badly approximable numbers has cardinality the continuum [22].

Theorem 6. *The following statements are equivalent:*

- i. $\theta \in \mathfrak{B}$.
- ii. ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) = 0$ *for all* $\mu \geq \nu$.
- iii. ${}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta) = {}^*\mathbb{PR}_{\varepsilon}$. *In particular, for all* $\widehat{\mu} \in {}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta)$, $\widehat{\mu}^{+} = \widehat{\mu}$.
- iv. $\widehat{\mu} = \widehat{\nu}$ *for every best growth decay pair: that is* ${}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta) = {}^*\mathbb{PR}^{\mathrm{bd}}_{\varepsilon}(\theta)$.

Proof. i. $\Rightarrow$ ii. If $\theta \in \mathfrak{B}$ then for all non-zero ${}^*n \in {}^*\mathbb{Z}(\theta)$ we have $|{}^*n| \cdot |\varepsilon({}^*n)| \geq C$. If there exists $\mu \geq \nu$ with $0 \neq {}^*n \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ then in ${}^*\mathbb{PR}_{\varepsilon}$,

$$|{}^*n| \cdot \mu \geq |{}^*n| \cdot \nu \geq |{}^*n| \cdot \nu({}^*n) \geq 1 = \text{ the } {}^*\mathbb{PR}\text{-class of } C,$$

implying that $|{}^*n| \cdot \mu \notin {}^*\mathbb{PR}_{\varepsilon}$ and ${}^*n \notin {}^*\mathbb{Z}^{\mu}$. ii. $\Rightarrow$ i. If ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) = 0$ for all $\mu \geq \nu$, then for each ${}^*n \in {}^*\mathbb{Z}(\theta)$, $|{}^*n| \cdot \varepsilon({}^*n) \geq \delta > 0$ where $\delta \in \mathbb{R}$. We can choose delta uniformly: if not, then by a diagonal sequence argument we could produce an element ${}^*\mathbb{Z}(\theta)$ for which $|{}^*n| \cdot \varepsilon({}^*n)$ is infinitesimal (i.e. we could produce a non trivial element of ${}^*\mathbb{Z}^{\nu({}^*n)}_{\nu({}^*n)}(\theta)$) violating the hypothesis. i. $\Rightarrow$ iii. $\theta \in \mathfrak{B} \Leftrightarrow$ the partial quotients a_i are uniformly bounded $\Leftrightarrow$ the successive ratios of best denominators q_{i+1}/q_i are uniformly bounded. Now given $\mu \in {}^*\mathbb{PR}_{\varepsilon}$ let $\{n_i\} \subset \mathbb{N}_{+}$ represent μ^{-1}. For each i let q_{k_i} be the largest best denominator with $q_{k_i} \leq n_i$ so that $n_i < q_{k_i+1}$. By hypothesis there exists a constant $B > 1$ so that $q_{k_i+1} < Bq_{k_i}$. It follows that $n_i = b_i q_{k_i}$ with $1 \leq b_i < B$. Then the growth of the class ${}^*\widehat{q}$ is equal to μ. iii. $\Rightarrow$ i. If $\theta \notin \mathfrak{B}$, choose $\widehat{\mu}$ so that if $\{q_{n_i}\}$ represents $\widehat{\mu}^{-1}$ then q_{n_i+1}/q_{n_i} is monotone and unbounded. Then the successor sequence $\{q^{+}_{n_i} = q_{n_i+1}\}$ defines a distinct element $\widehat{\mu}^{+} \in {}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta)$ with $\widehat{\mu}^{+} < \widehat{\mu}$. It follows that ${}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta) \neq {}^*\mathbb{PR}_{\varepsilon}$. i. $\Leftrightarrow$ iv. From Dirichlet's Theorem and the definition of $\mathfrak{B}$, $\widehat{\mu} = \widehat{\nu}$ for every best growth decay pair $\Leftrightarrow$ $1 > |{}^*\widehat{q}| \cdot \varepsilon({}^*\widehat{q}) \geq C$ for every best denominator class for some uniform $C > 0 \Leftrightarrow \theta \in \mathfrak{B}$. ☐

Recall that $\theta \in \mathbb{R} - \mathbb{Q}$ which is not badly approximable is called **well approximable**: that is, $\lim_{n\to\infty} \inf \ n\|n\theta\| = 0$. We denote the set of well approximable numbers by

$$\mathfrak{W} = \mathbb{R} - (\mathbb{Q} \cup \mathfrak{B}).$$

Theorem 7. *Let* $\theta \in \mathbb{R} - \mathbb{Q}$. *The following statements are equivalent:*

- i. $\theta \in \mathfrak{W}$.
- ii. *There exists* $\mu \in {}^*\mathbb{PR}_{\varepsilon}$ *such that* ${}^*\mathbb{Z}^{\mu}_{\mu}(\theta) \neq 0$.
- iii. *There exists* $\widehat{\mu} \in {}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta)$ *such that* $\widehat{\mu}^{+} < \widehat{\mu}$. *In particular,* ${}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}(\theta)$ *is not a dense order.*
- iv. $\widehat{\mu} > \widehat{\nu}$ *for some best growth best decay pair.*

Proof. i. $\Leftrightarrow$ ii. By Theorem 6, $\theta \in \mathfrak{B} \Rightarrow {}^*\mathbb{Z}^{\mu}_{\mu}(\theta) = 0$ for all μ. On the other hand if $\theta \in \mathfrak{W} \Rightarrow$ there exists $\mu \geq \nu$ with ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \neq 0$. By the order-reversing property of the growth filtration, the latter implies that ${}^*\mathbb{Z}^{\nu}_{\nu}(\theta) \neq 0$. i. $\Leftrightarrow$ iii. If $\theta \in \mathfrak{W}$ then there exists a best denominator sequence $\{q_{n_i}\}$ for which the sucessor $\{q^+_{n_i} = q_{n_i+1}\}$ satisfies $q^+_{n_i}/q_{n_i} \to \infty$. The other direction follows from Theorem 6. iii. $\Leftrightarrow$ iv. Immediate from Theorem 6. $\square$

Let $\kappa > 1$. Recall [4] that θ is **very well approximable (of exponent κ)** if the set of $n \in \mathbb{N}$ for which $\|n\theta\| < n^{-\kappa}$ has infinite cardinality i.e.

$$\liminf_{n\to\infty} n^{\lambda}\|n\theta\| = 0 \tag{11}$$

for all $1 \leq \lambda < \kappa$. We denote the set of very well approximable numbers of exponent κ by $\mathfrak{W}_\kappa$. If θ is very well approximable for some exponent then we will say simply that θ is **very well approximable**; the set of such numbers is denoted $\mathfrak{W}_{>1} = \bigcup \mathfrak{W}_\kappa$. The inclusion $\mathfrak{W}_{>1} \subset \mathfrak{W}$ is proper and we write $\mathfrak{W}_1 = \mathfrak{W} - \mathfrak{W}_{>1}$ = the set of **well but not very well approximable numbers**.

Theorem 8. *Let $\theta \in \mathbb{R} - \mathbb{Q}$. The following statements are equivalent:*

i. $\theta \in \mathfrak{W}_\kappa$, $\kappa > 1$.

ii. *There exists $\mu \in {}^*\mathbb{PR}_\varepsilon$ such that*

$$\bigcap_{\lambda\in[1,\kappa)} {}^*\mathbb{Z}^{\mu}_{\mu^\lambda}(\theta) \neq 0.$$

In particular, ${}^*\mathbb{Z}^{\nu}_{\nu}(\theta) \neq 0$ for all $\mu^\kappa < \nu \leq \mu$.

iii. $\widehat{\mu}^\kappa > \widehat{\nu}$ *for some best growth decay pair.*

Proof. i. $\Rightarrow$ ii. $\theta \in \mathfrak{W}_\kappa$ for $\kappa \Leftrightarrow$ there exists ${}^*n \in {}^*\mathbb{Z}(\theta)$ such that $\nu({}^*n) < \mu({}^*n)^\lambda$ for all $1 \leq \lambda < \kappa$. Since ${}^*\mathbb{PR}_\varepsilon$ is a dense linear order, we may find $\mu \in {}^*\mathbb{PR}_\varepsilon$ such that $\nu({}^*n) < \mu^\lambda < \mu({}^*n)^\lambda$ for all $1 \leq \lambda < \kappa$. Then $\mu < \mu({}^*n)$ and therefore

$${}^*n \in {}^*\mathbb{Z}^{\mu}_{\nu({}^*n)}(\theta) \subset {}^*\mathbb{Z}^{\mu}_{\mu^\lambda}(\theta)$$

for all $1 \leq \lambda < \kappa$. ii. $\Rightarrow$ i. Let ${}^*n \in \bigcap_{\lambda\in[1,\kappa)} {}^*\mathbb{Z}^{\mu}_{\mu^\lambda}(\theta)$. Then ${}^*n \cdot \mu \in {}^*\mathbb{PR}_\varepsilon$ and $\nu({}^*n) < \mu^\lambda$ for all $1 \leq \lambda < \kappa$. It follows that

$${}^*n^\lambda \cdot \nu({}^*n) < ({}^*n \cdot \mu)^\lambda \in {}^*\mathbb{PR}_\varepsilon$$

which implies (11). i. $\Leftrightarrow$ iii. $\theta \in \mathfrak{W}_\kappa$, $\kappa > 1 \Leftrightarrow$ (11) holds for all $1 \leq \lambda < \kappa \Leftrightarrow$ for some best denominator sequence ${}^*\widehat{q}$, (11) holds for all $1 \leq \lambda < \kappa$. $\square$

The **Liouville numbers** are those which are very well approximable for every exponent $\kappa > 1$; they are denoted $\mathfrak{W}_\infty$. For any $\mu \in {}^*\mathbb{PR}_\varepsilon$, write $(\mu^\infty, \mu] = \bigcup_{\kappa>1}(\mu^\kappa, \mu]$. The next result follows immediately from Theorem 8: we recall from §1 that $\bar{\mu}$ is the orbit of μ with respect to the Frobenius action of $({}^*\mathbb{R}_{\mathrm{fin}})^\times_+$. For $\widehat{\mu}, \widehat{\nu}$ a best growth-decay pair, the corresponding Frobenius orbits are denoted $\bar{\widehat{\mu}}, \bar{\widehat{\nu}}$.

Corollary 1. *Let $\theta \in \mathbb{R} - \mathbb{Q}$. The following statements are equivalent:*

i. $\theta \in \mathfrak{W}_\infty$.

ii. *There exists $\mu \in {}^*\mathbb{PR}_\varepsilon$ such that*

$$\bigcap_{\lambda\in[1,\infty)} {}^*\mathbb{Z}^{\mu}_{\mu^\lambda}(\theta) \neq 0.$$

In particular, ${}^*\mathbb{Z}^{\nu}_{\nu}(\theta) \neq 0$ for all $\nu \in (\mu^\infty, \mu]$.

iii. $\bar{\widehat{\mu}} > \bar{\widehat{\nu}}$ *for some best growth decay pair.*

4. Flat Spectra

The line $\mu = \nu$ represents a critical divide whose intersection with $\mathrm{Spec}(\theta)$ gives a new invariant of θ which is strongly influenced by patterns found in the sequence of partial quotients; as opposed to the full spectrum which is essentially determined by the exponent.

We define the **flat spectrum** of θ to be the set

$$\mathrm{Spec}_{\mathrm{flat}}(\theta) = \left\{ \mu \in {}^*\mathbb{PR}_{\varepsilon} \mid {}^*\mathbb{Z}^{\mu}_{\mu}(\theta) \neq 0 \right\}.$$

By Proposition 12, $\mathrm{Spec}_{\mathrm{flat}}(\theta) = {}^*\mathbb{PR}_{\varepsilon}$ for all $\theta \in \mathbb{Q}$, and by Theorem 6, $\mathrm{Spec}_{\mathrm{flat}}(\theta) = \emptyset$ for all $\theta \in \mathfrak{B}$. Therefore we will restrict attention in this section to $\theta \in \mathfrak{W}$ = the set of well approximable numbers.

Suppose that ${}^*m \in {}^*\mathbb{Z}(\theta)$ can be factored

$$ {}^*m = {}^*x \cdot {}^*n $$

for ${}^*x \in {}^*\mathbb{Z}$ and ${}^*n \in {}^*\mathbb{Z}(\theta)$. If $\nu = \nu({}^*n)$ and ${}^*x \in {}^*\mathbb{Z}^{\nu}$ then it follows that

$$ \varepsilon({}^*m) = {}^*x \cdot \varepsilon({}^*n), \quad \text{i.e.} \quad \nu({}^*m) = {}^*x \cdot \nu. $$

In this case we refer to ${}^*m = {}^*x \cdot {}^*n$ as an **ideological factorization** and speak of ${}^*m$ as being an **ideological multiple** of ${}^*n$. Note that in this case ${}^*m^{\perp} = {}^*x \cdot {}^*n^{\perp}$. Conversely, the action

$$ {}^*\mathbb{Z}^{\nu} \times {}^*\mathbb{Z}_{\nu}(\theta) \to {}^*\mathbb{Z}(\theta), \quad ({}^*x, {}^*n) \mapsto {}^*x \cdot {}^*n, $$

has image consisting of ideological multiples. There may be factorizations of ${}^*m$ which are not ideological.

Call ${}^*m \in {}^*\mathbb{Z}(\theta)$ a **multibest denominator** if there is an ideological factorization

$$ {}^*m = {}^*x \cdot {}^*\widehat{q} $$

for some best denominator ${}^*\widehat{q}$. We have

$$ \widehat{\nu} \leq \nu({}^*m) = {}^*x \cdot \widehat{\nu}, \quad \mu({}^*m) = {}^*x^{-1} \cdot \widehat{\mu} \leq \widehat{\mu}. \tag{12} $$

Recall that if ${}^*\widehat{q}$ is the class of $\{q_{n_i}\}$, we denote by ${}^*\widehat{q}^{+}$ the successor class ${}^*\{q_{n_i+1}\}$ and by ${}^*\widehat{q}^{-}$ the predecessor class ${}^*\{q_{n_i-1}\}$. Then if ${}^*a$ denotes the class of $\{a_{n_i+1}\}$ (the corresponding sequence of partial quotients), we have

$$ {}^*\widehat{q}^{+} = {}^*a \cdot {}^*\widehat{q} + {}^*\widehat{q}^{-}. $$

We say that ${}^*\widehat{q}$ has **infinite partial quotient** if ${}^*a$ is infinite. Note that ${}^*\widehat{q}$ has infinite partial quotient $\Leftrightarrow$ $\widehat{\mu}^{+} < \widehat{\mu}$. We apply the same terminology to a multibest class ${}^*m = {}^*x \cdot {}^*\widehat{q}$ if ${}^*\widehat{q}$ has infinite partial quotient. An element $\theta \in \mathfrak{W}$ will possess best classes ${}^*\widehat{q}$ having *finite* partial quotient precisely when the sequence $\{a_i\}$ of partial quotients has an infinite bounded subsequence e.g. when $\theta = e = [2, 1, 2, 1, 1, 4, 1, 1, 6, \dots]$.

Theorem 9. *Let* $\theta \in \mathfrak{W}$. *Then*

$$ {}^*\mathbb{Z}^{\mu}_{\mu}(\theta) = \left\{ {}^*m \text{ multibest with infinite partial quotient} \;\middle|\; \nu({}^*m) \leq \mu < \mu({}^*m) \right\}. \tag{13} $$

Proof. We may assume without loss of generality that $\theta > 0$. Clearly the set on the right hand side of (13) is contained in ${}^*\mathbb{Z}^{\mu}_{\mu}(\theta)$. Suppose now that $0 \neq {}^*m \in {}^*\mathbb{Z}^{\mu}_{\mu}(\theta)$.

Then we may find monotone representatives $\{s_i\}$ of μ and $\{m_i\}$ of ${}^*m$ for which $m_is_i \to 0$ and $\|m_i\theta\| \leq s_i$. From the first fact we get $s_i < m_i^{-1}$ so that we may write

$$\|m_i\theta\| \leq s_i < m_i^{-1}. \tag{14}$$

In particular, if $\{m_i^{\perp}\}$ is a representative of the dual ${}^*m^{\perp}$ then $|\theta - (m_i^{\perp}/m_i)| < m_i^{-2}$. By Grace's Theorem (Theorem 10 of [19]), the $m_i^{\perp}/m_i$ are (intermediate) convergents of θ: that is, for each i there exists $n = n_i$, $r = r_i$ nonnegative integers with $m_i/m_i^{\perp} = p_{n,r}/q_{n,r}$, where

$$p_{n,r} := rp_{n+1} + p_n, \quad q_{n,r} := rq_{n+1} + q_n,$$

$\{p_i\}$, $\{q_i\}$ are the sequences of numerators and denominators of the principal convergents (best approximations) of θ and $0 \leq r \leq a_{n+2}-1$. Moreover, Grace's Theorem further affirms that the possible values of r in this case are $r = 0, 1$ or $a_{n+2} - 1$. If we denote by ${}^*\widehat{q}$ the class of $\{q_{n_i}\}$, ${}^*\widehat{q}^{+}$ the class of $\{q_{n_i+1}\}$ and by ${}^*r$ the class of $\{r_i\}$ then we may write

$${}^*m = {}^*x \cdot {}^*\widehat{q}_{{}^*r} := {}^*x \cdot ({}^*r{}^*\widehat{q}^{+} + {}^*\widehat{q}), \quad {}^*m^{\perp} = {}^*x \cdot {}^*\widehat{p}_{{}^*r} := {}^*x \cdot ({}^*r{}^*\widehat{p}^{+} + {}^*\widehat{p}).$$

Note that the factorization ${}^*m = {}^*x \cdot {}^*\widehat{q}_{{}^*r}$ is ideological: for

$$\varepsilon({}^*m) = {}^*m\theta - {}^*m^{\perp} = {}^*x \cdot ({}^*\widehat{q}_{{}^*r}\theta - {}^*\widehat{p}_{{}^*r}) = {}^*x \cdot \varepsilon({}^*\widehat{q}_{{}^*r})$$

implying (since $\varepsilon({}^*m) \in {}^*\mathbb{R}_{\varepsilon}$) that ${}^*x \in {}^*\mathbb{Z}^{\nu}$ for $\nu = \nu({}^*\widehat{q}_{{}^*r})$.

We now show that ${}^*m$ is multibest having infinite partial quotient i.e. that ${}^*r = 0$ and ${}^*\widehat{q}_{{}^*r} = {}^*\widehat{q}$ has infinite partial quotient. To do this we will make use of a closed expression for the error term $\varepsilon({}^*\widehat{q}_{{}^*r})$ of the convergent ${}^*\widehat{q}_{{}^*r}$. First, let ${}^*a^{+}$ be the sequence class of $\{a_{n_i+2}\}$; thus the possibilities afforded by Grace's Theorem are ${}^*r = 0, 1, {}^*a^{+} - 1$. Now for any n we define θ_n by the formula $\theta = [a_1 \dots a_{n-1}\theta_n]$. In particular, $\theta_n = [a_na_{n+1}\dots] = a_n + \theta_{n+1}^{-1}$. Let ${}^*\theta$ be the class of $\{\theta_{n_i+2}\}$ and note that ${}^*\theta = {}^*a^{+} + ({}^*\theta^{+})^{-1}$ where ${}^*\theta^{+}$ is the class of $\{\theta_{n_i+3}\}$. Then the Lemma of §I.4 of [19] yields

$$|\varepsilon({}^*\widehat{q}_{{}^*r})| = \frac{{}^*\theta - {}^*r}{{}^*\theta{}^*\widehat{q}^{+} + {}^*\widehat{q}}.$$

and therefore $|\varepsilon({}^*m)| = {}^*x \cdot ({}^*\theta - {}^*r)/({}^*\theta{}^*\widehat{q}^{+} + {}^*\widehat{q})$.

$\boxed{\textit{Case } 1}$ ${}^*r = 0$. Since $\nu({}^*m) \leq \mu < \mu({}^*m)$, the multibest inequalities (12) imply that $\widehat{\nu} \leq \mu < \widehat{\mu}$ and hence $\widehat{\nu} < \widehat{\mu}$. Since ${}^*\theta \geq 1$, we have

$$\widehat{\nu} = \langle {}^*\widehat{q}^{+} + ({}^*\widehat{q}/{}^*\theta)\rangle^{-1} = \langle {}^*\widehat{q}^{+}\rangle^{-1} = \widehat{\mu}^{+}.$$

In particular, $\widehat{\mu}^{+} < \widehat{\mu}$ and therefore ${}^*\widehat{q}$ has infinite partial quotient. $\mu({}^*m) = {}^*x \cdot \langle {}^*\widehat{q}^{-1}\rangle = {}^*x \cdot \widehat{\mu}$. In other words, this case comprises precisely the multibest denominators ${}^*m$ having infinite partial quotient, for which $\nu({}^*m) \leq \mu < \mu({}^*m)$.

$\boxed{\textit{Case } 2}$ ${}^*r = 1$. Then $\nu({}^*\widehat{q}_{{}^*r}) = \mu({}^*\widehat{q}_{{}^*r}) = \widehat{\mu}^{+}$, so it is impossible that $\nu({}^*m) < \mu({}^*m)$, in view of the ideological factorization ${}^*m = {}^*x \cdot {}^*\widehat{q}_{{}^*r}$. Indeed, $\nu({}^*m) < \mu({}^*m)$ implies that ${}^*x \cdot \widehat{\mu}^{+} < {}^*x^{-1} \cdot \widehat{\mu}^{+}$ or $\langle {}^*x^2\rangle < 1$, impossible since ${}^*x$, being a sequence class of integers, is not infinitesimal.

$\boxed{\textit{Case } 3}$ ${}^*r = {}^*a^{+} - 1$. Then ${}^*\theta - {}^*r = 1 + ({}^*\theta^{+})^{-1} > 1$ and is bounded from above. Thus $\nu({}^*\widehat{q}_{{}^*r}) = \mu({}^*\widehat{q}_{{}^*r}) = \langle {}^*a^{+}\widehat{q}^{+}\rangle^{-1} = ({}^*a^{+})^{-1} \cdot \widehat{\mu}^{+}$, so as in $\boxed{\textit{Case } 2}$ we cannot have $\nu({}^*m) < \mu({}^*m)$. $\square$

Let ${}^*\mathbb{Z}^{\infty}_{\mathrm{b}}(\theta)$ be the set of best denominators having infinite partial quotient and let ${}^*\mathbb{Z}^{\infty}_{\mathrm{mb}}(\theta) \supset {}^*\mathbb{Z}^{\infty}_{\mathrm{b}}(\theta)$ be the set of multibest denominators having infinite partial quotient.

Corollary 2. *Let* $\theta \in \mathfrak{W}$. *Then*

$$\mathrm{Spec}_{\mathrm{flat}}(\theta) = \bigcup_{{}^*m\in {}^*\mathbb{Z}^{\infty}_{\mathrm{mb}}(\theta)} \big[\nu({}^*m), \mu({}^*m)\big) = \bigcup_{{}^*\widehat{q}\in {}^*\mathbb{Z}^{\infty}_{\mathrm{b}}(\theta)} \big[\widehat{\nu}, \widehat{\mu}\big). \tag{15}$$

In particular, $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ *has interior.*

Proof. The first equality follows immediately from Theorem 9. If ${}^*m = {}^*x \cdot {}^*\widehat{q}$ then

$$\big[\nu({}^*m), \mu({}^*m)\big) \subset \big[\widehat{\nu}, \widehat{\mu}\big)$$

giving the second equality. □

An interval $\big[\widehat{\nu}, \widehat{\mu}\big)$ appearing in (15) is called a **best interval**. These have been indicated in the portrait of "generic irrational" found in Figure 1.

Note 3. As specified in Theorem 8 and Corollary 1, for $\theta \in \mathfrak{W}_{>1}$, $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ contains power intervals of the form $\big[\widehat{\mu}^{\lambda}, \widehat{\mu}\big)$, $\lambda > 1$, and for $\theta \in \mathfrak{W}_{\infty}$, $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ contains Frobenius rays $\Phi_{(1,\infty)}(\widehat{\mu})$. If $\theta \in \mathfrak{W}_1$, then Corollary 2 says that although $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ contains no intervals of the form $\big[\widehat{\mu}^{\lambda}, \widehat{\mu}\big)$, it is nevertheless open. We can summarize this trichotomy by saying that the flat spectrum of elements of θ in $\mathfrak{W}_1$, $\mathfrak{W}_{>1}$ or $\mathfrak{W}_{\infty}$ contains components having *subpolynomial*, *polynomial* or *exponential connectivity*, respectively.

We will show that for $\theta \in \mathbb{R} - \mathbb{Q}$, $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ is a proper subset of ${}^*\mathbb{PR}_{\varepsilon}$. Let $\mu \in {}^*\mathbb{PR}_{\varepsilon}$, which we assume can be represented by a sequence $\{s_i\}$ of positive reals monotonically tending to 0. We say that μ is **shift invariant** if we may choose $\{s_i\}$ so that there exists $M > 0$ with $s_i/s_{i+1} < M$ for all i i.e. $s_{i+1} < s_i < M s_{i+1}$ for all i. If $\{s'_i\}$ is another monotone representative sequence representing μ then there exists a constant C for which ${}^*s - C{}^*s' \simeq 0$. Thus any two monotone representative sequences will have this property, and therefore being shift invariant is independent of the selected representative sequence. If μ is shift invariant and $\{s_i\}$ is a monotone representative then $\mathrm{s}(\mu) := \langle {}^*\{s_{i+1}\}\rangle = \mu$, hence the terminology. Let ${}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ be the set of shift invariant elements of ${}^*\mathbb{PR}_{\varepsilon}$. In what follows X^{C} denotes the complement of a set X.

Proposition 13. ${}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ *is clopen in* ${}^*\mathbb{PR}_{\varepsilon}$.

Proof. Suppose that $\mu \notin {}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$, represent it by ${}^*s = {}^*\{s_i\}$ with $s_i/s_{i+1} \to \infty$. Let $0 < {}^*\varepsilon \in {}^*\mathbb{R}_{\varepsilon}$ be such that $\varepsilon_i s_i/s_{i+1} \to \infty$. Define ${}^*r = {}^*\varepsilon{}^*s$ and $\mu' = \langle {}^*r\rangle$. Notice that $\mu' \notin {}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ and $\mu' < \mu$. Let $\nu \in (\mu', \mu)$ be represented by $\{x_i\}$ monotone and tending to 0 in which $r_i < x_i < s_i$ for all i. Then $x_i/x_{i+1} > r_i/s_{i+1} = \varepsilon_i s_i/s_{i+1} \to \infty$. Thus $(\mu', \mu) \subset ({}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon})^{\mathsf{C}}$. A similar argument shows that there exists $\mu'' > \mu$ with $(\mu, \mu'') \subset ({}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon})^{\mathsf{C}}$, thus ${}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ is closed. On the other hand, let $\mu \in {}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ be represented by $\{s_i\}$ with $s_i/s_{i+1} < M$ for all i. Choose $0 < {}^*\varepsilon \in {}^*\mathbb{R}_{\varepsilon}$ such that $\varepsilon_i/\varepsilon_{i+1} < N$ for all i, let ${}^*r = {}^*\varepsilon{}^*s$ and $\mu' = \langle {}^*r\rangle$. Let $\nu \in (\mu', \mu)$ be represented by $\{x_i\}$ monotone and tending to 0. Then we may write $x_i = M_i r_i$ with $M_i \to \infty$; note then that (trivially) $M_i/M_{i+1} < M'$ for some M'. Then $x_i/x_{i+1} < M'MN$ for all i and thus $(\mu', \mu) \subset {}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$. By the symmetry of the argument, we have $\mu \in (\mu', \mu'') \subset {}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ for some $\mu'' > \mu$, and so ${}^*\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ is open as well. □

Theorem 10. *Let* $\theta \in \mathbb{R} - \mathbb{Q}$*. Then* ${}^{\star}\mathbb{PR}^{\mathrm{sh}}_{\varepsilon} \subset \mathrm{Spec}_{\mathsf{flat}}(\theta)^{\mathsf{c}}$*. In particular,*

$$\mathrm{Spec}_{\mathsf{flat}}(\theta) \subsetneqq {}^{\star}\mathbb{PR}_{\varepsilon}.$$

Proof. If θ is badly approximable the result follows trivially from Theorem 6. So we assume θ is well approximable. Now suppose that $0 \neq {}^{*}m \in {}^{*}\mathbb{Z}^{\mu}_{\mu}(\theta)$ where $\mu \in {}^{\star}\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$. Then we may find monotone representatives $\{s_i\}$ of μ and $\{m_i\}$ of ${}^{*}m$ for which $m_i s_i \to 0$ and for which there exists M such that $s_i/s_{i+1} < M$ for all i. In particular, the first fact says that there exists an infinite sequence $\{R_i\}$ such that $s_i = (R_i m_i)^{-1}$. On the other hand, by Theorem 9, ${}^{*}m \in {}^{*}\mathbb{Z}^{\infty}_{\mathrm{wmb}}(\theta)$. Thus if ${}^{*}m = {}^{*}x \cdot {}^{*}\widehat{q}$ and we represent $\widehat{q} = \{q_{n_i}\}$ then $q_{n_{i+1}}/q_{n_i} \to \infty$. It follows that for any representative $\{x_i\}$ of ${}^{*}x$

$$\frac{s_i}{s_{i+1}} = \frac{R_{i+1} x_{i+1} q_{n_{i+1}}}{R_i x_i q_{n_i}} \longrightarrow \infty$$

(since $|x_i| \geq 1$ as ${}^{*}x \in {}^{*}\mathbb{Z}$) contradicting the shift invariance of μ.

$\square$

Note 4. The shift invariant set ${}^{\star}\mathbb{PR}^{\mathrm{sh}}_{\varepsilon}$ is greater (in the order $<$) than its complement in ${}^{\star}\mathbb{PR}_{\varepsilon}$; its elements may be characterized as the classes of "slow" infinitesimals. Theorem 10 says that the flat spectrum of θ irrational contains no slow indices.

5. Ideological Arithmetic

In this section we use the growth-decay filtration to provide the diophantine approximation groups with a partially defined multiplicative structure subject to matching conditions along the growth-decay indices. We begin with a result which describes the sense in which diophantine approximation groups generalize ideals.

Proposition 14. *For all* $\mu, \nu, \iota, \lambda \in {}^{\star}\mathbb{PR}_{\varepsilon}$ *there exists an action*

$${}^{*}\mathbb{Z}^{\nu[\iota]} \times {}^{*}\mathbb{Z}^{\mu[\lambda]}_{\nu}(\theta) \longrightarrow {}^{*}\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota}(\theta), \quad ({}^{*}a, {}^{*}n) \mapsto {}^{*}a{}^{*}n$$

in which $({}^{*}a{}^{*}n)^{\perp} = {}^{*}a{}^{*}n^{\perp}$*.*

Proof. For ${}^{*}a \in \mathbb{Z}^{\nu}$, ${}^{*}a{}^{*}n\theta = {}^{*}a{}^{*}n^{\perp} + {}^{*}a\varepsilon({}^{*}n)$. Since ${}^{*}a \cdot \nu < \iota \in {}^{\star}\mathbb{PR}_{\varepsilon}$, ${}^{*}a\varepsilon({}^{*}n) \in {}^{*}\mathbb{R}_{\varepsilon}$. $\square$

A **(fine) ideology** of ${}^{*}\mathbb{Z}$ is a subgroup $\mathfrak{a} \subset {}^{*}\mathbb{Z}$ equipped with a filtration by subgroups $\mathfrak{a} = \{\mathfrak{a}_{\nu}\}$ indexed by $\nu \in {}^{\star}\mathbb{PR}_{\varepsilon}$ for which

$${}^{*}\mathbb{Z}^{\nu} \cdot \mathfrak{a}^{\mu}_{\nu} \subset \mathfrak{a}^{\mu\cdot\nu} \quad ({}^{*}\mathbb{Z}^{\nu[\iota]} \cdot \mathfrak{a}^{\mu[\lambda]}_{\nu} \subset \mathfrak{a}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota})$$

for all $\nu \in {}^{\star}\mathbb{PR}_{\varepsilon}$ (for all $\mu, \nu, \iota, \lambda \in {}^{\star}\mathbb{PR}_{\varepsilon}$), where $\mathfrak{a}^{\mu}_{\nu} = \mathfrak{a}_{\nu} \cap {}^{*}\mathbb{Z}^{\mu}$ (where $\mathfrak{a}^{\mu[\lambda]}_{\nu} = \mathfrak{a}_{\nu} \cap {}^{*}\mathbb{Z}^{\mu[\lambda]}$). If one forgets the filtrations, what is left is the usual notion of ideal. By Proposition 14, diophantine approximation groups are ideologies.

More generally, one can define a **modulogy** (over ${}^{*}\mathbb{Z}$) as a bi-filtered abelian group $M = \{M^{\mu}_{\nu}\}$ in which there is an action ${}^{*}\mathbb{Z}^{\nu} \cdot M^{\mu}_{\nu} \subset M^{\mu\cdot\nu}$. A homomorphism $f : M \to N$ between modulogies is called a **modulogical homomorphism** if

MH1 it is filtered: $f(M^{\mu}_{\nu}) \subset N^{\mu}_{\nu}$ for all $\mu, \nu \in {}^{\star}\mathbb{PR}_{\varepsilon}$.

MH2 it respects the ${}^{*}\mathbb{Z}$ action: for all $\mu, \nu \in {}^{\star}\mathbb{PR}_{\varepsilon}$, ${}^{*}m \in {}^{*}\mathbb{Z}^{\nu}$ and $x \in M^{\mu}_{\nu}$

$$f({}^{*}m \cdot x) = {}^{*}m \cdot f(x).$$

If we set ${}^{*}\mathbb{Z}_{\nu} := {}^{*}\mathbb{Z}^{\nu}$ then ${}^{*}\mathbb{Z}$ is a modulogy over itself, and any ideology is a modulogy.

Proposition 15. *Let* $\theta, \eta \in \mathbb{R}$. *If* $\theta \asymp \eta$ *by* $A \in \mathrm{PGL}_2(\mathbb{Z})$ *then the induced isomorphism* $A : {}^*\mathbb{Z}(\theta) \cong {}^*\mathbb{Z}(\eta)$ *is modulogical.*

Proof. By Theorem 3 we already know that $\theta \asymp \eta$ by $A \in \mathrm{PGL}_2(\mathbb{Z})$ induces a filtered isomorphism. If ${}^*m \in {}^*\mathbb{Z}^{\nu}$, ${}^*n \in {}^*\mathbb{Z}^{\mu}_{\nu}$ then MH2 follows from:

$$A({}^*m \cdot {}^*n) = c({}^*m \cdot {}^*n)^{\perp} + d({}^*m \cdot {}^*n) = c^*m \cdot {}^*n^{\perp} + d^*m \cdot {}^*n = {}^*m \cdot A({}^*n).$$

□

The following result forms the basis of ideological arithmetic.

Theorem 11 (Ideological Arithmetic)**.** *Let* $\theta, \eta \in \mathbb{R}$. *Then*

$$ {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta) \cdot {}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}(\eta) \;\subset\; {}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta\eta) \;\cap\; {}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta+\eta) \;\cap\; {}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta-\eta). \tag{16}$$

In particular, ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \cdot {}^*\mathbb{Z}^{\nu}_{\mu}(\eta) \subset {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta\eta) \cap {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta+\eta) \cap {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta-\eta)$. *Moreover, for* ${}^*m \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ *and* ${}^*n \in {}^*\mathbb{Z}^{\nu}_{\mu}(\eta)$

$$({}^*m \cdot {}^*n)^{\perp_{\theta\eta}} = {}^*m^{\perp_\theta} \cdot {}^*n^{\perp_\eta} \quad \textit{and} \quad ({}^*m \cdot {}^*n)^{\perp_{\theta\pm\eta}} = {}^*m^{\perp_\theta} \cdot {}^*n \pm {}^*m \cdot {}^*n^{\perp_\eta} \tag{17}$$

Proof. We prove first that the left hand side of (16) is contained in ${}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta\eta)$. Given ${}^*m \in {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta)$ and ${}^*n \in {}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}(\eta)$,

$$\theta\eta({}^*m{}^*n) = {}^*m^{\perp}{}^*n^{\perp} + \varepsilon({}^*m)\varepsilon({}^*n) + \varepsilon({}^*m){}^*n^{\perp} + \varepsilon({}^*n){}^*m^{\perp}. \tag{18}$$

By hypothesis, the cross terms on the right hand side of (18), $\varepsilon({}^*m){}^*n^{\perp}$ and $\varepsilon({}^*n){}^*m^{\perp}$, are infinitesimals: here we are using (9). Thus $\theta\eta({}^*m{}^*n)$ is infinitesimal to ${}^*m^{\perp}{}^*n^{\perp}$ and ${}^*m{}^*n \in {}^*\mathbb{Z}(\theta\eta)$. In particular, $({}^*m{}^*n)^{\perp} = {}^*m^{\perp}{}^*n^{\perp}$, giving the product duality in (17). Moreover, $({}^*m{}^*n)\cdot(\mu\cdot\nu) = ({}^*m\cdot\mu)\cdot({}^*n\cdot\nu) < \iota\cdot\lambda$ so that ${}^*m{}^*n \in {}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}(\theta\eta)$. By (9) again, ${}^*m^{\perp}\cdot\nu({}^*n) \leq {}^*m^{\perp}\cdot\mu < \iota$, ${}^*n^{\perp}\cdot\nu({}^*m) \leq {}^*n^{\perp}\cdot\nu < \lambda$. Thus $\nu({}^*m{}^*n)$ satisfies the bound $\nu({}^*m{}^*n) \leq (\mu\cdot\nu) + \iota + \lambda$. Since ${}^*n$ is infinite, ${}^*n\cdot\nu < \lambda$ implies that $\nu < \lambda$ and therefore $\mu\cdot\nu < \mu\cdot\lambda < \lambda$. Hence $(\mu\cdot\nu) + \iota + \lambda = \iota + \lambda$, and ${}^*m{}^*n \in {}^*\mathbb{Z}_{\iota+\lambda}(\theta\eta)$ as claimed. As for the inclusion into ${}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta+\eta)$, this follows from the additive analog of (18),

$$(\theta+\eta)({}^*m{}^*n) = {}^*m^{\perp}{}^*n + {}^*m{}^*n^{\perp} + \varepsilon({}^*m){}^*n^{\perp} + \varepsilon({}^*n){}^*m^{\perp}.$$

The proof of the inclusion into ${}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta-\eta)$ is identical. □

The product defined by (16) will be referred to as the **growth-decay** or **ideological product**. While the ideological product has image in ${}^*\mathbb{Z}(\theta\eta)$, it is not the case that it has image in ${}^*\mathbb{Z}(\theta\eta^{-1})$: in contrast with the additive situation, in which the image of (16) is contained in both ${}^*\mathbb{Z}(\theta+\eta)$ and ${}^*\mathbb{Z}(\theta-\eta)$. That the ideological product yields diophantine approximations of the product, sum and difference has to do with the fact that it is essentially the precursor of the product of associated "two generator" ideologies (generated by "decoupled" numerators and denominators). This will be taken up in §15.

Note 5. By definition of the growth-decay trifiltration, $\mu < \iota$, $\nu < \lambda$ so that $\mu\cdot\nu < \iota+\lambda$. Thus the image of the product (16) is contained in the groups indexed by the slow components of $\mathrm{Spec}(\theta\eta)$, $\mathrm{Spec}(\theta+\eta)$.

There is no harm in stating the obvious: the formulas (17) for the additive and multiplicative duals are just the formulas for the numerators of fractional sum and product:

$$\frac{{}^*m^{\perp_\theta}}{{}^*m}\cdot\frac{{}^*n^{\perp_\eta}}{{}^*n}=\frac{({}^*m{}^*n)^{\perp_{\theta\eta}}}{{}^*m{}^*n},\qquad \frac{{}^*m^{\perp_\theta}}{{}^*m}\pm\frac{{}^*n^{\perp_\eta}}{{}^*n}=\frac{{}^*m^{\perp_\theta}\,{}^*n\pm{}^*m{}^*n^{\perp_\eta}}{{}^*m{}^*n}=\frac{({}^*m{}^*n)^{\perp_{\theta\pm\eta}}}{{}^*m{}^*n},$$

formulas which are compatible with their standard parts: the product and sum/difference of θ and η. When $\theta,\eta\in\mathbb{Q}$ the growth decay product is just the product on the fine growth filtration:

$${}^*\mathbb{Z}^{\mu[\iota]}(\theta)\cdot{}^*\mathbb{Z}^{\nu[\lambda]}(\eta)\subset{}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}(\theta\eta)\cap{}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}(\theta\pm\eta)$$

which reduces upon restriction to standard parts to $\mathbb{Z}(\theta)\cdot\mathbb{Z}(\eta)=\mathbb{Z}(\theta\eta)\cap\mathbb{Z}(\theta\pm\eta)$. This is in keeping with the fact that the diophantine approximation group of an element of $a/b\in\mathbb{Q}$ is the ideal ${}^*(b)$ generated by its denominator.

Let us compare ideological arithmetic with ideal arithmetic in $\mathbb{Z}$. For $a\in\mathbb{Z}$, let ${}^*(a)=$ the ideal generated by a in ${}^*\mathbb{Z}$. Then given $a,b\in\mathbb{Z}$, clearly ${}^*\mathbb{Z}(a)^\perp={}^*(a)$, ${}^*\mathbb{Z}(b)^\perp={}^*(b)$, ${}^*\mathbb{Z}(a+b)^\perp={}^*(a+b)$, ${}^*\mathbb{Z}(ab)^\perp={}^*(ab)$. The map (16) induces on the level of dual groups a pair of bilinear maps corresponding to the sum and the product. The map corresponding to the product is

$${}^*\mathbb{Z}(a)^\perp\times{}^*\mathbb{Z}(b)^\perp\longrightarrow{}^*\mathbb{Z}(ab)^\perp,\quad({}^*m\cdot a,{}^*n\cdot b)\mapsto({}^*m\cdot a,{}^*n\cdot b)^{\perp_{ab}}={}^*m{}^*n\cdot ab,$$

which corresponds exactly to the usual product of ideals.

On the other hand, the map of dual groups corresponding to the sum is

$${}^*\mathbb{Z}(a)^\perp\times{}^*\mathbb{Z}(b)^\perp\longrightarrow{}^*\mathbb{Z}(a+b)^\perp\subsetneqq{}^*(a)+{}^*(b)$$

since it is given by

$$({}^*m\cdot a,{}^*n\cdot b)\mapsto({}^*m\cdot a,{}^*n\cdot b)^{\perp_{a+b}}={}^*m{}^*n\cdot(a+b).$$

In particular, the image of the duality map is not all of ${}^*(a)+{}^*(b)$, since the elements belonging to the latter are *independent* linear combinations of a,b. This situation will be addressed in §15.

Let $\mu\geq\nu$. We will write

$$\theta\,{}_{\mu}\!\owedge_{\nu}\,\eta,\quad\left(\theta\,{}_{\mu[\iota]}\!\owedge_{\nu[\lambda]}\,\eta\right)\tag{19}$$

to indicate that both of ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$, ${}^*\mathbb{Z}^{\nu}_{\mu}(\eta)$ (both of ${}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta)$, ${}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}(\eta)$) are non-zero, so that the growth-decay product is non-trivial. When $\mu>\nu$ strictly, then $\theta\,{}_{\mu}\!\owedge_{\nu}\,\eta$ exactly when ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ is non-trivial, by Theorem 4. In particular, for $\mu>\nu$, $\theta\,{}_{\mu}\!\owedge_{\nu}\,\eta$ implies that $\theta\,{}_{\mu'}\!\owedge_{\nu'}\,\eta$ whenever $\mu\geq\mu'>\nu'\geq\nu$.

The relation ${}_{\mu}\!\owedge_{\nu}$ is not symmetric (commutative) i.e. (19) does not imply that $\eta\,{}_{\mu}\!\owedge_{\nu}\,\theta$; in fact, as we will see below, (19) does not even imply that $\eta\,{}_{\mu'}\!\owedge_{\nu'}\,\theta$ for *some* pair $(\mu',\nu')\in{}^*\mathbb{PR}_\varepsilon$ with $\mu'\geq\nu'$.

In view of Theorem 11, only the "tri-filtered" relation ${}_{\mu[\iota]}\!\owedge_{\nu[\lambda]}$ can be iterated, and in view of Note 5, iterated compositions move *to the left.* More precisely the iterated relation

$$\xi\,{}_{\mu'[\iota']}\!\owedge_{\nu'[\lambda']}\left(\theta\,{}_{\mu[\iota]}\!\owedge_{\nu[\lambda]}\,\eta\right)$$

is *defined* provided $\iota+\lambda\leq\mu'$, $\nu'\leq\mu\cdot\nu$ and $\lambda'\geq\iota\cdot\lambda$. Intuitively, as one continues to iterate, the nonvanishing spectrum of the real number appearing on the left will become progressively larger.

The ideological product is natural with respect to multiplication and the duality/reciprocal maps: $({}^*m\cdot{}^*n)^{\perp_{\theta\eta}} = {}^*m^{\perp_\theta}\cdot{}^*n^{\perp_\eta}$ for ${}^*m \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ and ${}^*n \in {}^*\mathbb{Z}^{\nu}_{\mu}(\eta)$. That is, in terms of the growth-decay bi-filtration,

$$\begin{tikzcd}
{}^*\mathbb{Z}^{\mu}_{\nu}(\theta)\times{}^*\mathbb{Z}^{\nu}_{\mu}(\eta) \arrow[r, "\cdot"] \arrow[d, "\cong", "\perp\times\perp"'] & {}^*\mathbb{Z}^{\mu\cdot\nu}(\theta\eta) \arrow[d, "\cong"', "\perp"] \\
{}^*\mathbb{Z}^{\mu}_{\nu}(\theta^{-1})\times{}^*\mathbb{Z}^{\nu}_{\mu}(\eta^{-1}) \arrow[r, "\cdot"] & {}^*\mathbb{Z}^{\mu\cdot\nu}((\theta\eta)^{-1})
\end{tikzcd}$$

with a similar diagram for the fine growth-decay tri-filtration. If we fix ${}^*m \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ then multiplication by ${}^*m$, ${}^*n \mapsto {}^*m\cdot{}^*n$, defines a linear map from ${}^*\mathbb{Z}^{\nu}_{\mu}(\eta)$ onto its image in ${}^*\mathbb{Z}^{\mu\cdot\nu}(\theta\eta)$.

For the remainder of this section we will regard the growth-decay product as defining a bilinear map to the product approximation group

$$ {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}(\theta)\cdot{}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}(\eta) \longrightarrow {}^*\mathbb{Z}^{\mu\cdot\nu[\iota\cdot\lambda]}_{\iota+\lambda}(\theta\eta). $$

All statements which follow will have a corresponding additive counterpart, obtained by replacing the word "divisor" by "summand".

Let $\omega = \theta\eta$ and suppose that $\theta_{\mu}\oslash_{\nu}\eta$ for $\mu > \nu$. We say that θ is a μ/ν-**fast divisor** of ω (η is a ν/μ-**slow divisor** of ω) and we write

$$\theta\ {}_{\mu}{\Uparrow}_{\nu}\ \omega \quad (\eta\ {}_{\mu}{\Downarrow}_{\nu}\ \omega).$$

In addition, we write $\theta \Uparrow \omega$ ($\eta \Downarrow \omega$) to mean that $\theta\ {}_{\mu}{\Uparrow}_{\nu}\ \omega$ ($\eta\ {}_{\mu}{\Downarrow}_{\nu}\ \omega$) for some $\mu > \nu$. Fast divisors (slow divisors) have error terms which tend to zero more rapidly (slowly) than their denominators tend to infinity. These designations are not symmetric, and we will see below that the badly approximable numbers are never fast divisors.

Proposition 16. $\theta\ {}_{\mu}{\Uparrow}_{\nu}\ \omega$ *for all* $\mu > \nu \Leftrightarrow \theta \in \mathbb{Q}$.

Proof. ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ is nontrivial for all $\mu > \nu \Leftrightarrow \theta \in \mathbb{Q}$. □

If $\theta_{\mu}\oslash_{\mu}\eta$, we say that both θ and η are μ-**flat divisors** of ω and write

$$\theta \talloblong \omega = \theta \talloblong_{\mu} \omega\ , \quad \eta \talloblong \omega = \eta \talloblong_{\mu} \omega.$$

Thus $\theta \talloblong \omega \Leftrightarrow \mathrm{Spec}_{\mathrm{flat}}(\theta)\cap\mathrm{Spec}_{\mathrm{flat}}(\eta) \neq \emptyset$. If $\theta \Downarrow \omega$, $\theta \Uparrow \omega$ we will write $\theta \Updownarrow \omega$ and say that θ is an **elastic divisor**; if θ is elastic and $\theta \talloblong \omega$ as well then we will write $\theta\overline{\Updownarrow}\omega$ say that θ is a **strong elastic divisor** of ω. Note that if $\theta\overline{\Updownarrow}\omega$ and $\eta = \omega/\theta$ then $\eta\overline{\Updownarrow}\omega$ as well.

Proposition 17. $\theta\ {}_{\mu}{\Uparrow}_{\nu}\ \omega$, $\theta\ {}_{\mu}{\Downarrow}_{\nu}\ \omega$ *and* $\theta \Updownarrow_{\mu} \omega$ *for all* $\mu \geq \nu \Leftrightarrow \theta, \omega \in \mathbb{Q}$.

Proof. Trivial. □

If $\omega = \theta\eta$ but $\theta \not\Uparrow \omega$, $\theta \not\Downarrow \omega$ and $\theta \not\talloblong \omega$ we say that θ is an **antidivisor** and write

$$\theta \lightning \omega.$$

Note that if $\theta\lightning\omega$ and $\eta = \omega/\theta$ it is not necessarily the case that $\eta\lightning\omega$ (c.f. Theorem 12, c. below). If on the other hand for any ω, θ is a divisor (of some speed: fast, slow or flat) we say that θ is an **omnidivisor**.

We now examine the notions of divisibility described above with regard to the classes $\mathfrak{B}$, $\mathfrak{W}_1$, $\mathfrak{W}_{>1}$ and $\mathfrak{W}_{\infty}$.

Theorem 12. *Let* $\omega \in \mathbb{R}$.

a. *For all* $\theta \in \mathfrak{B}$, $\theta \not\Uparrow \omega$, $\theta \not\square \omega$.
b. *For all* $\theta \in \mathfrak{W}_1$, $\theta \not\Uparrow \omega$.
c. *If* $\omega = \theta\eta$, $\theta \in \mathfrak{B}$ *and* $\eta \in \mathfrak{B} \cup \mathfrak{W}_1$ *then* $\theta \not\updownarrow \omega$.
d. *If* $\omega = \theta\eta$, $\theta, \eta \in \mathfrak{W}_{>1}$ *then* $\theta, \eta \Updownarrow \omega$. *If moreover* θ *and* η *are equivalent then* $\theta, \eta \overline{\Updownarrow} \omega$.

Proof. a. & b. The spectrum of any element of $\mathfrak{B}$ (of $\mathfrak{W}_1$) consists exactly of pairs $\mu < \nu$ (consists of pairs which satisfy $\mu \leq \nu$), so flat or fast composition (fast composition) with θ is not possible. c. This follows trivially from the definitions. d. For any $\theta \in \mathfrak{W}_{>1}$ there exists pair $\mu > \nu$ in $\mathrm{Spec}(\theta)$ so $\theta\,{}_\mu\oslash_\nu\,\eta$. Switching the roles of θ and η, for appropriate $\mu > \nu$, $\eta\,{}_\mu\oslash_\nu\,\theta$ as well. Thus $\theta, \eta \Updownarrow \omega$. If θ and η are equivalent then $\mathrm{Spec}_{\mathrm{flat}}(\theta) = \mathrm{Spec}_{\mathrm{flat}}(\eta)$ implying $\theta, \eta \overline{\Updownarrow} \omega$. ☐

Corollary 3. *The set of omnidivisors is precisely* $\mathfrak{W}_{>1}$.

Proof. By Theorem 12, parts a., b., the set of omnidivisors is contained in $\mathfrak{W}_{>1}$; by part d., every element of $\theta \in \mathfrak{W}_{>1}$ is an omnidivisor. ☐

A subset $X \subset \mathbb{R}$ is called an **antiprime set** if for all $\theta_1, \theta_2 \in X$, $\theta_1, \theta_2 \not\updownarrow \omega = \theta_1\theta_2$.

Proposition 18. $\mathfrak{B}$ *is the unique maximal antiprime set in* $\mathbb{R}$.

Proof. Note that $\mathfrak{B}$ is an anti-prime set: $\theta_1, \theta_2 \not\updownarrow \theta_1\theta_2$ for all $\theta_1, \theta_2 \in \mathfrak{B}$. Moreover, if we add another element $\eta \notin \mathfrak{B}$ we lose the defining property since for such η, $\eta \,\square\, \eta^2$. On the other hand, any $\theta \in \mathbb{R} - \mathfrak{B}$ is composable with itself, so there are no antiprime sets containing elements not in $\mathfrak{B}$. ☐

Thus we shall refer to $\mathfrak{B}$ as the **antiprimes** of ideological arithmetic. We say that ω has an **antiprime decomposition** if $\omega = \theta\eta$ for $\theta, \eta \in \mathfrak{B}$. For example, every $q \in \mathbb{Q}$ which is not a square has the antiprime decomposition $q = \sqrt{q}\sqrt{q}$. Antiprime decompositions are outside the realm of the version of ideological arithmetic presented here, to analyze them requires the finer arithmetic of *symmetric diophantine approximations*, the subject of §7.

Theorem 13. *Every non zero real number admits an antiprime decomposition.*

Proof. Let $F(n) = \{\theta \in \mathfrak{B} \mid \forall i\ a_i(\theta) \leq n\}$. By [17], every real number $r > 1$ is a product $r = \theta\eta$ of elements of $F(4)$. Since $\mathfrak{B}$ is closed under inversion, the claim follows. ☐

A set X is said to be **(strongly) ideologically generated** by $X_0 \subset X$ if for all $\omega \in X$, there exist $\theta_1, \theta_2 \in X_0$ such that $\theta_1\theta_2 = \omega$ with $\theta_1, \theta_2 \Updownarrow \omega$ ($\theta_1, \theta_2 \overline{\Updownarrow} \omega$).

Corollary 4. *The set of Liouville numbers* $\mathfrak{W}_\infty$ *ideologically generates* $\mathbb{R}$.

Proof. By [9] every real number ω may be written as a product of $\theta_1, \theta_2 \in \mathfrak{W}_\infty$ and clearly $\theta_1, \theta_2 \Updownarrow \omega$. ☐

6. Flat Arithmetic

In this section we consider the commutative relation ${}_\mu\oslash_\mu$, which is best understood using the sequence of partial quotients $\theta = [a_0 a_1 \ldots]$. As alluded to at the beginning of §4, this makes the determination of the flat product somewhat transverse to the linear classification $\mathfrak{W}$, $\mathfrak{W}_\kappa$, $\mathfrak{W}_\infty$.

Given $\theta \in \mathfrak{W}$, the basic problem is to determine the set of $\eta \in \mathfrak{W}$ which have a flat product with θ:

$$\Omega(\theta) = \{\eta \in \mathfrak{W} |\ \theta_{\mu} \owedge_{\mu} \eta \text{ for some } \mu \in \mathrm{Spec}_{\mathrm{flat}}(\theta)\}.$$

It is clear that $\Omega(\theta)$ is a projective linear invariant:

Proposition 19. *If* $\theta \Bumpeq \eta$ *then* $\Omega(\theta) = \Omega(\eta)$ *and* $\Omega(\theta) \supset \{A(\theta) |\ A \in \mathrm{PGL}_2(\mathbb{Z})\}$.

Proof. Immediate from Proposition 9. □

The following Proposition gives a simple criterion in terms of best classes for when flat products are defined.

Lemma 1. *Let* $\theta, \eta \in \mathfrak{W}$. *Then there exists* $\mu \in {}^*\mathbb{PR}_{\varepsilon}$ *such that* $\theta_{\mu} \owedge_{\mu} \eta \Leftrightarrow$ *there exist best classes* ${}^*\widehat{q} \in {}^*\mathbb{Z}(\theta)$, ${}^*\widehat{q}' \in {}^*\mathbb{Z}(\eta)$ *having infinite partial quotient such that either*

$$\widehat{\mu}^+ \leq (\widehat{\mu}')^+ < \widehat{\mu} \quad \text{or} \quad (\widehat{\mu}')^+ \leq \widehat{\mu}^+ < \widehat{\mu}'. \tag{20}$$

Proof. By Corollary 2, a flat composition is defined $\Leftrightarrow$ there exists best classes having infinite partial quotient ${}^*\widehat{q} \in {}^*\mathbb{Z}^{\infty}_{\mathrm{b}}(\theta)$, ${}^*\widehat{q}' \in {}^*\mathbb{Z}^{\infty}_{\mathrm{b}}(\eta)$ for which

$$[\widehat{\nu}, \widehat{\mu}) \cap [\widehat{\nu}', \widehat{\mu}') \neq \emptyset. \tag{21}$$

Because ${}^*\widehat{q} \in {}^*\mathbb{Z}(\theta)$ has infinite partial quotient, $\widehat{\nu} = \widehat{\mu}^+ = \langle {}^*\widehat{q}^+ \rangle^{-1} < \widehat{\mu} = \langle {}^*\widehat{q} \rangle^{-1}$, since

$$\widehat{\nu} = \left\langle \frac{{}^*\theta}{{}^*\theta {}^*\widehat{q}^+ + {}^*\widehat{q}} \right\rangle = \langle {}^*\widehat{q}^+ \rangle^{-1} = \widehat{\mu}^+.$$

Thus (21) is equivalent to $[\widehat{\mu}^+, \widehat{\mu}) \cap [(\widehat{\mu}')^+, \widehat{\mu}') \neq \emptyset$ which in turn is equivalent to (20). □

We begin by introducing a conspicuous class of numbers for which flat products are always defined: $\theta = [a_0 a_1 \dots] \in \mathfrak{W}$ is called **resolute** if $\lim a_i = \infty$ (i.e. there are no bounded subsequences). For example, $\theta = [1, 2, 3, \dots]$ is resolute whereas $e = [2; 1, 2, 1, 1, 4, 1, 1, \dots]$ is not. It is fairly easy to produce Liouville numbers which are resolute and Liouville numbers which are not:

Example 1. Let $\theta = [a_0 a_1 \dots]$ be defined inductively by taking $a_0 = a_1 = 1$ and $a_{n+1} = q_n^{n-1}$ for $n \geq 1$. Then $\| q_n \theta \| < q_{n+1}^{-1} < q_n^{-n}$ so that θ is a resolute Liouville number. If instead we define $a_n = q_{n-1}^n$ for n even and $a_n = 1$ for n odd, then for n odd, we have $\| q_n \theta \| < q_{n+1}^{-1} < q_n^{n+1}$ so that θ is Liouville but irresolute. The Liouville number $L(m) = \sum_{j=0}^{\infty} m^{-(j+1)!}$ is irresolute for all $m > 1$ since 1 occurs infinitely often in its sequence of partial quotients [24].

Define the relation

$$\theta \triangleright_{\mu} \eta$$

if $\theta_{\mu} \owedge_{\mu} \eta$ and there exists ${}^*\widehat{q} \in {}^*\mathbb{Z}^{\mu}_{\mu}(\theta)$, ${}^*\widehat{q}' \in {}^*\mathbb{Z}^{\mu}_{\mu}(\eta)$ with $\widehat{\mu}^+ \leq (\widehat{\mu}')^+ < \widehat{\mu}$.

Theorem 14. *If* $\theta \in \mathfrak{W}$ *is resolute then for all* $\eta \in \mathfrak{W}$ *there exists* $\mu \in \mathrm{Spec}_{\mathrm{flat}}(\theta)$ *with* $\theta \triangleright_{\mu} \eta$.

Proof. For $\theta \in \mathfrak{W}$ resolute every best class ${}^*\widehat{q}$ has infinite partial quotient. Now let ${}^*\widehat{q}'$ be a best class for η with infinite partial quotient, represented by the sequence $\{q'_{n_i}\}$. For each i choose q_{n_i} so that $q_{n_i} \leq q'_{n_i+1} \leq q_{n_i+1}$. Then denoting ${}^*\widehat{q} = {}^*\{q_{n_i}\}$, we have ${}^*\widehat{q} \leq ({}^*\widehat{q}')^+ \leq \widehat{q}^+$. Since ${}^*\widehat{q}$ has infinite partial quotient we must have either $\langle {}^*\widehat{q} \rangle < \langle ({}^*\widehat{q}')^+ \rangle$ or $\langle ({}^*\widehat{q}')^+ \rangle < \langle {}^*\widehat{q}^+ \rangle$. If the former is true, this implies the first inequality in (20). If not, $\langle {}^*\widehat{q}\rangle = \langle ({}^*\widehat{q}')^+\rangle$; then replace ${}^*\widehat{q}$ by ${}^*\widehat{q}^- = {}^*\{q_{n_i-1}\}$ to get the desired inequality. $\square$

With Lemma 1 this gives:

Corollary 5. *If* $\theta \in \mathfrak{W}$ *is resolute,* $\Omega(\theta) = \mathbb{R} - \mathfrak{B}$.

For any class $\mathfrak{C}$ of real numbers, we denote by $\mathfrak{C}^{\rm res}$ the resolute members. For example, $\mathfrak{B}^{\rm res} = \varnothing$ and $\mathfrak{W}^{\rm res}_{\infty} \subsetneqq \mathfrak{W}_{\infty}$.

Corollary 6. *If* $\theta, \eta \in \mathfrak{W}^{\rm res}_{>1}$ *and* $\omega = \theta\eta$ *then* $\theta, \eta \overline{\Updownarrow} \omega$. *In particular,* $\theta, \eta \in \mathfrak{W}^{\rm res}_{>1}$ *strongly ideologically generates* $\mathfrak{W}^{\rm res}_{>1} \cdot \mathfrak{W}^{\rm res}_{>1}$.

We will now consider the other extreme: pairs of real numbers which may not be flat composed at all. Constructing such pairs depends on finding real numbers which have large gaps in their flat spectra: that is, having increasingly long blocks of partial quotients all of whose members are uniformly bounded. More precisely we say that θ has an **abyss** if there exist a sequence of consecutive blocks of partial quotients $G = \{B_n\}$, $B_n = \{a_{i_n}, \dots, a_{i_n+k_n}\}$ which are pairwise disjoint, each element of which is bounded by a constant M independent of n, such that the block length $k_n \to \infty$. In this event we say that θ is **abyssal**; if θ is not abyssal, it is called **abyssless**. For example: any resolute number is abyssless; e is abyssless yet irresolute, as is the Liouville number $L(m)$ [24]. Thus the set of abyssal real numbers is a strict subset of the irresolute real numbers. On the other hand, one can easily create Liouville numbers which are abyssal following a procedure similar to that used in *Note* 1.

Proposition 20. $\theta \in \mathfrak{W}$ *is abyssal* $\Leftrightarrow$ $\mathrm{Spec}_{\rm flat}(\theta)$ *is disconnected.*

Proof. $\Rightarrow$ Suppose θ is abyssal. Then we may find a best class ${}^*\widehat{q}$ for which each element of the bi-infinite sequence of predecessors and successors

$$\cdots {}^*\widehat{q}^{--} < {}^*\widehat{q}^{-} < {}^*\widehat{q} < {}^*\widehat{q}^{+} < {}^*\widehat{q}^{++} < \cdots$$

defines the same element of ${}^*\mathbb{P}\mathbb{R}$. From this it follows that $\widehat{\mu} = \widehat{\nu} = \langle {}^*\widehat{q}\rangle^{-1} \notin \mathrm{Spec}_{\rm flat}(\theta)$. On the other hand, since $\theta \in \mathfrak{W}$, there exist best classes ${}^*\widehat{q}_1, {}^*\widehat{q}_2$ with infinite partial quotient for which $\widehat{\nu}_1 < \widehat{\nu} < \widehat{\nu}_2$. $\Leftarrow$ If θ is abyssless then any best class ${}^*\widehat{q}$ which has bounded partial quotient is finitely many successors as well as finitely many predecessors away from a best class with infinite bounded quotient. This implies that the common growth and decay class of ${}^*\widehat{q}$, $\widehat{\mu} = \widehat{\nu}$, belongs to $\mathrm{Spec}_{\rm flat}(\theta)$ since it occurs as the right and left endpoint of a pair of adjacent best intervals $[\widehat{\nu}', \widehat{\mu}')$, $[\widehat{\nu}'', \widehat{\mu}'')$ corresponding to neighboring best classes with infinite bounded quotient. Thus $\mathrm{Spec}_{\rm flat}(\theta)$ is connected. $\square$

We have the following strengthening of Corollary 5.

Theorem 15. *If* θ *is abyssless then* $\Omega(\theta) = \mathbb{R} - \mathfrak{B}$. *For any* $\eta \in \mathbb{R} - \mathfrak{B}$, $\mathrm{Spec}_{\rm flat}(\eta) \subset \mathrm{Spec}_{\rm flat}(\theta)$.

Proof. Let $\eta \in \mathbb{R} - \mathfrak{B}$ and let ${}^*\widehat{q}' \in {}^*\mathbb{Z}(\eta)$ be a best class with infinite partial quotient for η. There exist best classes ${}^*\widehat{q}_1, {}^*\widehat{q}_2$ with infinite partial quotient for θ such that ${}^*\widehat{q}_1 < {}^*\widehat{q}' < ({}^*\widehat{q}')^+ < {}^*\widehat{q}_2$. By the connectivity of $\mathrm{Spec}_{\rm flat}(\theta)$, $[\widehat{\nu}', \widehat{\mu}') \subset \mathrm{Spec}_{\rm flat}(\theta)$. $\square$

Corollary 7. *If* $\theta \in \mathbb{R} - \mathfrak{B}$ *(*$\theta \in \mathfrak{W}_{>1} \cup \mathbb{Q}$*) is abyssless,* $\theta \mathbin{[\!]} \omega$ *(*$\theta \overline{\Updownarrow} \omega$*) for all* $\omega \in \mathbb{R}$ *with* $\theta\omega^{-1} \notin \mathfrak{B}$.

Corollary 8. *If* θ, η *are abyssless, then* $\mathrm{Spec}_{\mathrm{flat}}(\eta) = \mathrm{Spec}_{\mathrm{flat}}(\theta)$.

We call this common connected set of Corollary 8 the **flat interval** ${}^{*}\mathbb{PR}_{\mathrm{flat}}$. In particular, ${}^{*}\mathbb{PR}_{\varepsilon} - {}^{*}\mathbb{PR}_{\mathrm{flat}}$ is disjoint from all flat spectra and includes the shift invariant elements. Note also that it is clear that there exist abyssless θ, η which are not equivalent yet their flat spectra coincide: so $\mathrm{Spec}_{\mathrm{flat}}(\theta)$ is not a complete invariant of θ. An abyssless number θ is an **omniflatdivisor**: an element $\theta \in \mathfrak{W}$ in which $\theta \,\square\, \omega$ for all $\omega \in \mathbb{R}$ with $\theta\omega^{-1} \notin \mathfrak{B}$.

The next result shows that the set $\Omega(\theta)$ need not be equal to $\mathbb{R} - \mathfrak{B}$.

Theorem 16. *There exist abyssal* $\theta, \eta \in \mathfrak{W}$ *such that* $\theta_{\mu} \oslash_{\mu} \eta$ *is undefined for all* μ. *In fact, one can find such a pair in which* $\theta \in \mathfrak{W}_{\kappa}$, $\eta \in \mathfrak{W}_{\kappa'}$ *for any* $\kappa, \kappa' \in [1, \infty]$.

Proof. We construct θ, η by way of partial fractions. We begin by specifying the initial partial fractional segment $[\mathbf{1}_1 a_{m_1}]$ of θ, where $\mathbf{1}_1$ is a large block of ones of size $m_1 - 1$. Let $q_{m_1} = a_{m_1} q_{m_1 - 1} + q_{m_1 - 2}$ be the best denominator corresponding to a_{m_1}. Now specify the initial segment for η, $[\mathbf{1}_1' b_{n_1}]$ so that for N_1 a fixed integers with $2N_1 < n_1$, there are

i. N_1 best denominators associated to the 1's of $\mathbf{1}_1'$ that are less than q_{m_1} i.e.
$$q_1', \dots, q_{N_1}' < q_{m_1}.$$
If necessary we go back and choose m_1 larger so that this can be done.

ii. N_1 best denominators associated to the 1's of $\mathbf{1}_1'$ which are greater than q_{m_1}:
$$q_{n_1 - 1}', \dots, q_{n_1 - 1 - N_1}' > q_{m_1}.$$

The next step is to select $M_2 > N_1$ so that if $\mathbf{1}_2$ is a block of $1's$ having at least $2M_2$ elements then at least M_2 of the new best denominators associated to the augmented sequence $[\mathbf{1}_1 a_{m_1 + 1} \mathbf{1}_2]$ are less than q_{n_1}' i.e.
$$q_{m_1 + 1}, \dots, q_{m_1 + M_2} < q_{n_1}'.$$
Then choose $m_2 > 2M_2$ so that if $\mathbf{1}_2$ has m_2 elements than at least M_2 of the new best denominators associated to the augmented sequence $[\mathbf{1}_1 a_{m_1} \mathbf{1}_2]$ are greater than q_{n_1}':
$$q_{m_1 + m_2}, \dots, q_{m_1 + m_2 - M_2} > q_{n_1}'.$$
Let $a_{m_1 + m_2} > a_{m_1}$ and consider $[\mathbf{1}_1 a_{m_1} \mathbf{1}_2 a_{m_1 + m_2}]$, applying the above procedure to specify $[\mathbf{1}_1' b_{n_1} \mathbf{1}_2' b_{n_1 + n_2}]$, where $b_{n_1} < b_{n_1 + n_2}$. Inductively we build the sequence of partial quotients of θ and η in this way, arranging that the non 1 partial quotients
$$a_{m_1 + \cdots + m_k}, b_{n_1 + \cdots + n_k},$$
as well as the M_k, N_k, tend to ∞. See Figure 2.

Consider the sequences
$$\{q_{m_1 + \cdots + m_k - 1}\}_{k=1}^{\infty}, \quad \{q'_{n_1 + \cdots + n_k - 1}\}_{k=1}^{\infty} \tag{22}$$
and let ${}^{*}\widehat{q}$, ${}^{*}\widehat{q}'$ be best classes for θ, η having infinite partial quotient. By construction of the partial fraction sequences, ${}^{*}\widehat{q}$, ${}^{*}\widehat{q}'$ are classes of sequences formed from the elements of the corresponding sequences of (22). Moreover, our choices of blocks of 1's in the partial quotients of θ and η ensure that

- $({}^{*}\widehat{q}')^{+} \notin [{}^{*}\widehat{q}, {}^{*}\widehat{q}^{+}]$ and that $({}^{*}\widehat{q}')^{+}, {}^{*}\widehat{q}, {}^{*}\widehat{q}^{+}$ define distinct classes in ${}^{*}\mathbb{PR}$.
- ${}^{*}\widehat{q}^{+} \notin [{}^{*}\widehat{q}', ({}^{*}\widehat{q}')^{+}]$ and that ${}^{*}\widehat{q}^{+}, {}^{*}\widehat{q}', ({}^{*}\widehat{q}')^{+}$ define distinct classes in ${}^{*}\mathbb{PR}$.



Figure 2. Construction of θ, η.

This implies that the growth decay interval $[\widehat{\mu}^+, \widehat{\mu})$ corresponding to ${}^*\widehat{q}$ cannot contain the decay $(\widehat{\mu}')^+$ of ${}^*\widehat{q}'$, and *vice verca*. By Lemma 1, $\theta\,{}_{\mu}\!\oslash_{\mu}\eta$ is undefined for all μ. By choosing the non 1 partial fractions appropriately we can ensure that $\theta \in \mathfrak{W}_{\kappa}$, $\eta \in \mathfrak{W}_{\kappa'}$ for any fixed $\kappa, \kappa' \in [1, \infty]$. □

It follows that the problems of determining $\Omega(\theta)$ and flat composability are non trivial. Nevertheless, it seems plausible that the techniques in [24] can be extended to show that the Liouville numbers occurring in the sum and product representations [9] are abyssless.

Conjecture. *$\mathfrak{W}_{\infty}$ strongly ideologically generates $\mathbb{R}$.*

7. Symmetric Diophantine Approximations and Lorentzian Groupologies

The study of special approximations in which the the error term is dominated by a function $\psi : \mathbb{Z} - \{0\} \to \mathbb{R}$ has held, from the very beginning, a distinguished position in the subject of Diophantine Approximation. Classically, for a fixed ψ, one looks for conditions on θ which guarantee the existence of infinitely many solutions to the inequality $|n\theta - m| < |\psi(n)|^{-1}$, or equivalently, in the language of this paper, a single solution to

$$|\varepsilon({}^*n)| = |{}^*n\theta - {}^*n^{\perp}| < |\psi({}^*n)|, \quad {}^*n \in {}^*\mathbb{Z}(\theta). \tag{23}$$

The Theorems of Dirichlet, Liouville and Roth all fall within this genre. More generally, if one only specifies convergence properties of the sum $\sum \psi(n)$, one seeks (e.g. Khintchine's Theorem) to measure the size of the set of real numbers having solutions to (23).

In this section, we will shift the emphasis from one of existence to a qualitative study of the solution set:

$$ {}^*\mathbb{Z}(\theta|\psi) := \{0 \neq {}^*n \in {}^*\mathbb{Z}(\theta) |\text{ there exists } C > 0 \text{ such that } |\varepsilon({}^*n)| < C|\psi({}^*n)|\} \cup \{0\}, $$

focusing on the extent to which the set ${}^*\mathbb{Z}(\theta|\psi)$ has interesting arithmetic structure. We note that trivially ${}^*\mathbb{Z}(\theta|\psi)$ is closed under the operation of taking additive inverses.

We begin by fixing the choice $\psi(x) = x^{-1}$: note then that

$$ {}^*\mathbb{Z}\big(\theta|x^{-1}\big) = \{{}^*n \in {}^*\mathbb{Z}(\theta) | \nu({}^*n) \leq \mu({}^*n)\}\,. \tag{24} $$

It turns out that the same hypothesis used to describe the ideological product structure of the ${}^*\mathbb{Z}(\theta)$ can be used to deduce an *additive* structure in ${}^*\mathbb{Z}\big(\theta|x^{-1}\big)$. To this end, denote

$$ {}^*\mathbb{Z}^{\mu}_{\nu}\big(\theta|x^{-1}\big) := {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \cap {}^*\mathbb{Z}(\theta|x^{-1}). $$

Note that the defining condition $\nu({}^*n) \leq \mu({}^*n)$ in (24) *does not* imply that ${}^*\mathbb{Z}^{\mu}_{\nu}\big(\theta|x^{-1}\big) = 0$ for $\mu < \nu$. Recall that tropical subtraction in ${}^*\mathbb{PR}$ is defined

$$ \mu - \nu := \min(\mu, \nu). $$

Theorem 17. *Let* $\mu, \nu, \iota, \lambda \in {}^*\mathbb{PR}_{\varepsilon}$. *Then*

$$ {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}\big(\theta|x^{-1}\big) + {}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}\big(\theta|x^{-1}\big) \subset {}^*\mathbb{Z}^{\mu-\nu[\iota+\lambda]}\big(\theta|x^{-1}\big). $$

In particular,

$$ {}^*\mathbb{Z}^{\mu}_{\nu}\big(\theta|x^{-1}\big) + {}^*\mathbb{Z}^{\nu}_{\mu}\big(\theta|x^{-1}\big) \subset {}^*\mathbb{Z}^{\mu-\nu}\big(\theta|x^{-1}\big). $$

Proof. Let ${}^*m \in {}^*\mathbb{Z}^{\mu[\iota]}_{\nu}\big(\theta|x^{-1}\big)$, ${}^*n \in {}^*\mathbb{Z}^{\nu[\lambda]}_{\mu}\big(\theta|x^{-1}\big)$. Then

$$ \left|({}^*m + {}^*n)^2\theta - ({}^*m + {}^*n)({}^*m^{\perp} + {}^*n^{\perp})\right| \leq C + \left|2{}^*m{}^*n\theta - ({}^*m{}^*n^{\perp} + {}^*m^{\perp}{}^*n)\right| $$

for some constant C. But

$$ \begin{aligned} |2{}^*m{}^*n\theta - ({}^*m{}^*n^{\perp} + {}^*m^{\perp}{}^*n)| &= \theta^{-1}\left|({}^*m\theta - {}^*m^{\perp})({}^*n\theta - {}^*n^{\perp}) + ({}^*m{}^*n\theta^2 - {}^*m^{\perp}{}^*n^{\perp})\right| \\ &= \theta^{-1}\left|\varepsilon({}^*m)\cdot\varepsilon({}^*n) + \varepsilon({}^*m{}^*n)\right|. \end{aligned} $$

which is infinitesimal (here we are using Theorem 11 to conclude that $\varepsilon({}^*m{}^*n) = {}^*m{}^*n\theta^2 - {}^*m^{\perp}{}^*n^{\perp}$ is the error term of a diophantine approximation, hence is infinitesimal). Thus ${}^*m + {}^*n \in {}^*\mathbb{Z}\big(\theta|x^{-1}\big)$. Finally we note that

$$ ({}^*m + {}^*n)\cdot(\mu - \nu) \leq {}^*m\cdot(\mu - \nu) + {}^*n\cdot(\mu - \nu) < \iota + \lambda. $$

Thus ${}^*m + {}^*n \in {}^*\mathbb{Z}^{\mu-\nu[\iota+\lambda]}\big(\theta|x^{-1}\big)$. □

With its sum partially defined along the growth-decay filtration, we refer to ${}^*\mathbb{Z}\big(\theta|x^{-1}\big)$ as a **groupology**; the sum is referred to as the **growth-decay sum** or **groupological sum**. We say that the groupology ${}^*\mathbb{Z}\big(\theta|x^{-1}\big)$ is **barren** if there are no non-trivial grupological sums.

Corollary 9. *Let* $\theta \in \mathbb{R} - \mathbb{Q}$. *Then* ${}^*\mathbb{Z}\big(\theta|x^{-1}\big)$ *is barren* $\Leftrightarrow$ $\theta \in \mathfrak{B}$.

Proof. $\Leftarrow$ Immediate from Theorem 6. $\Rightarrow$ If $\theta \in \mathfrak{W}$ there exists μ for which ${}^*\mathbb{Z}^{\mu}_{\mu}\big(\theta|x^{-1}\big) \neq 0$. In this case we have the non-trivial growth-decay sum

$$ {}^*\mathbb{Z}^{\mu}_{\mu}\big(\theta|x^{-1}\big) + {}^*\mathbb{Z}^{\mu}_{\mu}\big(\theta|x^{-1}\big) \subset {}^*\mathbb{Z}^{\mu}\big(\theta|x^{-1}\big). $$

□

Recall [13] the real vector space ${}^{\bullet}\mathbb{R} := {}^*\mathbb{R}/{}^*\mathbb{R}_{\varepsilon} \supset \mathbb{R}$ of **extended reals**; we define a function $|\cdot|_{\theta} : {}^*\mathbb{Z} \to {}^{\bullet}\mathbb{R}_{+}$ by

$$ |{}^*n|_{\theta} := \big(|{}^*n| \cdot \|{}^*n\theta\|\big)^{1/2} \mod {}^*\mathbb{R}_{\varepsilon}. $$

Note that when ${}^*n \in {}^*\mathbb{Z}(\theta)$, $\|{}^*n\theta\| = |\varepsilon({}^*n)|$. By definition, $|{}^*n|_{\theta} \in \mathbb{R}_{+}$ for all ${}^*n \in {}^*\mathbb{Z}\big(\theta|x^{-1}\big)$. Somewhat abusively, we refer to $|\cdot|_{\theta}$ as the **θ-norm**; while it is technically not a norm, it may be viewed as a generalized norm in a sense which will soon be made clear. We have immediately:

Proposition 21. *$|\cdot|_{\theta} \equiv 0$ on ${}^*\mathbb{Z}^{\mu}_{\nu}\big(\theta|x^{-1}\big)$ for $\mu \geq \nu$.*

Note 6. For each $\theta \in \mathfrak{B}$, let $C_{\theta} > 0$ be the supremum of constants C for which $\|n\theta\| < Cx^{-1}$ has only finitely many solutions. The set of such C_{θ} as one ranges over $\theta \in \mathfrak{B}$ is called the Lagrange spectrum [22]. Note that if $\theta \in \mathfrak{B}$ and $C_{\theta} > 0$ is the associated element of the Lagrange spectrum then for all $0 \neq {}^*n \in {}^*\mathbb{Z}\big(\theta|x^{-1}\big)$

$$ |{}^*n|_{\theta} \geq C_{\theta}^{1/2}. $$

Thus for badly approximable numbers, the θ-norm is always positive on non 0 elements.

For arbitrary $\theta \in \mathbb{R} - \mathbb{Q}$, do there exist any ${}^*n \in {}^*\mathbb{Z}\big(\theta|x^{-1}\big)$ for which $|{}^*n|_{\theta} \neq 0$? If there exists such an ${}^*n$ then there are positive real constants $c < C$ such that some representative sequence $\{n_i\}$ satisfies the inequality

$$ \frac{c}{n_i} \leq |n_i\theta - n_i^{\perp}| \leq \frac{C}{n_i} \tag{25} $$

i.e. $\langle |{}^*n| \cdot |\varepsilon({}^*n)| \rangle = \langle |{}^*n| \rangle \cdot \nu({}^*n) = 1$, or equivalently,

$$ \mu({}^*n) = \nu({}^*n). $$

We call such a class ${}^*n$ a **symmetric diophantine approximation**, the set of which union 0 is denoted

$$ {}^*\mathbb{Z}^{\mathrm{sym}}(\theta) := |\cdot|_{\theta}^{-1}(0,\infty) \cup \{0\} \subset {}^*\mathbb{Z}\big(\theta|x^{-1}\big). $$

We have trivially that ${}^*n \in {}^*\mathbb{Z}^{\mathrm{sym}}(\theta) \Leftrightarrow N \cdot {}^*n \in {}^*\mathbb{Z}^{\mathrm{sym}}(\theta)$ for $N \in \mathbb{Z} - \{0\}$.

We will show that ${}^*\mathbb{Z}^{\mathrm{sym}}(\theta) \neq 0$ for all $\theta \in \mathbb{R} - \mathbb{Q}$. For each $\nu \in {}^{\star}\mathbb{PR}_{\varepsilon}$, write

$$ {}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) = \{{}^*n \in {}^*\mathbb{Z}^{\mathrm{sym}}(\theta) |\ \nu({}^*n) = \nu\}, $$

so that ${}^*\mathbb{Z}^{\mathrm{sym}}(\theta) = \bigcup_{\nu} {}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$. The following Proposition identifies ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$ as the part of ${}^*\mathbb{Z}_{\nu}(\theta)$ inhabiting the narrow space between the intersection of the slow diophantine approximations of decay ν and the flat diophantine approximations of decay ν.

Proposition 22. *Let $\theta \in \mathbb{R}$. Then ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) = \Big(\bigcap_{\mu<\nu} {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)\Big) - {}^*\mathbb{Z}^{\nu}_{\nu}(\theta)$.*

Proof. ${}^*n \in {}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) \Leftrightarrow {}^*n \cdot \nu = {}^*n \cdot \nu({}^*n) = 1$ in ${}^{\star}\mathbb{PR} \Leftrightarrow {}^*n \notin {}^*\mathbb{Z}^{\nu}_{\nu}(\theta)$ and ${}^*n \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$ for all $\mu < \nu$. □

Corollary 10. *If* $\theta \in \mathfrak{B}$ *then*

$$^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) = \bigcap_{\mu<\nu} {}^*\mathbb{Z}^{\mu}_{\nu}(\theta).$$

In particular, $^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$ *is a subgroup of* $^*\mathbb{Z}_{\nu}(\theta)$.

Recall (see §3) the set of best growths $^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon} \subset {}^*\mathbb{PR}_{\varepsilon}$.

Theorem 18. *For all* $\theta \in \mathbb{R}-\mathbb{Q}$ *and all* $\nu \in {}^*\mathbb{PR}^{\mathrm{bg}}_{\varepsilon}$, $^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) \neq 0$. *In particular, if* $\theta \in \mathfrak{B}$ *then* $^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta) \neq 0$ *for all* $\nu \in {}^*\mathbb{PR}_{\varepsilon}$.

Proof. For $\theta \in \mathfrak{B}$ the result is obviously true: any best class $^*\widehat{q} \in {}^*\mathbb{Z}^{\mathrm{sym}}(\theta)$, and moreover, we may realize any growth index ν as a best growth by Theorem 6. Now assume that $\theta \notin \mathfrak{B} \cup \mathbb{Q}$. Let $\theta = [a_1 a_2 \dots]$. By Grace's Theorem [19], the intermediate best denominator (= denominator of the intermediate convergent) $q_{n,r} = rq_{n+1} + q_n$, where $0 \leq r < a_{n+2}$, satisfies

$$1 < |q_{n,r}| \cdot \|q_{n,r}\theta\|$$

for all $r \neq 0, 1, a_{n+2}-1$. For any infinite sequence of such intermediate convergents we take $c = 1$ in (25). On the other hand (Lemma I.4 of [19]), we have

$$|q_{n,r}| \cdot \|q_{n,r}\theta\| = \frac{q_{n,r}(\theta_{n+2}-r)}{\theta_{n+2}q_{n+1}+q_n} = \frac{(rq_{n+1}+q_n)(\theta_{n+2}-r)}{\theta_{n+2}q_{n+1}+q_n}$$

where (as in Theorem 9) θ_i is defined by the equation $\theta = [a_1 \dots a_{i-1}\theta_i]$. We note that $r < a_{n+2} < \theta_{n+2}$. Dividing out numerator and denominator by the dominant term $\theta_{n+2}q_{n+1}$ gives

$$\begin{aligned} |q_{n,r}| \cdot \|q_{n,r}\theta\| &= \frac{\Big((r/\theta_{n+2}) + \big(q_n/(\theta_{n+2}q_{n+1})\big)\Big)(\theta_{n+2}-r)}{1+\big(q_n/(\theta_{n+2}q_{n+1})\big)} \\ &< r + \frac{q_n}{q_{n+1}} - r\left(\frac{r}{\theta_{n+2}} + \frac{q_n}{\theta_{n+2}q_{n+1}}\right) \\ &< r+1. \end{aligned}$$

Then if we choose r_i bounded and $\neq 0, 1, a_{n_i+2}-1$, the class $^*n = {}^*\{q_{n_i,r_i}\}$ will be symmetric. It follows that for any best class $^*\widehat{q}$ there is a symmetric class $^*n$ such that $\mu(^*\widehat{q}) = \mu(^*n)$. □

Corollary 11. *If* $\theta \in \mathfrak{B}$ *then* $^*\mathbb{Z}^{\mathrm{sym}}(\theta)$ *contains both the best denominator classes as well as the intermediate best denominator classes.*

Proof. From the proof of Theorem 18, we know that the intermediate best denominators of the form $^*n = {}^*\{q_{n_i,r_i}\}$, where $r_i \neq 0, 1, a_{n_i+2}-1$ and is uniformly bounded, belong to $^*\mathbb{Z}^{\mathrm{sym}}(\theta)$. On the other hand, since $\theta \in \mathfrak{B}$, the best denominators (which occur for $r_i = 0, 1$) as well as the consecutive difference $^*\widehat{q}^{+} - {}^*\widehat{q}$ (which occurs for $r_i = a_{n_i+2}-1$) belong to $^*\mathbb{Z}^{\mathrm{sym}}(\theta)$. □

Note 7. As the above paragraphs show, the function $|\cdot|_{\theta}$ is nontrivial for all $\theta \in \mathbb{R}-\mathbb{Q}$. We take a moment to contrast it with its rational and p-adic analogs.

i. If $\theta = q = a/b \in \mathbb{Q}$ then $|\cdot|_q \equiv 0$ on ${}^*\mathbb{Z}(\theta) = {}^*\mathbb{Z}(\theta|x^{-1})$. For ${}^*n \in {}^*\mathbb{Z}$ arbitrary, $|{}^*n|_q = c \cdot |{}^*n|$ where $c = a'/b$ for some $a' < b$. In fact, $|\cdot|_q$ induces a function
$$|\cdot|_q : {}^*\mathbb{Z}/{}^*\mathbb{Z}(\theta) \cong \mathbb{Z}/b\mathbb{Z} \longrightarrow {}^*\mathbb{Q}/{}^*\mathbb{Z} \cong \mathbb{Q}/\mathbb{Z}.$$

ii. If $\xi \in \mathbb{Q}_p = p$-adic numbers and we use the p-adic absolute value to define the distance-to-the-nearest-integer function $\|\cdot\|$, then $\|\cdot\| \equiv 0$ on ${}^*\mathbb{Z}(\xi) = {}^*\mathbb{Z}(\xi|x^{-1})$. Therefore, for ${}^*n \in {}^*\mathbb{Z}$ arbitrary, $|{}^*n|_\xi \leq |\xi|_p$; if $\xi \in \hat{\mathbb{Z}}_p$ then $|\cdot|_\xi \equiv 0$.

◆

Recall that the **Littlewood conjecture** asserts that for any pair $\theta, \eta \in \mathfrak{B}$,
$$\liminf |n|\|n\theta\|\|n\eta\| = 0.$$

Observation 1. Given $\theta, \eta \in \mathfrak{B}$, suppose that $\exists \nu \in {}^*\mathbb{PR}_\varepsilon$ such that
$$\{0\} \subsetneqq {}^*\mathbb{Z}^{\mathrm{sym}}_\nu(\theta) \cap {}^*\mathbb{Z}^\nu(\eta) \quad \text{or} \quad \{0\} \subsetneqq {}^*\mathbb{Z}^\nu(\theta) \cap {}^*\mathbb{Z}^{\mathrm{sym}}_\nu(\eta).$$
Then the Littlewood conjecture holds for the pair θ, η.

In view of Observation 1 it would be of interest to find as precise a description as possible of ${}^*\mathbb{Z}^{\mathrm{sym}}(\theta)$. When $\theta = \varphi = (\sqrt{5}+1)/2$ is the golden mean there are many symmetric diophantine approximations which are neither intermediate nor principle convergents; we characterize them now. Recall [29] that every natural number has a unique **Zeckendorf representation**

$$N = F_{i_1} + \cdots + F_{i_k} \tag{26}$$

where F_k = the kth Fibonacci number, and the sequence $i_1 < \cdots < i_k$ consists of non consecutive integers ≥ 2. Using Binet's formula [20]
$$F_k = \frac{\varphi^k - (-1)^k \varphi^{-k}}{\sqrt{5}}$$
one has the following

Lemma 2. *Let $N \in \mathbb{N}$ have the Zeckendorf representation (26). Then $\|N\varphi\| < \varphi^{-n}$ $\Leftrightarrow$*

- $i_1 \geq n+1$.
- $i_1 = n$ *and* $i_2 - i_1$ *is odd and* ≥ 3.

Proof. See [8]. □

Define the **Zeckendorf degree** of the representation (26) to be
$$\text{Z-deg}(N) := i_k - i_1.$$
For ${}^*n \in {}^*\mathbb{Z}$ the Zeckendorf degree is an element of ${}^*\mathbb{N}$.

Theorem 19. *Let φ be the golden mean. Then*
$${}^*\mathbb{Z}^{\mathrm{sym}}(\varphi) = \{{}^*n \in {}^*\mathbb{Z}(\varphi) | \text{ Z-deg}({}^*n) < \infty\}.$$

Proof. Let ${}^*n = {}^*\{n_i\}$ and let M be a bound for Z-deg$({}^*n)$. Then by Lemma 2 and Binet's formula

$$|{}^*n|\cdot|\varepsilon({}^*n)| \leq C\varphi^M$$

for $C > 0$ a constant that depends only on φ. On the other hand, if Z-deg$({}^*n) \in {}^*\mathbb{N} - \mathbb{N}$ then

$$|{}^*n|\cdot|\varepsilon({}^*n)| \geq \varphi^{\text{Z-deg}({}^*n)} \mod {}^*\mathbb{R}_\varepsilon .$$

□

Returning to the Littlewood conjecture, we see that the following is true. Let

$$Z_D(\varphi) \subset \mathbb{Z}[\varphi]$$

be the set of $\mathbb{Q}(\sqrt{5})$-integers of the form

$$f(\varphi) = \varphi^{i_1} + \cdots + \varphi^{i_k}$$

with $i_1 < \cdots < i_k$ non consecutive, $i_1 \geq 2$ and relative degree $i_k - i_1 < D$.

Corollary 12. *Let* $\theta \in \mathbb{R}$, $\theta \notin \mathbb{Q}(\sqrt{5})$. *If for some* D

$$\{\|f(\varphi)\theta\| \; f(\varphi) \in Z_D(\varphi)\}$$

is dense in $[0, 1/2)$ *then the Littlewood conjecture holds for the pair* (φ, θ).

Note 8. The hypothesis of Corollary 12 in the case of $D = 1$ reduces to Chowla's conjecture in the case of the golden mean [21].

✠

We turn to the matter of the general arithmetic structure of ${}^*\mathbb{Z}^{\rm sym}(\theta)$, describing ${}^*\mathbb{Z}^{\rm sym}(\theta) = \{{}^*\mathbb{Z}^{\rm sym}_\nu(\theta)\}$ as a subgroupology of ${}^*\mathbb{Z}(\theta|x^{-1})$. For every sign pair $\sigma \in \{\pm\}^2$, let ${}^*\mathbb{Z}(\theta)_\sigma$ be the monoid consisting of 0 and those diophantine approximations ${}^*n \in {}^*\mathbb{Z}(\theta)$ for which $(\text{sign}({}^*n), \text{sign}(\varepsilon({}^*n))) = \sigma$. Let ${}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma = {}^*\mathbb{Z}^{\rm sym}_\nu(\theta) \cap {}^*\mathbb{Z}(\theta)_\sigma$.

Theorem 20. *Let* $\theta \in \mathbb{R}$.

1. ${}^*\mathbb{Z}^{\rm sym}(\theta) = \{{}^*\mathbb{Z}^{\rm sym}_\nu(\theta)\}$ *satisfies*
$$ {}^*\mathbb{Z}^{\rm sym}_\nu(\theta) + {}^*\mathbb{Z}^{\rm sym}_\nu(\theta) \subset {}^*\mathbb{Z}\big(\theta|x^{-1}\big).$$
2. *For* $\sigma = (+,+)$ *or* $(-,-)$ *and* $\nu \in {}^*\mathbb{P}\mathbb{R}_\varepsilon$, ${}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma$ *is a monoid:*
$$ {}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma + {}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma \subset {}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma .$$

Proof. The first assertion is a consequence of the inequality

$$|{}^*m + {}^*n|\cdot|\varepsilon({}^*m + {}^*n)| \leq M + |{}^*n||\varepsilon({}^*m)| + |{}^*m||\varepsilon({}^*n)| :$$

as the terms $|{}^*n||\varepsilon({}^*m)|$, $|{}^*m||\varepsilon({}^*n)|$ are bounded. The second assertion follows by noting that for ${}^*m, {}^*n \in {}^*\mathbb{Z}^{\rm sym}_\nu(\theta)_\sigma$,

$$\langle|{}^*m + {}^*n|\rangle = \langle|{}^*m|\rangle + \langle|{}^*n|\rangle = \nu^{-1}, \quad \langle|\varepsilon({}^*m + {}^*n)|\rangle = \langle|\varepsilon({}^*m)|\rangle + \langle|\varepsilon({}^*n)|\rangle = \nu.$$

□

Corollary 13. *If* $\theta \in \mathfrak{B}$ *then* ${}^*\mathbb{Z}^{\rm sym}(\theta) = \{{}^*\mathbb{Z}^{\rm sym}_\nu(\theta)\}$ *is a family of groups satisfying*

$$ {}^*\mathbb{Z}^{\nu[\iota]} \cdot {}^*\mathbb{Z}^{\rm sym}_\nu(\theta) \subset {}^*\mathbb{Z}^{\nu^2}_\iota(\theta). \tag{27}$$

Proof. If $\theta \in \mathfrak{B}$ then ${}^*\mathbb{Z}\big(\theta|x^{-1}\big) = {}^*\mathbb{Z}^{\mathrm{sym}}(\theta)$. By 1. of Theorem 20, ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$ is a group for all $\nu \in {}^*\mathbb{PR}_{\varepsilon}$. The property (27) follows immediately from the definitions. $\square$

The symmetric set ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$ has, in addition, a fractional addition/multiplication law which generalizes the ideological product of Theorem 11, and which is nontrivial even for $\theta \in \mathfrak{B}$. To formulate it it is necessary to work with numerator denominator pairs rather than just denominators. Thus let

$$ {}^*\mathbb{Z}^{1,1}(\theta) = \{({}^*n^{\perp}, {}^*n) |\, {}^*n \in {}^*\mathbb{Z}(\theta)\} = \{({}^*m, {}^*n) \in {}^*\mathbb{Z}^2 |\, \langle|{}^*n\theta - {}^*m|\rangle < 1\} \subset {}^*\mathbb{Z}^2 $$

be the associated group of numerator denominator pairs of diophantine approximations. The canonical isomorphism ${}^*\mathbb{Z}^{1,1}(\theta) \cong {}^*\mathbb{Z}(\theta)$ induces the growth-decay filtration ${}^*\mathbb{Z}^{1,1}(\theta) = \{({}^*\mathbb{Z}^{1,1})^{\mu}_{\nu}(\theta)\}$. The set of symmetric numerator denominator pairs is denoted

$$ ({}^*\mathbb{Z}^{1,1})^{\mathrm{sym}}_{\nu}(\theta) \subset ({}^*\mathbb{Z}^{1,1})_{\nu}(\theta). $$

Now define

$$ {}^*\mathring{\mathbb{Z}}^{1,1}(\theta) = \{({}^*m, {}^*n) \in {}^*\mathbb{Z}^2 |\, \langle|{}^*n\theta - {}^*m|\rangle \leq 1\} \supsetneq {}^*\mathbb{Z}^{1,1}(\theta). $$

Note that $\mathbb{Z}^2 \subset {}^*\mathring{\mathbb{Z}}^{1,1}(\theta)$, and if $({}^*m, {}^*n) \in {}^*\mathring{\mathbb{Z}}^{1,1}(\theta) - \mathbb{Z}^2$ then both ${}^*m$ and ${}^*n$ are infinite. In addition, for all infinite $({}^*m, {}^*n) \in {}^*\mathring{\mathbb{Z}}^{1,1}(\theta)$, ${}^*m/{}^*n \simeq \theta$ in ${}^*\mathbb{R}$. If we form the quotient group

$$ {}^*\overline{\mathbb{Z}}^{1,1}(\theta) := {}^*\mathring{\mathbb{Z}}^{1,1}(\theta)/\mathbb{Z}^2 $$

then every class ${}^*\bar{n} \in {}^*\overline{\mathbb{Z}} := {}^*\mathbb{Z}/\mathbb{Z}$ (= the group of universes in ${}^*\mathbb{Z}$, a densely ordered group) determines a unique numerator ${}^*\bar{m} \in {}^*\overline{\mathbb{Z}}$ for which $({}^*\bar{m}, {}^*\bar{n}) \in {}^*\overline{\mathbb{Z}}^{1,1}(\theta)$ and we write ${}^*\bar{n}^{\perp} = {}^*\bar{m}$. Since elements of ${}^*\mathbb{Z}^{1,1}(\theta)$ are already uniquely determined by their denominator, there is an induced inclusion

$$ {}^*\mathbb{Z}^{1,1}(\theta) \hookrightarrow {}^*\overline{\mathbb{Z}}^{1,1}(\theta). $$

For $({}^*\bar{n}^{\perp}, {}^*\bar{n}) \in {}^*\overline{\mathbb{Z}}^{1,1}(\theta)$ write $\nu({}^*\bar{n}) = \nu({}^*n)$ if ${}^*n \in {}^*\bar{n}$ belongs to ${}^*\mathbb{Z}(\theta)$; otherwise write $\nu({}^*\bar{n}) = 1$. In addition, write $\mu({}^*\bar{n}) = \langle {}^*n^{-1}\rangle$ for any ${}^*n \in {}^*\bar{n}$, which is evidently independent of the choice of representative.

Let ${}^*\mathbb{PR}_{\leq 1} = {}^*\mathbb{PR}_{\varepsilon} \cup \{1\}$. Now for $\mu \in {}^*\mathbb{PR}_{\varepsilon}, \nu \in {}^*\mathbb{PR}_{\leq 1}$ define

$$ ({}^*\overline{\mathbb{Z}}^{1,1})^{\mu}_{\nu}(\theta) = \{({}^*\bar{m}, {}^*\bar{n}) |\, {}^*\bar{n} \cdot \mu \in {}^*\mathbb{PR}_{\varepsilon}, \nu({}^*\bar{n}) \leq \nu\}. $$

Note that for all $\nu < 1$, $({}^*\overline{\mathbb{Z}}^{1,1})^{\mu}_{\nu}(\theta) = ({}^*\mathbb{Z}^{1,1})^{\mu}_{\nu}(\theta) \cong {}^*\mathbb{Z}^{\mu}_{\nu}(\theta)$. The proof of the following Theorem is left to the reader, who will note that it is available for elements of $\mathfrak{B}$, providing the latter with a weak form of ideological arithmetic.

Theorem 21 (Symmetric Ideological Arithmetic). *For any* $\theta, \eta \in \mathbb{R}$ *and all* $\nu \in {}^*\mathbb{PR}_{\varepsilon}$*, there are maps*

$$ \cdot, \pm\, : ({}^*\mathbb{Z}^{1,1})^{\mathrm{sym}}_{\nu}(\theta) \times ({}^*\mathbb{Z}^{1,1})^{\mathrm{sym}}_{\nu}(\eta) \longrightarrow ({}^*\overline{\mathbb{Z}}^{1,1})^{\nu^2}_{1}(\theta \pm \eta), ({}^*\overline{\mathbb{Z}}^{1,1})^{\nu^2}_{1}(\theta\eta) $$

given by fractional product, sum and difference of pairs.

We now make precise the extent to which we may regard $|\cdot|_{\theta}$ as a generalized norm: to begin, the following result suggests that we may view $|\cdot|_{\theta}$ as a pseudo-norm on the groupology ${}^*\mathbb{Z}\big(\theta|x^{-1}\big)$.

Theorem 22. *The restriction of the function* $|\cdot|_{\theta}$ *to* ${}^*\mathbb{Z}(\theta|x^{-1})$ *obeys the non-archimedean triangle inequality for all defined sums.*

Proof. Let ${}^*n_1 \in {}^*\mathbb{Z}^{\mu}_{\nu}\big(\theta|x^{-1}\big)$, ${}^*n_2 \in {}^*\mathbb{Z}^{\nu}_{\mu}\big(\theta|x^{-1}\big)$ so that the sum ${}^*n_1 + {}^*n_2$ is defined. Then

$$|{}^*n_1 + {}^*n_2|^2_\theta \leq (|{}^*n_1| + |{}^*n_2|)(|\varepsilon({}^*n_1)| + |\varepsilon({}^*n_2)|) \mod {}^*\mathbb{R}_\varepsilon.$$

By hypothesis we have that the elements $|{}^*n_1|\cdot|\varepsilon({}^*n_2)|$, $|{}^*n_2|\cdot|\varepsilon({}^*n_1)|$ are infinitesimal, so that

$$|{}^*n_1 + {}^*n_2|^2_\theta \leq |{}^*n_1|^2_\theta + |{}^*n_2|^2_\theta.$$

However by Proposition 21, for $\mu \geq \nu$, $|{}^*n_1|^2_\theta = 0$, implying $|{}^*n_1 + {}^*n_2|_\theta \leq \max(|{}^*n_1|_\theta, |{}^*n_2|_\theta)$. □

Since at least one of the elements of any defined groupological sum already has θ-norm 0, Theorem 22 is a bit of a cheat. A more interesting norm-theoretic interpretation of $|\cdot|_\theta$ may be obtained by restricting to ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$. As it turns out, it is more natural to contemplate a *Minkowskian* formulation of $|\cdot|_\theta$.

Given ${}^*m, {}^*n \in {}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$, consider the following symmetric function in two-variables:

$$[{}^*m, {}^*n]_\theta := \frac{1}{2}\left({}^*m\varepsilon({}^*n) + {}^*n\varepsilon({}^*m)\right) \mod {}^*\mathbb{R}_\varepsilon \in \mathbb{R}$$

and write $[{}^*m]_\theta := [{}^*m, {}^*m]^{\frac{1}{2}}_\theta$ so that $|{}^*m|_\theta = |[{}^*m]_\theta|$. In particular, $|\cdot|_\theta$ is the "Minkowski norm" associated to $[\cdot,\cdot]_\theta$. We say that ${}^*m$ is **time-like** if $[{}^*m]^2_\theta > 0$ and **space-like** if $[{}^*m]^2_\theta < 0$. The time-like elements correspond to the signs $\sigma = (+,+), (-,-)$ and the space-like correspond to the signs $\sigma = (+,-), (-,+)$. We say that time-like elements point in the same direction if their sign σ is the same: the elements with sign $(+,+)$ are interpreted as future pointing. The function $[\cdot,\cdot]_\theta$ extends by the same formula to all of ${}^*\mathbb{Z}(\theta|x^{-1}) = \{{}^*\mathbb{Z}^{\mu}_{\nu}(\theta|x^{-1})\}$. We call an element ${}^*n \in {}^*\mathbb{Z}(\theta|x^{-1})$ **light-like** if $[{}^*m]_\theta = 0$ e.g. if ${}^*m \in {}^*\mathbb{Z}^{\mu}_{\nu}(\theta|x^{-1})$ for some $\mu \geq \nu$. The time-like and space-like elements of ${}^*\mathbb{Z}(\theta|x^{-1})$ are exactly the elements of ${}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$. If we reverse our clocks and view diophantine approximations as material particles "departing from" θ ($\mu({}^*n)$ = time and $\varepsilon({}^*n)$ = space) then $|{}^*n|_\theta$ is nothing more than the initial speed. When $\theta \in \mathfrak{B}$, we have the inequality

$$|[{}^*m]|^2_\theta \geq C_\theta \tag{28}$$

where C_θ is the corresponding element of the Lagrange spectrum. We may interpret C_θ as being Planck's constant for the system defined by the symmetric diophantine approximations of θ, where the inequality (28) plays the role of Heisenberg's uncertainty principle.

Theorem 23. *For all* ${}^*m_1, {}^*m_2 \in {}^*\mathbb{Z}^{\mathrm{sym}}_{\nu}(\theta)$,

$$[{}^*m_1 + {}^*m_2, {}^*n]_\theta = [{}^*m_1, {}^*n]_\theta + [{}^*m_2, {}^*n]_\theta.$$

If moreover ${}^*m_1, {}^*m_2$ *are time-like and point in the same direction, then they satisfy the reverse triangle inequality:*

$$[{}^*m_1 + {}^*m_2]_\theta \geq [{}^*m_1]_\theta + [{}^*m_2]_\theta.$$

Proof. The first statement is immediate. To prove the second it will suffice to prove the following reverse Cauchy inequality:

$$[{}^*m, {}^*n]_\theta \geq [{}^*m_1]_\theta [{}^*m_2]_\theta.$$

Indeed, suppose the latter is true, and $^*m_1, {^*m_2}$ are time-like of the same sign, then immediately:

$$\begin{aligned}[][^*m_1 + {^*m_2}]_\theta^2 = ({^*m_1} + {^*m_2})(\varepsilon({^*m_1}) + \varepsilon({^*m_2})) &= [^*m_1]_\theta^2 + [^*m_2]_\theta^2 + 2[^*m_1, {^*m_2}]_\theta \\ &\geq ([^*m_1]_\theta + [^*m_2]_\theta)^2.\end{aligned}$$

But the reverse Cauchy inequality follows from:

$$[^*m_1, {^*m_2}]_\theta^2 - [^*m_1]_\theta^2[^*m_2]_\theta^2 = \frac{1}{4}({^*m_1}\varepsilon({^*m_2}) - {^*m_2}\varepsilon({^*m_1}))^2 \geq 0.$$

□

We call $^*\mathbb{Z}^{\rm sym}(\theta) = \{{^*\mathbb{Z}}^{\rm sym}_\nu(\theta)\}$ equipped with the Minkowskian pairing $[\cdot,\cdot]_\theta$ a **Lorentzian groupology**. The Minkowskian norm has the following compatibility with the symmetric ideological product:

Proposition 23. *Suppose that* $^*m \in {^*\mathbb{Z}}^{\rm sym}_\nu(\theta), {^*n} \in {^*\mathbb{Z}}^{\rm sym}_\nu(\eta)$ *are either both time-like or both space-like. Then* $|[^*m^*n]_{\theta\eta}| = \infty$ *so that*

$$|[^*m^*n]_{\theta\eta}| > [^*m]_\theta[^*n]_\eta.$$

Proof. First note that $\varepsilon(^*m^*n) = {^*n}\varepsilon(^*m) + {^*m}\varepsilon(^*n) + \varepsilon(^*m)\varepsilon(^*n)$ and by hypothesis $^*n\varepsilon(^*m) \simeq r$, $^*n\varepsilon(^*m) \simeq s$ with r, s non-zero reals. These non-zero reals will be of the same sign if $^*m, {^*n}$ are either both time-like or both space-like: in this event, $|\varepsilon(^*m^*n)| \simeq |r| + |s| > 0$, hence $|[^*m^*n]_{\theta\eta}|$ is infinite and the result is trivially true. □

Theorem 24. *The group* $\mathrm{PGL}_2(\mathbb{Z})$ *acts by Lorentzian isometries: that is, for all* $^*m, {^*n} \in {^*\mathbb{Z}}^{\rm sym}_\nu(\theta)$,

$$[^*m, {^*n}]_\theta = [A(^*m), A(^*n)]_{A(\theta)}.$$

In particular, if $\theta \mathrel{\overset{\cdot}{\underset{\cdot}{\sim}}} \eta$ *then* $^*\mathbb{Z}^{\rm sym}(\theta) \cong {^*\mathbb{Z}}^{\rm sym}(\eta)$ *as Lorentzian groupologies.*

Proof. If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then as was seen in Theorem 3, $A(^*m) = c^*m^\perp + d^*m$ and $\varepsilon(A(^*m)) = (c\theta + d)^{-1}\varepsilon(^*m)$, with similar formulas for $A(^*n)$ and $\varepsilon(A(^*n))$. Thus

$$A(^*m)\varepsilon(A(^*m)) = \frac{c^*m^\perp + d^*m}{c\theta + d}\varepsilon(^*m) = \frac{c(^*m^\perp/^*m) + d}{c\theta + d}{^*m}\varepsilon(^*m) \simeq {^*m}\varepsilon(^*m)$$

and the result follows. □

We now define a family of norms indexed by general exponents. For each $\kappa > 1$ consider the function $x^{-\kappa}$ and the set $^*\mathbb{Z}(\theta|x^{-\kappa}) \subset {^*\mathbb{Z}}(\theta|x^{-1})$ with its associated "norm" function

$$|^*m|_{\theta,\kappa} = |^*m^\kappa\varepsilon(^*n)| \mod {^*\mathbb{R}_\varepsilon} \in \mathbb{R}.$$

Much of the discussion developed above for the case $\kappa = 1$ extends to $\kappa > 1$. We summarize the situation for $\kappa > 1$ leaving the straightforward verifications to the reader.

a. The set of **κ -symmetric diophantine approximations**, defined

$$^*\mathbb{Z}^{\rm sym}_\kappa(\theta) = \{{^*\mathbb{Z}}^{\rm sym}_{\nu,\kappa}(\theta)\} := |\cdot|^{-1}_{\theta,\kappa}(0,\infty) \subset {^*\mathbb{Z}}(\theta|x^{-1}),$$

satisfies the obvious analogue of Theorem 20.

b. If $\kappa \neq \kappa'$ then $^*\mathbb{Z}^{\rm sym}_\kappa(\theta) \cap {^*\mathbb{Z}}^{\rm sym}_{\kappa'}(\theta) = \{0\}$.

c. If θ has exponent κ then $^*\mathbb{Z}^{\rm sym}_\kappa(\theta) = {^*\mathbb{Z}}(\theta|x^{-\kappa})$ is a family of groups, and $^*\mathbb{Z}^{\rm sym}_{\kappa'}(\theta) = 0$ for all $\kappa' > \kappa$.

d. If $\theta \Bumpeq \eta$ by $A \in \mathrm{PGL}_2(\mathbb{Z})$ then we have the following analogue of Theorem 24): for all ${}^*m \in {}^*\mathbb{Z}(\theta|x^{-\kappa})$, $|A({}^*m)|_{A(\theta),\kappa} = |{}^*m|_{\theta,\kappa}$.

We may also consider nonstandard exponents: for any ${}^*\kappa \in {}^*\mathbb{R}_+$, ${}^*\kappa > 1$,

$${}^*\mathbb{Z}(\theta|x^{-{}^*\kappa}) \subset \bigcap_{{}^*\kappa>\kappa>1} {}^*\mathbb{Z}(\theta|x^{-\kappa}),$$

equipped with its associated norm function $|\cdot|_{\theta,{}^*\kappa}$ with which we may define the ${}^*\kappa$ symmetric diophantine approximations ${}^*\mathbb{Z}^{\mathrm{sym}}_{{}^*\kappa}(\theta) = \{{}^*\mathbb{Z}^{\mathrm{sym}}_{\nu,{}^*\kappa}(\theta)\}$. Note that if $\theta \notin \mathfrak{W}_\infty$ then ${}^*\mathbb{Z}^{\mathrm{sym}}_{{}^*\kappa}(\theta) = 0$ for all ${}^*\kappa$ infinite nonstandard. In general we have

$${}^*\mathbb{Z}(\theta|x^{-1}) = \bigcup_{{}^*\kappa>1} {}^*\mathbb{Z}^{\mathrm{sym}}_{{}^*\kappa}(\theta).$$

We refer to ${}^*\mathbb{Z}(\theta)$, with the family $\{({}^*\mathbb{Z}(\theta|x^{-{}^*\kappa}), |\cdot|_{\theta,{}^*\kappa})\}$ as a **Frechet Lorentzian groupology**: for any ${}^*m \in {}^*\mathbb{Z}(\theta)$, $|{}^*m|_{\theta,{}^*\kappa} = 0$ for all ${}^*\kappa \Leftrightarrow {}^*m = 0$. We call an ideological homomorphism

$$f : {}^*\mathbb{Z}(\theta) \to {}^*\mathbb{Z}(\eta)$$

a **Frechet Lorentzian isometry** or simply an **isometry** if it preserves the Frechet Lorentzian norms.

Theorem 25. $\theta \Bumpeq \eta \Rightarrow {}^*\mathbb{Z}(\theta) \cong {}^*\mathbb{Z}(\eta)$ *by isometric ideological isomorphism.*

Proof. The proof of Theorem 24 follows through identically to show that $A \in \mathrm{PGL}_2\mathbb{Z}$ acts by Frechet Lorentzian isometries. ☐

Conjecture. $\theta \Bumpeq \eta \Leftrightarrow {}^*\mathbb{Z}(\theta) \cong {}^*\mathbb{Z}(\eta)$ *by an isometric ideological isomorphism.*

8. IDEOLOGICAL ARITHMETIC OF $\mathcal{O}$-APPROXIMATION GROUPS

We refer to §5 of [13] for notation and definitions. Let $K/\mathbb{Q}$ be a finite extension of degree d and let $\mathbb{K} \cong \mathbb{R}^r \times \mathbb{C}^s \cong \mathbb{R}^d$ be the Minkowski space: the archimedean part of the K-adeles, a finite-dimensional $\mathbb{R}$-algebra. We regard $\mathbb{R} \subset \mathbb{K}$ via the diagonal embedding. Denote by $\mathrm{N} : {}^*\mathbb{K} \to {}^*\mathbb{R}$ the norm map: ${}^*\boldsymbol{z} = ({}^*z_i) \mapsto \mathrm{N}({}^*\boldsymbol{z}) = {}^*z_1 \cdots {}^*z_d$.

K-Tropical Semi-ring

Consider the ring ${}^*\mathbb{K}_{\mathrm{fin}} \subset {}^*\mathbb{K}$ of elements all of whose coordinates are bounded. The group of units is the subgroup ${}^*\mathbb{K}^\times_{\mathrm{fin}}$ of elements all of whose coordinates are non-infinitesimal and non-infinite. The multiplicative quotient

$${}^*\mathbb{PK} = {}^*\mathbb{K}/{}^*\mathbb{K}^\times_{\mathrm{fin}}$$

is partially ordered and directed along the coordinates. We will denote elements of ${}^*\mathbb{PK}$ by $\boldsymbol{\mu}$. There is a diagonal inclusion ${}^*\mathbb{PR} \hookrightarrow {}^*\mathbb{PK}$; the image of μ will be denoted μ (not bold).

As in the case of $K = \mathbb{Q}$, ${}^*\mathbb{PK}$ has the structure of a tropical semi-ring with respect to the induced product, and the sum defined

$$\boldsymbol{\mu} + \boldsymbol{\mu}' = (\mu_1 + \mu_1', \dots, \mu_d + \mu_d').$$

Note that $\boldsymbol{\mu} + \boldsymbol{\mu}'$ is the least element greater than $\boldsymbol{\mu}, \boldsymbol{\mu}'$. Elements of ${}^*K$ act by multiplication on the left of ${}^*\mathbb{PK}$, respecting in the style of Proposition 4 the tropical structure. The norm (product of coordinates) map induces a multiplicative map $\mathrm{N} : {}^*\mathbb{PK} \to {}^*\mathbb{PR}$.

Note that ${}^*\mathbb{P}\mathbb{C} := {}^*\mathbb{C}/{}^*\mathbb{C}^\times_{\mathrm{fin}}$ is isomorphic to ${}^*\mathbb{P}\mathbb{R}$ since ${}^*\mathbb{C}^\times_{\mathrm{fin}} \supset \mathbb{S}^1$. Thus the non archimedean valuation $\langle\cdot\rangle$, extended in the obvious way to ${}^*\mathbb{C}$, takes values in ${}^*\mathbb{P}\mathbb{R}$. We have

$$ {}^*\mathbb{P}\mathbb{K} \cong ({}^*\mathbb{P}\mathbb{R})^r \times ({}^*\mathbb{P}\mathbb{C})^s \cong {}^*\mathbb{P}\mathbb{R}^{r+s}, $$

so that every element $\boldsymbol{\mu}$ may be written in coordinate form $(\mu_1, \dots, \mu_{r+s})$.

Let $\langle\cdot\rangle : {}^*\mathbb{K} \to {}^*\mathbb{P}\mathbb{K}$ the the canonical projection. Let ${}^*\mathbb{K}_\varepsilon \subset {}^*\mathbb{K}_{\mathrm{fin}}$ be the (neither maximal nor prime) ideal of elements all of whose coordinates are infinitesimal and denote ${}^*\mathbb{P}\mathbb{K}_\varepsilon = {}^*\mathbb{K}_\varepsilon/{}^*\mathbb{K}^\times_{\mathrm{fin}}$. Define also

$$ {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)} = \{\boldsymbol{\mu} \in {}^*\mathbb{P}\mathbb{K} |\ \mathrm{N}(\boldsymbol{\mu}) \in {}^*\mathbb{P}\mathbb{R}_\varepsilon\} \supset {}^*\mathbb{P}\mathbb{K}_\varepsilon. $$

While ${}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$ is not closed with respect to $+$, it is downwardly closed with respect to the partial order ($\boldsymbol{\mu} \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$ and $\boldsymbol{\mu}' < \boldsymbol{\mu}$ implies $\boldsymbol{\mu}' \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$) and moreover, each interval $(-\infty, \boldsymbol{\mu}] = \{\boldsymbol{\mu}' |\ \boldsymbol{\mu}' \leq \boldsymbol{\mu}\} \subset {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$ is a sub tropical semi ring. Note that if we define

$$ {}^*\mathbb{P}\mathbb{K}_{\mathrm{Tr}(\boldsymbol{\varepsilon})} = \{\boldsymbol{\mu} |\ \mathrm{Tr}(\boldsymbol{\mu}) := \mu_1 + \cdots + \mu_{r+s} = \max(\mu_i) \in {}^*\mathbb{P}\mathbb{R}_\varepsilon\}, $$

then ${}^*\mathbb{P}\mathbb{K}_{\mathrm{Tr}(\boldsymbol{\varepsilon})} = {}^*\mathbb{P}\mathbb{K}_\varepsilon$.

Let $\langle\!\langle\cdot\rangle\!\rangle : {}^*\mathbb{P}\mathbb{K} \to {}^*\mathbb{P}\mathbb{R}$ be the map induced by composing the euclidean norm $\|\cdot\|$ on ${}^*\mathbb{K}$ (valued in ${}^*\mathbb{R}_+$) with $\langle\cdot\rangle$: in other words, $\langle\!\langle\cdot\rangle\!\rangle = \langle\|\cdot\|\rangle$. In particular, we have

$$ \langle\!\langle\boldsymbol{\mu}\rangle\!\rangle = \mathrm{Tr}(\boldsymbol{\mu}) = \mu_1 + \cdots + \mu_d = \max \mu_i. $$

K-Ideology

For each $\boldsymbol{\mu} \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$ define

$$ {}^*\mathcal{O}^{\boldsymbol{\mu}} = \{{}^*\alpha \in {}^*\mathcal{O} |\ {}^*\alpha \cdot \boldsymbol{\mu} \in {}^*\mathbb{P}\mathbb{K}_\varepsilon\} \subset {}^*\mathcal{O}. $$

If $\boldsymbol{\mu} < \boldsymbol{\mu}'$ then ${}^*\mathcal{O}^{\boldsymbol{\mu}} \supset {}^*\mathcal{O}^{\boldsymbol{\mu}'}$ however if $\boldsymbol{\mu}$ and $\boldsymbol{\mu}'$ are unrelated, the associated groups are unrelated by inclusion. The fine growth subfiltration ${}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}$ is defined by ${}^*\alpha\cdot\boldsymbol{\mu} < \boldsymbol{\iota}$ where $\boldsymbol{\iota} \in {}^*\mathbb{P}\mathbb{K}_\varepsilon$, a subgroup of ${}^*\mathcal{O}^{\boldsymbol{\mu}}$.

The use of elements $\boldsymbol{\mu} \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$ to index growth is natural in this context as there exist elements ${}^*\alpha \in {}^*\mathcal{O}$ having $\mathbb{K}$-coordinates $({}^*\alpha_1, \dots, {}^*\alpha_d)$ exhibiting inhomogeneous growth: having a bounded non-infinitesimal coordinate, an infinite coordinate and an infinitesimal coordinate. For example, if α is a Salem number, the class of the sequence $\{\alpha^i\}$ has a coordinate lying on ${}^*\mathbb{S}^1$, an infinitesimal coordinate and an infinite coordinate.

Now given $\boldsymbol{z} \in \mathbb{K}$ and $\boldsymbol{\nu} \in {}^*\mathbb{P}\mathbb{K}_\varepsilon$ we define ${}^*\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{z}) \subset {}^*\mathcal{O}(\boldsymbol{z})$ as the subgroup of ${}^*\alpha$ for which $\langle\boldsymbol{\varepsilon}({}^*\alpha)\rangle \leq \boldsymbol{\nu}$, where $\boldsymbol{\varepsilon}({}^*\alpha) = {}^*\alpha\boldsymbol{z} - {}^*\alpha^\perp$. We have

$$ \langle\boldsymbol{\varepsilon}({}^*\alpha + {}^*\alpha')\rangle \leq \langle\boldsymbol{\varepsilon}({}^*\alpha)\rangle + \langle\boldsymbol{\varepsilon}({}^*\alpha')\rangle \leq \boldsymbol{\nu} $$

so ${}^*\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{z})$ is a group. Here we note that we can extend the definition of ${}^*\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{z})$ to indices $\boldsymbol{\nu} \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$, since $(-\infty, \boldsymbol{\nu})$ is a sub tropical semi-ring. We have thus defined the K-ideology structure

$$ {}^*\mathcal{O}(\boldsymbol{z}) = \left\{{}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\boldsymbol{z})\right\}, \quad {}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\boldsymbol{z}) = {}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]} \cap {}^*\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{z}) $$

for $\boldsymbol{z} \in \mathbb{K}$ and $\boldsymbol{\mu}, \boldsymbol{\nu}, \boldsymbol{\iota} \in {}^*\mathbb{P}\mathbb{K}_{\mathrm{N}(\varepsilon)}$. Note that we have the compatibility ${}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \subset {}^*\mathcal{O}^{\mu}_{\nu}(\theta)$.

K-Nonvanishing Spectra

The K-nonvanishing spectrum is

$$\mathrm{Spec}_{\mathcal{O}}(\boldsymbol{z}) = \{(\boldsymbol{\mu},\boldsymbol{\nu}) |\, {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \neq \boldsymbol{0}\} \subset {}^\star\mathbb{P}\mathbb{K}^2_{\mathrm{N}(\varepsilon)}.$$

Note first that trivially

Proposition 24. *Let* $\theta \in \mathbb{R}$. *If* $\boldsymbol{\mu} < \boldsymbol{\nu} \in {}^\star\mathbb{P}\mathbb{K}_\varepsilon$ *then* ${}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\theta) \neq 0$.

Proof. Let $\mu < \nu \in {}^\star\mathbb{P}\mathbb{R}_\varepsilon$ be such that $\boldsymbol{\mu} < \mu < \nu < \boldsymbol{\nu}$. Then $0 \neq {}^*\mathbb{Z}^{\mu}_{\nu}(\theta) \subset {}^*\mathcal{O}^{\mu}_{\nu}(\theta) \subset {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\theta)$. ☐

Recall that $\theta \in \mathbb{R}$ is called a Pisot-Vijayaraghavan number if it is a real algebraic integer greater than 1 for which all of its conjugates have absolute value < 1. In what follows, we denote $P_{K,\theta}(X) = \prod(X - \theta_\tau)$ where the product is over the archimedean places τ of the field K and $\theta_\tau := \tau(\theta)$.

Theorem 26. *Let* $\theta \in \mathbb{R}$ *be a Pisot-Vijayaraghavan number with* $\theta \in \tau(\mathcal{O})$ *for some place* τ *and for which* $P_{K,\theta}(X)$ *is the minimal polynomial of* θ. *Then there exists* $\boldsymbol{\mu} \in {}^\star\mathbb{P}\mathbb{K}_\varepsilon$ *such that* ${}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\mu}}(\theta) \neq 0$.

Proof. Let $\vartheta \in \mathcal{O}$ be such that $\vartheta_{\tau_1} = \theta$, so that $|\vartheta_{\tau_i}| < 1$ for $i = 2, \dots d$. Let ${}^*\alpha \in {}^*\mathcal{O}$ be the class associated to the sequence $\vartheta, \vartheta^2, \vartheta^3, \dots$ and let ${}^*\alpha^\perp = \vartheta {}^*\alpha \in {}^*\mathcal{O}$. Then ${}^*\alpha \in {}^*\mathcal{O}(\theta)$ with dual ${}^*\alpha^\perp$: indeed ${}^*\alpha\theta - {}^*\alpha^\perp$ is the class of the vector sequence

$$\left\{\left(0, \vartheta^n_{\tau_2}(\theta - \vartheta_{\tau_2}), \dots, \vartheta^n_{\tau_d}(\theta - \vartheta_{\tau_d})\right)\right\}_{n=1}^\infty$$

which is infinitesimal since $|\vartheta_{\tau_i}| < 1$ for $i = 2, \dots d$. Let $\boldsymbol{\mu} \in {}^\star\mathbb{P}\mathbb{K}_\varepsilon$ be the class of ${}^*\alpha\theta - {}^*\alpha^\perp$. Then

$${}^*\alpha \cdot \boldsymbol{\mu} = (0, {}^*\alpha_{\tau_2} \cdot \mu_2, \dots, {}^*\alpha_{\tau_d} \cdot \mu_d) \in {}^\star\mathbb{P}\mathbb{K}_\varepsilon$$

since the components ${}^*\alpha_{\tau_2}, \dots, {}^*\alpha_{\tau_d}$ are themselves infinitesimal. Thus ${}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\mu}}(\theta) \neq 0$. ☐

Theorem 26 reveals that there are infinitely many antiprimes (quadratic Pisot Vijayaraghavan numbers) which possess a non-trivial flat spectra provided that we expand the field of approximants to one minimally containing them. In particular, such numbers cease to be antiprime, a phenomenon which may be described as the "splitting" of the nonvanishing spectrum.

K-Ideological Arithmetic

We have the following exact analogue of Theorem 11:

Theorem 27 (K-Ideological Arithmetic)**.** *Let* $\boldsymbol{z}, \boldsymbol{w} \in \mathbb{K}$. *Then*

$$
{}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\boldsymbol{z}) \cdot {}^*\mathcal{O}^{\boldsymbol{\nu}[\boldsymbol{\lambda}]}_{\boldsymbol{\mu}}(\boldsymbol{w}) \subset {}^*\mathcal{O}^{\boldsymbol{\mu}\cdot\boldsymbol{\nu}[\boldsymbol{\iota}\cdot\boldsymbol{\lambda}]}_{\boldsymbol{\iota}+\boldsymbol{\lambda}}(\boldsymbol{z}\boldsymbol{w}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}\cdot\boldsymbol{\nu}[\boldsymbol{\iota}\cdot\boldsymbol{\lambda}]}_{\boldsymbol{\iota}+\boldsymbol{\lambda}}(\boldsymbol{z}+\boldsymbol{w}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}\cdot\boldsymbol{\nu}[\boldsymbol{\iota}\cdot\boldsymbol{\lambda}]}_{\boldsymbol{\iota}+\boldsymbol{\lambda}}(\boldsymbol{z}-\boldsymbol{w}). \tag{29}
$$

Proof. If ${}^*\alpha \in {}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\boldsymbol{z})$ and ${}^*\beta \in {}^*\mathcal{O}^{\boldsymbol{\nu}[\boldsymbol{\lambda}]}_{\boldsymbol{\mu}}(\boldsymbol{w})$ then

$$({}^*\alpha + {}^*\beta) \cdot \boldsymbol{z}\boldsymbol{w} = (({}^*\alpha_1 + {}^*\beta_1)z_1w_1, \dots, ({}^*\alpha_d + {}^*\beta_d)z_dw_d)$$

and

$${}^*\alpha {}^*\beta \cdot \boldsymbol{z}\boldsymbol{w} = ({}^*\alpha_1 {}^*\beta_1 z_1 w_1, \dots, {}^*\alpha_d {}^*\beta_d z_d w_d).$$

The proof proceeds as in that of Theorem 11, implemented along the coordinates of ${}^*\mathbb{K}$. ☐

The remarks following Theorem 11 apply just as well to $\mathcal{O}$-ideological arithmetic. Here however non principal ideals are absent, not surprising since the definition of $\mathcal{O}$-diophantine approximation groups is made with regard to single elements of $\mathbb{K}$. In §15 we will produce the analogues of (classes) of two generator ideals by "decoupling" numerator denominator pairs. Non principal ideals also appear naturally as dual diophantine approximations of vectors, see §9.

For $\boldsymbol{z}, \boldsymbol{w} \in \mathbb{K}$, we write

$$\boldsymbol{z}_{\boldsymbol{\mu}}\textcircled{$\mathcal{O}$}_{\boldsymbol{\nu}}\boldsymbol{w} \tag{30}$$

when the product is nontrivial with respect to $\mathcal{O}$ and the parameters $\boldsymbol{\mu}, \boldsymbol{\nu}$. Thus when $\mathcal{O} = \mathbb{Z}$, $\textcircled{$\mathbb{Z}$} = \textcircled{$\wedge$}$. The symbol

$$^{*}\alpha\ _{\boldsymbol{\mu}}\textcircled{$\boldsymbol{z}\boldsymbol{w}$}_{\boldsymbol{\nu}}\,^{*}\beta$$

will indicate that the product $^{*}\alpha^{*}\beta$ is defined as one of diophantine approximations, subject to the condition that $^{*}\alpha \in {}^{*}\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z})$ resp. $^{*}\beta \in {}^{*}\mathcal{O}^{\boldsymbol{\nu}}_{\boldsymbol{\mu}}(\boldsymbol{w})$. The notions of $\mathcal{O}$-fast, $\mathcal{O}$-slow and $\mathcal{O}$-flat divisors are defined as in §5. When $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ are not related by the order, we say that the factors are **oscillatory divisors**.

Let $\mathrm{PGL}_2(\mathcal{O})$ be the projective linear group with entries in $\mathcal{O}$. Then $\mathrm{PGL}_2(\mathcal{O})$ acts on $\mathbb{K}$ and we write

$$\boldsymbol{z} \Bumpeq_K \boldsymbol{z}'$$

if there exists $A \in \mathrm{PGL}_2(\mathcal{O})$ such that $A(\boldsymbol{z}) = \boldsymbol{z}'$.

Theorem 28. *If* $\boldsymbol{z} \Bumpeq_K \boldsymbol{z}'$ *by* $A \in \mathrm{PGL}_2(\mathcal{O})$ *then* A *induces an ideological isomorphism*

$$A : {}^{*}\mathcal{O}(\boldsymbol{z}) \longrightarrow {}^{*}\mathcal{O}(\boldsymbol{z}').$$

If in addition we have $\boldsymbol{w} \in \mathbb{K}$ *and* $B \in \mathrm{PGL}_2(\mathcal{O})$ *then*

$$\boldsymbol{z}_{\boldsymbol{\mu}}\textcircled{$\mathcal{O}$}_{\boldsymbol{\nu}}\boldsymbol{w} \iff A(\boldsymbol{z})_{\boldsymbol{\mu}}\textcircled{$\mathcal{O}$}_{\boldsymbol{\nu}}B(\boldsymbol{w}).$$

Proof. Same idea as the proof of Theorem 3, implemented along place coordinates. ☐

Proposition 25. *Suppose that* $K/\mathbb{Q}$ *is Galois and* $\sigma \in \mathrm{Gal}(K/\mathbb{Q})$. *Then*

$$\boldsymbol{z}_{\boldsymbol{\mu}}\textcircled{$\mathcal{O}$}_{\boldsymbol{\nu}}\boldsymbol{w} \iff \sigma(\boldsymbol{z})_{\sigma(\boldsymbol{\mu})}\textcircled{$\mathcal{O}$}_{\sigma(\boldsymbol{\nu})}\sigma(w)$$

and

$$^{*}\alpha\ _{\boldsymbol{\mu}}\textcircled{$\boldsymbol{z}\boldsymbol{w}$}_{\boldsymbol{\nu}}\,^{*}\beta \iff \sigma(^{*}\alpha)_{\sigma(\boldsymbol{\mu})}\textcircled{$\sigma(\boldsymbol{z})\sigma(\boldsymbol{w})$}_{\sigma(\boldsymbol{\nu})}\sigma(^{*}\beta).$$

In particular, an elements status as an $\mathcal{O}$*-fast,* $\mathcal{O}$*-slow,* $\mathcal{O}$*-flat or* $\mathcal{O}$*-oscillatory divisor is preserved by the action of* $\sigma \in \mathrm{Gal}(K/\mathbb{Q})$.

Proof. Let $\sigma \in \mathrm{Gal}(K/\mathbb{Q})$, then

$$\sigma\left({}^{*}\mathcal{O}^{\boldsymbol{\mu}[\iota]}_{\boldsymbol{\nu}}(\boldsymbol{z})\right) = {}^{*}\mathcal{O}^{\sigma(\boldsymbol{\mu})[\sigma(\iota)]}_{\sigma(\boldsymbol{\nu})}\big(\sigma(\boldsymbol{z})\big).$$

It follows immediately that σ respects the K-ideological product as indicated in the statement of the Proposition. ☐

If we take if $\boldsymbol{z} = \boldsymbol{\theta} = (\theta, \dots, \theta)$ then the trace map $\mathrm{Tr} : \mathbb{K} \to \mathbb{R}$ defines a well-defined homomorphism of groups $\mathrm{Tr} : {}^{*}\mathcal{O}(\boldsymbol{\theta}) \to {}^{*}\mathbb{Z}(\theta)$. In addition, we have a well-defined map of projective classes $\mathrm{Tr} : {}^{\star}\mathbb{PK}_{\varepsilon} \longrightarrow {}^{\star}\mathbb{PR}_{\varepsilon}$ so that

$$\mathrm{Tr}\left({}^{*}\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{\theta})\right) \subset {}^{*}\mathbb{Z}_{\mathrm{Tr}(\boldsymbol{\nu})}(\theta).$$

Note that the trace map does *not* map ${}^{*}\mathcal{O}^{\boldsymbol{\mu}}(\boldsymbol{\theta})$ to ${}^{*}\mathbb{Z}^{\mathrm{Tr}(\boldsymbol{\mu})}(\theta)$.

On the other hand, the norm map, as we have seen, induces $\mathrm{N} : {}^{\star}\mathbb{PK}_{\mathrm{N}(\varepsilon)} \longrightarrow {}^{\star}\mathbb{PR}_{\varepsilon}$, however it does not define a map from ${}^{*}\mathcal{O}_{\boldsymbol{\nu}}(\boldsymbol{\theta})$ to ${}^{*}\mathbb{Z}_{\mathrm{N}(\boldsymbol{\nu})}(\theta)$. Instead it yields a map of fine growth filtrations

$$\mathrm{N} : {}^{*}\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]} \longrightarrow {}^{*}\mathbb{Z}^{\mathrm{N}(\boldsymbol{\mu})[\mathrm{N}(\boldsymbol{\iota})]}.$$

As an immediate corollary, we may deduce that for $\boldsymbol{\mu}$ to be a growth rate for elements of ${}^{*}\mathcal{O}$, the product of the infinitesimal coordinates must dominate the product of the non infinitesimal coordinates:

Proposition 26. *If* $\boldsymbol{\mu} \in {}^{\star}\mathbb{PK}$ *and* ${}^{*}\mathcal{O}^{\boldsymbol{\mu}} \neq 0$ *then* $\boldsymbol{\mu} \in {}^{\star}\mathbb{PR}_{\mathrm{N}(\varepsilon)}$.

Proof. This follows since every element of ${}^{*}\mathbb{Z}$ is infinite, so $\mathrm{N}({}^{*}\boldsymbol{\alpha}) \cdot \mathrm{N}(\boldsymbol{\mu}) \in {}^{\star}\mathbb{PR}_{\varepsilon}$ can only occur if $\mathrm{N}(\boldsymbol{\mu})$ is infinitesimal. □

From the above paragraphs, we deduce the following broad principle:

> Growth is multiplicative and not additive. Decay is additive but not multiplicative.

In general, if we seek to return to the ground ring ${}^{*}\mathbb{Z}$ using either the norm or the trace, one of the ideological parameters must be sacrificed. Nevertheless, there exist specific situations when the anomalous parameter can be controlled.

Proposition 27. *Let* $K/\mathbb{Q}$ *be of degree* 2 *and let* σ *be the nontrivial element of its Galois group. Then for any* $\boldsymbol{z} \in \mathbb{K}$,

$$\mathrm{N}\left({}^{*}\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\sigma(\boldsymbol{\mu})}(\boldsymbol{z})\right) \subset {}^{*}\mathbb{Z}^{\mathrm{N}(\boldsymbol{\mu})[\mathrm{N}(\boldsymbol{\iota})]}_{\mathrm{Tr}(\boldsymbol{\iota})}\big(\mathrm{N}(\boldsymbol{z})\big).$$

Proof. Let ${}^{*}\boldsymbol{\alpha} \in {}^{*}\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\sigma(\boldsymbol{\mu})}(\boldsymbol{z})$. Write ${}^{*}\boldsymbol{\alpha} = ({}^{*}\alpha_1, {}^{*}\alpha_2), {}^{*}\boldsymbol{\alpha}^{\perp} = ({}^{*}\alpha_1^{\perp}, {}^{*}\alpha_2^{\perp})$ as well as $\boldsymbol{\varepsilon}({}^{*}\boldsymbol{\alpha}) = (\varepsilon({}^{*}\alpha_1), \varepsilon({}^{*}\alpha_2))$. It is immediate that $\mathrm{N}({}^{*}\boldsymbol{\alpha}) = {}^{*}\alpha_1{}^{*}\alpha_2 \in {}^{*}\mathbb{Z}^{\mathrm{N}(\boldsymbol{\mu})[\mathrm{N}(\boldsymbol{\iota})]}$. On the other hand,

$$\begin{aligned} \mathrm{N}({}^{*}\boldsymbol{\alpha}) \cdot \mathrm{N}(\boldsymbol{z}) - \mathrm{N}({}^{*}\boldsymbol{\alpha}^{\perp}) &= \mathrm{N}(\boldsymbol{\varepsilon}({}^{*}\boldsymbol{\alpha})) + {}^{*}\alpha_1^{\perp}({}^{*}\alpha_2 z_2 - {}^{*}\alpha_2^{\perp}) + {}^{*}\alpha_2^{\perp}({}^{*}\alpha_1 z_1 - {}^{*}\alpha_1^{\perp}) \\ &= \mathrm{N}(\boldsymbol{\varepsilon}({}^{*}\boldsymbol{\alpha})) + {}^{*}\alpha_1^{\perp} \cdot \varepsilon({}^{*}\alpha_2) + {}^{*}\alpha_2^{\perp} \cdot \varepsilon({}^{*}\alpha_1). \end{aligned} \tag{31}$$

Since ${}^{*}\boldsymbol{\alpha} \in {}^{*}\mathcal{O}_{\sigma(\boldsymbol{\mu})}(\boldsymbol{z})$, the image of (31) by $\langle\cdot\rangle$ belongs to ${}^{\star}\mathbb{PR}_{\varepsilon}$. Since

$$\langle \mathrm{N}(\boldsymbol{\varepsilon}({}^{*}\boldsymbol{\alpha}))\rangle < \langle {}^{*}\alpha_1^{\perp} \cdot \varepsilon({}^{*}\alpha_2)\rangle, \langle {}^{*}\alpha_2^{\perp} \cdot \varepsilon({}^{*}\alpha_1)\rangle$$

we may disregard $\langle \mathrm{N}(\boldsymbol{\varepsilon}({}^{*}\boldsymbol{\alpha}))\rangle$, and therefore the image of (31) by $\langle\cdot\rangle$ is bounded by $\mathrm{Tr}(\boldsymbol{\iota})$. □

Note 9. In view of the nature of the image growth-decay indices occurring in Proposition 27, we cannot use the norm to push products down of the form presented in Theorem 27.

There is a similar sort of result for the trace. Given $\boldsymbol{\mu} \in {}^*\mathbb{PK}$, define the **lower trace** to be

$$\mathrm{tr}(\boldsymbol{\mu}) := \min \mu_i .$$

Note that if $\mu \in {}^*\mathbb{PR}$ then $\mathrm{tr}(\mu) = \mathrm{Tr}(\mu) = \mu$.

Proposition 28. *Let $K/\mathbb{Q}$ be of degree d. Then for any $\theta \in \mathbb{R}$,*

$$\mathrm{Tr}\left({}^*\mathcal{O}^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\theta)\right) \subset {}^*\mathbb{Z}^{\mathrm{tr}(\boldsymbol{\mu})[\mathrm{Tr}(\boldsymbol{\iota})]}_{\mathrm{Tr}(\boldsymbol{\nu})}(\theta).$$

If $\boldsymbol{\nu} = -\infty$, the result is valid for θ replaced by $\gamma \in K \subset \mathbb{K}$.

Proof. We have already observed that the trace map preserves decay for $\theta \in \mathbb{R}$. If the decay is $-\infty$ then the trace map preserves the decay for $\gamma \in K$. On the other hand, the inequality ${}^*\alpha \cdot \boldsymbol{\mu} < \boldsymbol{\iota}$ may be rewritten $({}^*\alpha_1 \cdot \mu_1, \cdots, {}^*\alpha_d \cdot \mu_d) < (\iota_1, \ldots, \iota_d)$. It follows that

$$\mathrm{Tr}({}^*\alpha) \cdot \mathrm{tr}(\boldsymbol{\mu}) = (\alpha_1 + \cdots + \alpha_d) \cdot \min(\mu_i) < \iota_1 + \cdots + \iota_d = \mathrm{Tr}(\boldsymbol{\iota}).$$

□

Corollary 14. *If $\mu, \nu \in {}^*\mathbb{PR}_{\varepsilon}$, $\theta, \eta \in \mathbb{R}$ and ${}^*\alpha \ {}_{\mu}\textcircled{\theta\eta}_{\nu} {}^*\beta$ then $\mathrm{Tr}({}^*\alpha) \ {}_{\mu}\textcircled{\theta\eta}_{\nu} \mathrm{Tr}({}^*\beta)$. If $\nu = -\infty$ the result holds for $\gamma, \delta \in K$.*

Thus the trace map respects the *ideological arithmetic* of classical (principal) ideals.

K-Classification

Diophantine approximation by the K-integers $\mathcal{O}$, as formulated in [13], is *global*, since it is performed with respect to all the archimedean places at once – in contrast to the classical notion of Diophantine approximation by elements of K [5], which is *local*, framed with respect to a fixed archimedean place. It is therefore reasonable to define K-versions of the usual linear classification of the reals as described in §3. Note that the classification theory of Koksma, though based on approximations by algebraic numbers, is not field specific, and is therefore not relevant to the considerations of this section.

We say that $\boldsymbol{z} \in \mathbb{K} - K$ is

- ***K*-badly approximable** if ${}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\mu}}(\boldsymbol{z}) = 0$ for all $\boldsymbol{\mu} \in {}^*\mathbb{PK}_{\varepsilon}$.
- ***K*-well approximable** if it is not K-badly approximable.
- ***K*-very well approximable** of exponent κ if there exists $\boldsymbol{\mu} \in {}^*\mathbb{PK}_{\varepsilon}$ such that

$$\bigcap_{\lambda \in [1,\kappa)} {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\mu}^{\lambda}} \neq 0.$$

- ***K*-Liouville** if it is K-very well approximable of exponent κ for all $\kappa > 1$.

Denote these classes by $\mathfrak{B}(K)$, $\mathfrak{W}(K)$, $\mathfrak{W}_{\kappa}(K)$ and $\mathfrak{W}_{\infty}(K)$, any one of which is denoted $\mathfrak{C}(K)$; the $K = \mathbb{Q}$ counterpart is simply denoted $\mathfrak{C}$. Note that for any class $\mathfrak{C}$ there exists $\theta \in \mathfrak{C}$ such that $\theta \notin \sigma(K)$ for every archimedean place σ of K.

Theorem 29. *Let $\theta \in \mathfrak{C}$ be such that $\theta \notin \sigma(K)$ for every archimedean place σ of K. Then $\theta \in \mathfrak{C}(K)$. In particular, the class $\mathfrak{C}(K)$ is nontrivial for all $K/\mathbb{Q}$ of finite degree.*

Proof. The only nontrivial case is when $\mathfrak{C}(K)=\mathfrak{B}(K)$. Suppose that there exists $\boldsymbol{\mu}\in{}^{\star}\mathbb{PK}_{\varepsilon}$ with ${}^{*}\mathcal{O}^{\mu}_{\boldsymbol{\mu}}(\theta)\neq 0$. Since $\theta\notin\sigma(K)$ for every archimedean place σ, none of the coordinates of $\boldsymbol{\mu}$ are equal to $-\infty$. Therefore, we may find a diagonal element $\mu=(\mu,\dots,\mu)$, where $\mu\in{}^{\star}\mathbb{PR}_{\varepsilon}$, such that $\boldsymbol{\mu}<\mu<1$ in which ${}^{*}\alpha\cdot\mu\in\mathbb{PK}_{\varepsilon}$. For such and element we have *a fortiori* that $\boldsymbol{\nu}({}^{*}\alpha)\leq\mu$ so that ${}^{*}\alpha\in{}^{*}\mathcal{O}^{\mu}_{\mu}(\theta)\neq 0$ as well. Applying Proposition 28, and using the fact that $\mathrm{Tr}(\mu)=\mathrm{tr}(\mu)=\mu$ we obtain $0\neq\mathrm{Tr}({}^{*}\alpha)\in{}^{*}\mathbb{Z}^{\mu}_{\mu}(\theta)$, which is a contradiction. $\square$

The K-classes defined above are Galois natural: if $K/\mathbb{Q}$ is a Galois extension, then for any K-class $\mathfrak{C}(K)$ and $\sigma\in\mathrm{Gal}(K/\mathbb{Q})$ we have $\sigma(\mathfrak{C}(K))=\mathfrak{C}(K)$, acting as the identity on $\mathfrak{C}\subset\mathfrak{C}(K)$. If $\theta\in\mathfrak{C}$ but $\theta\notin\mathfrak{C}(K)$ then we say that θ **splits** in K. For example,Theorem 26 says that any quadratic Pisot-Vijayaraghavan number θ with a conjugate in K splits in K. The factorization symbols $\Uparrow_K$, $\Downarrow_K$, $\Updownarrow_K$ and $[\!]_K$ have the expected meanings in the K-context, which are Galois natural in a sense which the reader may formulate.

9. MATRIX IDEOLOGICAL ARITHMETIC

Let Θ be a real matrix of size $r\times s$. In [13] we defined the **inhomogeneous diophantine approximation group** of Θ to be

$$ {}^{*}\mathbb{Z}^{s}(\Theta)=\{{}^{*}\boldsymbol{n}\in{}^{*}\mathbb{Z}^{s}|\ \exists{}^{*}\boldsymbol{n}^{\perp}\in{}^{*}\mathbb{Z}^{r}\text{ s.t. }\varepsilon({}^{*}\boldsymbol{n}):=\Theta{}^{*}\boldsymbol{n}-{}^{*}\boldsymbol{n}^{\perp}\ \in{}^{*}\mathbb{R}^{r}_{\varepsilon}\}. $$

The corresponding **homogeneous diophantine approximation group** was defined by

$$ {}^{*}\widetilde{\mathbb{Z}}^{s}(\Theta)=\mathrm{Ker}(\perp)=\{{}^{*}\boldsymbol{n}\in{}^{*}\mathbb{Z}^{s}(\Theta)|\ \Theta{}^{*}\boldsymbol{n}\in{}^{*}\mathbb{R}^{r}_{\varepsilon}\}<{}^{*}\mathbb{Z}^{s}(\Theta). $$

Thus, if ${}^{*}\mathbb{Z}^{r}(\Theta)^{\perp}$ denotes the group of duals then ${}^{*}\mathbb{Z}^{s}(\Theta)/{}^{*}\widetilde{\mathbb{Z}}^{s}(\Theta)\cong{}^{*}\mathbb{Z}^{r}(\Theta)^{\perp}$. Note that if Θ is square invertible then ${}^{*}\widetilde{\mathbb{Z}}^{s}(\Theta)=0$.

In this section we will develop ideological arithmetic for the groups ${}^{*}\mathbb{Z}^{s}(\Theta)$. First, for ${}^{*}\boldsymbol{n}=({}^{*}n_1,\dots,{}^{*}n_s)\in{}^{*}\mathbb{Z}^{s}$, the **house norm** is defined

$$ \overline{|{}^{*}\boldsymbol{n}|}:=\max_{j=1,\dots,s}|{}^{*}n_j| \tag{32} $$

and $\mu({}^{*}\boldsymbol{n}):=\langle\overline{|{}^{*}\boldsymbol{n}|}\rangle^{-1}\in{}^{\star}\mathbb{PR}$. Then for $\mu\in{}^{\star}\mathbb{PR}_{\varepsilon}$ the set

$$ ({}^{*}\mathbb{Z}^{s})^{\mu}=\left\{{}^{*}\boldsymbol{n}\middle|\ \mu^{1/s}<\mu({}^{*}\boldsymbol{n})\right\}=\left\{{}^{*}\boldsymbol{n}\middle|\ \overline{|{}^{*}\boldsymbol{n}|}\cdot\mu^{1/s}\in{}^{\star}\mathbb{PR}_{\varepsilon}\right\} $$

forms a group: for ${}^{*}\boldsymbol{m},{}^{*}\boldsymbol{n}\in({}^{*}\mathbb{Z}^{s})^{\mu}$,

$$ \overline{|{}^{*}\boldsymbol{m}+{}^{*}\boldsymbol{n}|}\cdot\mu^{1/s}\leq\left(\overline{|{}^{*}\boldsymbol{m}|}+\overline{|{}^{*}\boldsymbol{n}|}\right)\cdot\mu^{1/s}\in{}^{\star}\mathbb{PR}_{\varepsilon}. $$

The set $({}^{*}\mathbb{Z}^{s})^{\mu[\iota]}$ of elements of $({}^{*}\mathbb{Z}^{s})^{\mu}$ which in addition satisfy $\overline{|{}^{*}\boldsymbol{n}|}\cdot\mu^{1/s}<\iota$ forms a subgroup.

For $\boldsymbol{\varepsilon}=(\varepsilon_1,\dots,\varepsilon_r)\in{}^{*}\mathbb{R}^{r}_{\varepsilon}$, define $\overline{|{}^{*}\boldsymbol{\varepsilon}|}$ as in (32). Given ${}^{*}\boldsymbol{n}\in{}^{*}\mathbb{Z}^{s}(\Theta)$ denote by $\nu({}^{*}\boldsymbol{n})=\langle\overline{|\varepsilon({}^{*}\boldsymbol{n})|}\rangle$: then

$$ ({}^{*}\mathbb{Z}^{s})_{\nu}(\Theta)=\{{}^{*}\boldsymbol{n}\in{}^{*}\mathbb{Z}^{s}(\Theta)|\ \nu({}^{*}\boldsymbol{n})\leq\nu^{1/r}\} $$

is also a group. We have thus defined the **matrix ideology**

$$ {}^{*}\mathbb{Z}^{s}(\Theta)=\{({}^{*}\mathbb{Z}^{s})^{\mu}_{\nu}(\Theta)\}=\{({}^{*}\mathbb{Z}^{s})^{\mu[\iota]}_{\nu}(\Theta)\}. $$

In general, the dual group $({}^{*}\mathbb{Z}^{r})(\Theta)^{\perp}$ admits a filtration by growth only, $\{({}^{*}\mathbb{Z}^{r})^{\mu}(\Theta)^{\perp}\}$, except when ${}^{*}\widetilde{\mathbb{Z}}^{s}(\Theta)=0$ e.g. when Θ is square invertible. Nevertheless, we have

Lemma 3. *Let* ${}^*\boldsymbol{n} \in ({}^*\mathbb{Z}^s)^{\mu[\lambda]}(\Theta)$*. Then*

$$\overline{|{}^*\boldsymbol{n}^{\perp}|} \cdot \mu^{1/s} < \lambda,$$

that is, ${}^*\boldsymbol{n}^{\perp} \in ({}^*\mathbb{Z}^r)^{\mu^{r/s}[\lambda]}$*. If* Θ *is invertible of dimension* $r{\times}r$ *then* ${}^*\boldsymbol{n} \in ({}^*\mathbb{Z}^r)^{\mu[\lambda]}(\Theta) \Leftrightarrow {}^*\boldsymbol{n}^{\perp} \in ({}^*\mathbb{Z}^r)^{\mu[\lambda]}(\Theta)^{\perp}$*.*

Proof. We must show that $\langle\overline{|{}^*\boldsymbol{n}^{\perp}|}\rangle \cdot (\mu^{r/s})^{1/r} = \langle\overline{|{}^*\boldsymbol{n}^{\perp}|}\rangle \cdot \mu^{1/s} \in {}^{\star}\mathbb{PR}_{\varepsilon}$. But this follows immediately from:

$$\overline{|{}^*\boldsymbol{n}^{\perp}|} = \overline{|\Theta{}^*\boldsymbol{n}|} = \max_{i=1,\dots,r} \Big|\sum_{j=1}^{s} \Theta_{ij}{}^*n_j\Big| \leq s\overline{|\Theta|} \cdot \overline{|{}^*\boldsymbol{n}|}$$

where $\overline{|\Theta|} = \max_{ij} |\Theta_{ij}|$. In particular, $\overline{|{}^*\boldsymbol{n}^{\perp}|} \cdot \mu < \lambda$. The last statement follows from symmetry of argument. □

The relevant arithmetic operations for matrix ideological arithmetic are given by the Kronecker product and sum [18]. Let Θ, Θ' be real matrices of dimensions $r \times s$, $r' \times s'$. Denote by $\Theta \otimes \Theta'$ the **Kronecker (or tensor) product** i.e. the $rr' \times ss'$ block matrix

$$\Theta \otimes \Theta' = \begin{pmatrix} \theta_{11}\Theta' & \cdots & \theta_{1s}\Theta' \\ \vdots & \ddots & \vdots \\ \theta_{r1}\Theta' & \cdots & \theta_{rs}\Theta' \end{pmatrix}$$

where $\Theta = (\theta_{ij})$. If $r = s$ and $r' = s'$ the **Kronecker sum** and **Kronecker difference** are the $rr' \times rr'$ matrices

$$\Theta \oplus \Theta' = \Theta \otimes I_{r'} + I_r \otimes \Theta', \quad \Theta \ominus \Theta' = \Theta \otimes I_{r'} - I_r \otimes \Theta'.$$

Neither the Kronecker product nor the Kronecker sum/difference are commutative, nor do they satisfy the distributive law. If we assume Θ, Θ' are square and denote by $\sigma(\Theta)$, $\sigma(\Theta')$ their spectra then $\sigma(\Theta{\otimes}\Theta') = \sigma(\Theta){\cdot}\sigma(\Theta')$, $\sigma(\Theta{\oplus}\Theta') = \sigma(\Theta){+}\sigma(\Theta')$ and $\sigma(\Theta \ominus \Theta') = \sigma(\Theta) - \sigma(\Theta')$.

Denote by $\widetilde{\mathcal{M}}(\mathbb{R})$ the monoid of all real matrices with respect to the Kronecker product, and by $\mathcal{M}(\mathbb{R})$ the submonoid of square matrices, equipped further with the Kronecker sum/difference. Note that there is a monomorphism $(\mathbb{R}, +, \times) \hookrightarrow (\mathcal{M}(\mathbb{R}), \oplus, \otimes)$. Observe that if $\boldsymbol{m}$, $\boldsymbol{n}$ are vectors of size s resp. s' then

$$(\Theta \otimes \Theta')(\boldsymbol{m} \otimes \boldsymbol{n}) = \Theta(\boldsymbol{m}) \otimes \Theta'(\boldsymbol{n}); \tag{33}$$

if $r = s$ and $r' = s'$ and $\boldsymbol{m}$, $\boldsymbol{n}$ are vectors of size r resp. r' then

$$(\Theta \oplus \Theta')(\boldsymbol{m} \otimes \boldsymbol{n}) = \Theta(\boldsymbol{m}) \otimes \boldsymbol{n} + \boldsymbol{m} \otimes \Theta'(\boldsymbol{n}), \tag{34}$$

with a similar formula for the Kronecker difference. If we think of $(\Theta\boldsymbol{m}, \boldsymbol{m})$ as a "vector numerator denominator pair", then (33) is the denominator addition formula and (34) is the numerator addition formula.

Theorem 30 (Matrix Ideological Arithmetic)**.** *Let* Θ, Θ' *be real matrices of dimensions* $r \times s$, $r' \times s'$*. Then there is a well-defined bilinear pairing:*

$$\otimes : ({}^*\mathbb{Z}^s)^{\mu^s[\iota]}_{\nu^r}(\Theta) \times ({}^*\mathbb{Z}^{s'})^{\nu^{s'}[\lambda]}_{\mu^{r'}}(\Theta') \longrightarrow ({}^*\mathbb{Z}^{ss'})^{\mu^{ss'}\cdot\nu^{ss'}[\iota\cdot\lambda]}_{\iota+\lambda}(\Theta \otimes \Theta') \tag{35}$$

defined by the Kronecker product of vectors. If $r = s$ *and* $r' = s'$ *then the Kronecker product also defines a pairing*

(36)

$$\otimes : ({}^*\mathbb{Z}^r)^{\mu^r[\iota]}_{\nu^r}(\Theta) \times ({}^*\mathbb{Z}^{r'})^{\nu^{r'}[\lambda]}_{\mu^{r'}}(\Theta') \longrightarrow ({}^*\mathbb{Z}^{rr'})^{\mu^{rr'}\cdot\nu^{rr'}[\iota\cdot\lambda]}_{\iota+\lambda}(\Theta \oplus \Theta') \cap ({}^*\mathbb{Z}^{rr'})^{\mu^{rr'}\cdot\nu^{rr'}[\iota\cdot\lambda]}_{\iota+\lambda}(\Theta \ominus \Theta').$$

In particular, when $r = r' = s = s' = 1$ *we recover Theorem 11.*

Proof. We will prove (35); the proof of (36) is similar and is left to the reader. The argument amounts to replacing scalar product by tensor product in the calculations found in Theorem 11, taking into account the dimensional normalizations used to define the filtrations. Let $^*\boldsymbol{m} \in ({}^*\mathbb{Z}^s)^{\mu^s[\iota]}_{\nu^r}(\Theta)$, $^*\boldsymbol{n} \in ({}^*\mathbb{Z}^{s'})^{\nu^{s'}[\lambda]}_{\mu^{r'}}(\Theta')$. First observe that

$$\mu({}^*\boldsymbol{m} \otimes {}^*\boldsymbol{n}) = \mu({}^*\boldsymbol{m}) \cdot \mu({}^*\boldsymbol{n}) > \mu \cdot \nu = (\mu^{ss'} \cdot \nu^{ss'})^{1/ss'}$$

which implies $^*\boldsymbol{m} \otimes {}^*\boldsymbol{n} \in ({}^*\mathbb{Z}^{ss'})^{\mu^{ss'}\cdot\nu^{ss'}[\iota\cdot\lambda]}$. Denote by Θ_i the ith row of Θ, Θ'_j the jth row of Θ' and by $\odot$ the dot product. The rr' vector $\Theta \otimes \Theta'({}^*\boldsymbol{m} \otimes {}^*\boldsymbol{n})$ has coordinates indexed by $(i, j) \in I \times J$ where the latter is given the linear dictionary order. The (i, j) coordinate is given by

$$\begin{aligned}(\Theta_i \odot {}^*\boldsymbol{m}) \cdot (\Theta_j \odot {}^*\boldsymbol{n}) &= ({}^*\boldsymbol{m}_i^\perp + \varepsilon({}^*\boldsymbol{m})_i) \cdot ({}^*\boldsymbol{n}_j^\perp + \varepsilon({}^*\boldsymbol{n})_j) \\ &= {}^*\boldsymbol{m}_i^\perp {}^*\boldsymbol{n}_j^\perp + \varepsilon({}^*\boldsymbol{m})_i {}^*\boldsymbol{n}_j^\perp + {}^*\boldsymbol{m}_i^\perp \varepsilon({}^*\boldsymbol{n})_j + \varepsilon({}^*\boldsymbol{m})_i \varepsilon({}^*\boldsymbol{n})_j\end{aligned}$$

i.e.

(37)

$$\Theta \otimes \Theta'({}^*\boldsymbol{m} \otimes {}^*\boldsymbol{n}) - {}^*\boldsymbol{m}^\perp \otimes {}^*\boldsymbol{n}^\perp = \varepsilon({}^*\boldsymbol{m}) \otimes {}^*\boldsymbol{n}^\perp + {}^*\boldsymbol{m}^\perp \otimes \varepsilon({}^*\boldsymbol{n}) + \varepsilon({}^*\boldsymbol{m}) \otimes \varepsilon({}^*\boldsymbol{n}).$$

The term $\varepsilon({}^*\boldsymbol{m}) \otimes \varepsilon({}^*\boldsymbol{n})$ can be disregarded because upon passage to ${}^\star\mathbb{PR}_\varepsilon$ it is strictly less than the absolute values of the images of the other two terms on the right hand side of (37). Now

$$\langle\overline{|\varepsilon({}^*\boldsymbol{m}) \otimes {}^*\boldsymbol{n}^\perp|}\rangle = \overline{|{}^*\boldsymbol{n}^\perp|} \cdot \nu({}^*\boldsymbol{m}) \leq \overline{|{}^*\boldsymbol{n}^\perp|} \cdot \nu < \lambda,$$

where we have used Lemma 3. Similarly one shows that $\langle\overline{|{}^*\boldsymbol{m}^\perp \otimes \varepsilon({}^*\boldsymbol{n})|}\rangle < \iota$ so that $^*\boldsymbol{m} \otimes {}^*\boldsymbol{n} \in ({}^*\mathbb{Z}^{ss'})_{\iota+\lambda}(\Theta \otimes \Theta')$ as claimed. □

The characterization of classes of real numbers in terms of the nonvanishing spectrum found in §3 can be generalized to real matrices using the nonvanishing spectrum

$$\mathrm{Spec}(\Theta) = \{(\mu, \nu) |\, ({}^*\mathbb{Z}^s)^\mu_\nu(\Theta) \neq 0\}.$$

By viewing Θ as the family of linear forms $\{\Theta_i\}_{i=1}^r$ in s variables, we will obtain the familiar correspondence between the geometry of $\mathrm{Spec}(\Theta)$ and approximation classes.

We say that Θ is **rational** if there exists a vector $\boldsymbol{m} \in \mathbb{Z}^s$ such that $\Theta(\boldsymbol{n}) \in \mathbb{Z}^r$; otherwise we say that Θ is **irrational**. Denote the set of rational real matrices by $\tilde{\mathcal{Q}}(\mathbb{R})$ and by $\mathcal{Q}(\mathbb{R})$ the subset of square rational real matrices. Then both $\tilde{\mathcal{Q}}(\mathbb{R})$ and by $\mathcal{Q}(\mathbb{R})$ are closed with respect to $\otimes$, and $\mathcal{Q}(\mathbb{R})$ is closed with respect to $\oplus, \ominus$ as well. Of course, the restriction of the monomorphism $\mathbb{R} \hookrightarrow \mathcal{M}(\mathbb{R})$ to $\mathbb{Q}$ lies in $\mathcal{Q}(\mathbb{R})$.

Proposition 29. *For all* $\Theta \in \tilde{\mathcal{M}}(\mathbb{R})$, $\mathrm{Spec}(\Theta) \supset \{(\mu, \nu)|\mu < \nu\}$. *If* $\Theta \in \tilde{\mathcal{Q}}(\mathbb{R})$ *then* $\mathrm{Spec}(\Theta) = {}^\star\mathbb{PR}^2_\varepsilon$.

Proof. For $\Theta \in \tilde{\mathcal{M}} - \tilde{\mathcal{Q}}(\mathbb{R})$, this is an straightforward adaptation of the proof of Theorem 4 using Schmidt's "Dirichlet Theorem" for families of linear forms (Theorem 3A on page 36 of [22]). If $\Theta \in \tilde{\mathcal{Q}}(\mathbb{R})$ then $({}^*\mathbb{Z}^s)_{-\infty}(\Theta) \neq 0$ from which we get $\mathrm{Spec}(\Theta) = {}^\star\mathbb{PR}^2_\varepsilon$. □

A matrix Θ is **badly approximable** if there exists a constant $C > 0$ such that for all ${}^{*}\boldsymbol{n} \in {}^{*}\mathbb{Z}^{s}(\Theta)$,

$$\overline{|{}^{*}\boldsymbol{n}|}^{s} \cdot \overline{|\varepsilon({}^{*}\boldsymbol{n})|}^{r} > C.$$

This definition is equivalent to the definition given in [22] in terms of the family of linear forms $\{\Theta_i\}_{i=1}^{r}$. In what follows, $\mathrm{Spec}_{\mathrm{flat}}(\Theta)$ = the intersection of $\mathrm{Spec}(\Theta)$ with the diagonal.

Theorem 31. *$\Theta \in \tilde{\mathcal{M}}(\mathbb{R}) - \tilde{\mathcal{Q}}(\mathbb{R})$ is badly approximable $\Leftrightarrow \mathrm{Spec}(\Theta) = \{(\mu, \nu) | \mu < \nu\}$ $\Leftrightarrow \mathrm{Spec}_{\mathrm{flat}}(\Theta) = \emptyset$.*

Proof. This follows directly from the definition of being badly approximable. $\square$

Denote by $\tilde{\mathcal{B}}(\mathbb{R}) \supset \mathfrak{B}$ the set of badly approximable real matrices, by $\tilde{\mathcal{W}}(\mathbb{R}) = \tilde{\mathcal{M}}(\mathbb{R}) - (\tilde{\mathcal{B}}(\mathbb{R}) \cup \tilde{\mathcal{Q}}(\mathbb{R})) \supset \mathfrak{W}$ the set of well approximable matrices. In addition, we have the notion of **very well approximable matrices** $\tilde{\mathcal{W}}_{\kappa}(\mathbb{R})_{r,s}$ of exponent $\kappa > s$: in which for any $\lambda \in (s, \kappa)$ the infinitesimal equality

$$\overline{|{}^{*}\boldsymbol{n}|}^{\kappa} \cdot \overline{|\varepsilon({}^{*}\boldsymbol{n})|}^{r} \simeq 0.$$

has a solution. Write $\tilde{\mathcal{W}}_{>s}(\mathbb{R})_{r,s}$ for the set of very well approximable matrices of size $r \times s$. When $\kappa = \infty$ we get the **Liouville matrices** $\tilde{\mathcal{W}}_{\infty}(\mathbb{R})_{r,s}$. Let $\tilde{\mathcal{B}}(\mathbb{R})_{r,s}$ denote the space of badly approximable matrices of size $r \times s$.

Theorem 32. *Let $\tilde{\mathcal{C}}(\mathbb{R})_{r,s}$ be any of the classes $\tilde{\mathcal{B}}(\mathbb{R})_{r,s}$, $\tilde{\mathcal{W}}(\mathbb{R})_{r,s}$, $\tilde{\mathcal{W}}_{>s}(\mathbb{R})_{r,s}$, $\tilde{\mathcal{W}}_{\infty}(\mathbb{R})_{r,s}$ described in the previous paragraph. Then*

$$\tilde{\mathcal{C}}(\mathbb{R})^{T}_{r,s} := \left\{\Theta^{T} \middle| \Theta \in \tilde{\mathcal{C}}(\mathbb{R})_{r,s}\right\} = \tilde{\mathcal{C}}(\mathbb{R})_{s,r}.$$

Proof. Khintchine transference. $\square$

We now introduce a generalized notion of projective linear equivalence appropriate to the matrix setting. Let $\mathrm{GL}_{r,s}(\mathbb{Z}) := \mathrm{GL}_{r+s}(\mathbb{Z})$ be the group of $(r+s) \times (r+s)$ integral invertible matrices, partitioned in the following way

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A_{r\times r} & B_{r\times s} \\ C_{s\times r} & D_{s\times s} \end{pmatrix}. \tag{38}$$

Note that the product of matrices $M, M' \in \mathrm{GL}_{r,s}(\mathbb{Z})$ viewed in block form follows the familiar formula for 2×2 matrix multiplication e.g.

$$M'M = \begin{pmatrix} A' & B' \\ C' & D' \end{pmatrix}\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A'A + B'C & A'B + B'D \\ C'A + D'C & C'B + D'D \end{pmatrix}. \tag{39}$$

If $M' = M^{-1}$ then $A'A+B'C = I_r$, $C'B+D'D = I_s$, $A'B+B'D = O_{r,s}$ and $C'A+D'C = O_{s,r}$, where I_r is the $r \times r$ identity matrix, $O_{r,s}$ is the $r \times s$ zero matrix, etc.

Let Θ, Θ' be $r \times s$ real matrices. We write

$$\Theta \bumpeq_{r,s} \Theta'$$

if there exists $M \in \mathrm{GL}_{r,s}(\mathbb{Z})$ with

$$(A\Theta + B) = \Theta'(C\Theta + D),$$

or equivalently, if

$$M \begin{pmatrix} \Theta \\ I_s \end{pmatrix} = \begin{pmatrix} \Theta' \\ I_s \end{pmatrix} \big(C\Theta + D\big). \tag{40}$$

Note that $\bumpeq_{1,1}$ is just $\bumpeq$, the usual relation of projective linear equivalence for scalars.

Theorem 33. $\Bumpeq_{r,s}$ *is an equivalence relation.*

Proof. We begin with transitivity. Suppose that $\Theta \Bumpeq_{r,s} \Theta'$ and $\Theta' \Bumpeq_{r,s} \Theta''$ by matrices M, M'. Then $\Theta'(C\Theta + D) = (A\Theta + B)$ and $\Theta''(C'\Theta' + D') = (A'\Theta' + B')$. We want to show that $\Theta \Bumpeq_{r,s} \Theta''$ via $M'M$. By the product formula (39), this amounts to showing

$$\Theta''\Big(\big(C'A + D'C\big)\Theta + \big(C'B + D'D\big)\Big) = \big(A'A + B'C\big)\Theta + \big(A'B + B'D\big).$$

However this follows from:

$$\begin{aligned}\Theta''\Big(\big(C'A + D'C\big)\Theta + \big(C'B + D'D\big)\Big) &= \Theta''\Big(C'\big(A\Theta + B\big) + D'\big(C\Theta + D\big)\Big)\\ &= \Theta''\big(C'\Theta' + D'\big)\big(C\Theta + D\big)\\ &= \big(A'\Theta' + B'\big)\big(C\Theta + D\big)\\ &= A'\big(A\Theta + B\big) + B'\big(C\Theta + D\big)\\ &= \big(A'A + B'C\big)\Theta + \big(A'B + B'D\big).\end{aligned}$$

As for symmetry: suppose that $\Theta \Bumpeq_{r,s} \Theta'$ by M, understood in the sense of (40). Let $M' = M^{-1}$; we want to show that

$$M'\begin{pmatrix}\Theta' \\ I_s\end{pmatrix} = \begin{pmatrix}\Theta \\ I_s\end{pmatrix}\big(C'\Theta' + D'\big).$$

Multiplying both sides of (40) by M' gives

$$\begin{pmatrix}\Theta \\ I_s\end{pmatrix} = M'\begin{pmatrix}\Theta' \\ I_s\end{pmatrix}\big(C\Theta + D\big).$$

Therefore we must show that $(C\Theta + D)(C'\Theta' + D') = I_s$: in fact, it suffices to show that $(C'\Theta' + D')(C\Theta + D) = I_s$. This follows from:

$$\begin{aligned}(C'\Theta' + D')(C\Theta + D) =& C'(A\Theta + B) + D'(C\Theta + D)\\ =& (C'A + D'C)\Theta + (C'B + D'D)\\ =& O_{s,r}\Theta + I_s = I_s.\end{aligned}$$

□

The matrix M implying $\Theta \Bumpeq_{r,s} \Theta'$ is clearly a projective invariant; by abuse of notation we will sometimes write $M(\Theta) = \Theta'$. We have therefore produced a partially defined action of $\mathrm{PGL}_{r,s}(\mathbb{Z})$ on $\widetilde{\mathcal{M}}(\mathbb{R})_{r,s}$ generalizing the projective linear action of $\mathrm{PGL}_2(\mathbb{Z})$ on $\mathbb{R}$: M acts on Θ provided there exists Θ' for which $\Theta \Bumpeq_{r,s} \Theta'$ via M.

Recall from [13] the group of numerator denominator pairs

$${}^*\mathbb{Z}^{r,s}(\Theta) = \{({}^*\boldsymbol{m}^{\perp}, {}^*\boldsymbol{m}) |\, {}^*\boldsymbol{m} \in {}^*\mathbb{Z}^s(\Theta)\}.$$

Theorem 34. *Suppose that* $M \in \mathrm{PGL}_{r,s}(\mathbb{Z})$ *acts on* $\Theta \in \widetilde{\mathcal{M}}(\mathbb{R})_{r,s}$*. Then the map*

$$({}^*\boldsymbol{m}^{\perp}, {}^*\boldsymbol{m}) \longmapsto ({}^*\boldsymbol{n}^{\perp}, {}^*\boldsymbol{n}) := (A{}^*\boldsymbol{m}^{\perp} + B{}^*\boldsymbol{m}, C{}^*\boldsymbol{m}^{\perp} + D{}^*\boldsymbol{m}) \tag{41}$$

defines an isomorphism of groups of numerator denominator pairs ${}^*\mathbb{Z}^{r,s}(\Theta) \cong {}^*\mathbb{Z}^{r,s}(M(\Theta))$*. In particular,*

$$({}^*\mathbb{Z}^s)^{\mu[\iota]}_{\nu}(\Theta) \cong ({}^*\mathbb{Z}^s)^{\mu[\iota]}_{\nu}(M(\Theta))$$

for all $\mu, \nu, \iota \in {}^*\mathbb{PR}_\varepsilon$.

Proof. Suppose that $\Theta({}^*\boldsymbol{m}) = {}^*\boldsymbol{m}^\perp + \varepsilon({}^*\boldsymbol{m})$. Then we may write

$$({}^*\boldsymbol{n}^\perp, {}^*\boldsymbol{n}) = (A(\Theta({}^*\boldsymbol{m}) - \varepsilon({}^*\boldsymbol{m})) + B^*\boldsymbol{m}, C(\Theta({}^*\boldsymbol{m}) - \varepsilon({}^*\boldsymbol{m})) + D^*\boldsymbol{m}).$$

Therefore,

$$\begin{aligned} \Theta'^*\boldsymbol{n} &\simeq \Theta'(C\Theta + D)^*\boldsymbol{m} \\ &= (A\Theta + B)^*\boldsymbol{m} \\ &\simeq {}^*\boldsymbol{n}^\perp. \end{aligned}$$

We leave the proof that (41) is invertible and respects ideological parameters to the reader. $\square$

We define the relation

$$\Theta_\mu \owedge_\nu \Theta'$$

when the groups occurring in (35) (with their dimensional normalizations of μ, ν) are nontrivial. The extent to which elements of $\tilde{\mathcal{M}}(\mathbb{R})$ are involved in ideological factorizations can be delineated according to class e.g. the elements of $\tilde{\mathcal{B}}(\mathbb{R})$ are anti prime, the elements of $\tilde{\mathcal{W}}_{>s}(\mathbb{R})_{r,s}$ are the omnidivisors etc. We leave it to the reader to formulate the matrix analogue of the divisibility discussion found at the end of §5.

Corollary 15. *Let* $\Theta, \Theta' \in \tilde{\mathcal{M}}(\mathbb{R}) - \tilde{\mathcal{Q}}(\mathbb{R})$ *be real matrices. If* $M \in \mathrm{PGL}_{r,s}(\mathbb{Z}), N \in \mathrm{PGL}_{r',s'}(\mathbb{Z})$ *act on* Θ, Θ' *then* $\Theta_\mu \owedge_\nu \Theta' \Leftrightarrow M(\Theta)_\mu \owedge_\nu N(\Theta')$.

The $\mathcal{O}$-approximation theory may be merged with the matrix theory in the more or less expected way. Given $K/\mathbb{Q}$ a finite extension of degree d, consider the sets $\tilde{\mathcal{M}}(\mathbb{K}) \supset \mathcal{M}(\mathbb{K})$ of matrices and square matrices with entries in $\mathbb{K}$, equipped with the Kronecker product as well as the Kronecker sum in the square case. For $\boldsymbol{\Theta} \in \tilde{\mathcal{M}}(\mathbb{K})$ of dimension $r \times s$, the diophantine approximation group is denoted ${}^*\mathcal{O}^s(\boldsymbol{\Theta})$ (see [13]).

If ${}^*\boldsymbol{\alpha} = ({}^*\alpha_1, \dots, {}^*\alpha_s) \in \mathcal{O}^s$, let ${}^*\alpha_i \in \mathbb{K}$ have place-indexed coordinates

$${}^*\alpha_i = ({}^*\alpha_{i,\tau_1}, \dots, {}^*\alpha_{i,\tau_d})$$

and define

$$\overline{|{}^*\boldsymbol{\alpha}|} = (\max_{i=1,\dots,s} |{}^*\alpha_{i,\tau_1}|, \dots, \max_{i=1,\dots,s} |{}^*\alpha_{i,\tau_d}|) \in {}^*\mathbb{R}^d_+.$$

Write $\boldsymbol{\mu}({}^*\boldsymbol{\alpha}) = \langle\overline{|{}^*\boldsymbol{\alpha}|}\rangle^{-1}$ and similarly $\boldsymbol{\nu}({}^*\boldsymbol{\alpha}) = \overline{|\varepsilon({}^*\boldsymbol{\alpha})|}$. With these definitions we obtain the ideology

$${}^*\mathcal{O}^s(\boldsymbol{\Theta}) = \{({}^*\mathcal{O}^s)^{\boldsymbol{\mu}[\boldsymbol{\iota}]}_{\boldsymbol{\nu}}(\boldsymbol{\Theta})\}.$$

Ideological arithmetic in this setting takes the anticipated form upon combining Theorems 27 and 30.

Restrict to the case of row vectors: let $\boldsymbol{\theta} = (\theta_1, \dots, \theta_s) \in \mathbb{K}^s$, so that

$${}^*\mathcal{O}^s(\boldsymbol{\theta}) = \{{}^*\boldsymbol{\alpha} \in \mathcal{O}^s |\, \exists {}^*\alpha^\perp \in {}^*\mathcal{O} \text{ s.t. } \boldsymbol{\theta} \odot {}^*\boldsymbol{\alpha} \simeq {}^*\alpha^\perp \in {}^*\mathcal{O}\}$$

where $\odot$ is dot product. The dual group

$${}^*\mathcal{O}(\boldsymbol{\theta})^\perp = \{{}^*\alpha^\perp |\, \exists {}^*\boldsymbol{\alpha} \in {}^*\mathcal{O}^s \text{ s.t. } \boldsymbol{\theta} \odot {}^*\boldsymbol{\alpha} \simeq {}^*\alpha^\perp\} \subset {}^*\mathcal{O}$$

is the **nonprincipal diophantine approximation group** generated by the coordinates of $\boldsymbol{\theta}$. Note that if $\boldsymbol{\gamma} = (\gamma_1, \gamma_2)$ for $\gamma_1, \gamma_2 \in \mathcal{O}$ then ${}^*\mathcal{O}(\boldsymbol{\gamma})^\perp$ is the ideal generated by γ_1, γ_2 in ${}^*\mathcal{O}$. If the Kronecker product ${}^*\boldsymbol{\alpha} \otimes {}^*\boldsymbol{\beta}$ for ${}^*\boldsymbol{\alpha} \in {}^*\mathcal{O}^s(\boldsymbol{\theta})$, ${}^*\boldsymbol{\beta} \in {}^*\mathcal{O}^{s'}(\boldsymbol{\eta})$ defines via ideological arithmetic an element of ${}^*\mathcal{O}^{ss'}(\boldsymbol{\theta} \otimes \boldsymbol{\eta})$ then the corresponding duals multiply as elements of ${}^*\mathcal{O}$:

$$({}^*\boldsymbol{\alpha} \otimes {}^*\boldsymbol{\beta})^{\perp_{\boldsymbol{\theta} \otimes \boldsymbol{\eta}}} = {}^*\alpha^{\perp_{\boldsymbol{\theta}}}\, {}^*\beta^{\perp_{\boldsymbol{\eta}}}.$$

In particular, if we restrict to vectors whose entries belong to K we recover the multiplicative arithmetic of ideals:

- If $\boldsymbol{\gamma} \in K^s$ then ${}^*\mathcal{O}(\boldsymbol{\gamma})^{\perp}$ is the ideal generated by the entries of $\boldsymbol{\gamma}$. Thus there exists an element $\boldsymbol{\gamma}_0 \in K^2$ such that ${}^*\mathcal{O}(\boldsymbol{\gamma})^{\perp} = {}^*\mathcal{O}(\boldsymbol{\gamma}_0)^{\perp}$.
- The ideological arithmetic of the Kronecker product, on the level of dual groups, is the pairing

  $$ {}^*\mathcal{O}(\boldsymbol{\gamma})^{\perp} \cdot {}^*\mathcal{O}(\boldsymbol{\gamma}')^{\perp} \longrightarrow {}^*\mathcal{O}((\boldsymbol{\gamma} \otimes \boldsymbol{\gamma}')_0)^{\perp}. $$

  defined by the ordinary product in ${}^*\mathcal{O}$.

We end this section with a few notes regarding the further development of the matrix theory along the lines of the scalar theory:

(1) A theory of flat matrix arithmetic would appear to be, for the moment, out of reach since we do not have yet at our disposable a matrix continued fraction algorithm yielding a good notion of best approximations (in the sense of satisfying the matrix analogue of Dirichlet's Theorem), see for example [23].

(2) One may define on ${}^*\mathbb{Z}^s(\Theta)$ the Minkowskian norm $[{}^*\boldsymbol{m}]_{\Theta} := \overline{|{}^*\boldsymbol{m}|}^{s} \cdot \overline{|\varepsilon({}^*\boldsymbol{m})|}^{r}$ as well as the set of symmetric diophantine approximations

$$({}^*\mathbb{Z}^s)^{\mathrm{sym}}(\Theta) = \{{}^*\boldsymbol{m} \in {}^*\mathbb{Z}^s(\Theta) |\, 0 < [{}^*\boldsymbol{m}]_{\Theta} < \infty\} \cup \{0\}.$$

For $\Theta \in \tilde{\mathcal{B}}(\mathbb{R})$, $({}^*\mathbb{Z}^s)^{\mathrm{sym}}(\Theta)$ is non trivial; again, due to the lack of a good notion of matrix best approximations, we can only conjecture that for general $\Theta \in \tilde{\mathcal{M}}(\mathbb{R})$, $({}^*\mathbb{Z}^s)^{\mathrm{sym}}(\Theta) \neq 0$.

10. POLYNOMIAL IDEOLOGY AND RATIONAL MAPS

In this section we begin our study of the ideological structure of polynomial diophantine approximation. In order to ease notation, we will write from now on

$${}^*\mathfrak{Z} := {}^*\mathbb{Z}[X] \subset {}^*\tilde{\mathfrak{Z}} := {}^*(\mathbb{Z}[X]).$$

Note that an element ${}^*f \in {}^*\mathfrak{Z}$ is a finite degree polynomial in the ring ${}^*\mathbb{Z}$ whereas an element ${}^*f = {}^*\{f_i\} \in {}^*\tilde{\mathfrak{Z}}$ has a well-defined possibly infinite degree $\deg({}^*f) = {}^*\{\deg(f_i)\} \in {}^*\mathbb{N}$. For each ${}^*d \in {}^*\mathbb{N}$ we define the additive subgroup

$${}^*\tilde{\mathfrak{Z}}_{{}^*d} := \{{}^*f \in {}^*\tilde{\mathfrak{Z}} |\ \deg({}^*f) \leq {}^*d\}.$$

This gives ${}^*\tilde{\mathfrak{Z}}$ the structure of a filtered ring:

$${}^*\tilde{\mathfrak{Z}}_{{}^*d} \cdot {}^*\tilde{\mathfrak{Z}}_{{}^*e} \subset {}^*\tilde{\mathfrak{Z}}_{{}^*d + {}^*e}.$$

We will also want to consider the following coarsening of the degree filtration. Let

$${}^*\check{d} = \{{}^*d' \in {}^*\mathbb{N} |\ \exists {}^*p \in ({}^*\mathbb{Q}^{\mathrm{fin}})^{\times}_{+} \text{ such that } {}^*d' = {}^*p\,{}^*d\}.$$

The set of such ${}^*\check{d}$ is ordered; note that

$${}^*\check{d} + {}^*\check{e} = ({}^*d + {}^*e)^{\vee}.$$

(Indeed, if say ${}^*\check{d} < {}^*\check{e}$ and ${}^*d' = {}^*p{}^*d$, ${}^*e' = {}^*q{}^*e$ where ${}^*p, {}^*q \in ({}^*\mathbb{Q}^{\text{fin}})^{\times}_{+}$ then ${}^*d' + {}^*e' = {}^*x({}^*d + {}^*e)$ where

$$ {}^*x = \frac{{}^*p{}^*d + {}^*q{}^*e}{{}^*d + {}^*e} \simeq {}^*q \in ({}^*\mathbb{Q}^{\text{fin}})^{\times}_{+}, $$

since ${}^*d/{}^*e$ is infinitesimal.) Then for each ${}^*\check{d}$,

$$ {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}} := \bigcup_{{}^*d \in {}^*\check{d}} {}^*\tilde{\mathfrak{Z}}_{{}^*d} $$

is a *ring* filtered by subgroups and

$$ {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}} \cdot {}^*\tilde{\mathfrak{Z}}_{{}^*\check{e}} \subset {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d} + {}^*\check{e}} $$

makes of ${}^*\tilde{\mathfrak{Z}}$ a ring filtered by rings. Note that ${}^*\tilde{\mathfrak{Z}}_{\check{1}} = {}^*\mathfrak{Z}$.

For each $\theta \in \mathbb{R}$, the ring of (infinite-degree) polynomial diophantine approximations is defined

$$ {}^*\tilde{\mathfrak{Z}}(\theta) = \{ {}^*f \in {}^*\tilde{\mathfrak{Z}} \mid {}^*f(\theta) \simeq 0 \}. $$

The subring ${}^*\mathfrak{Z}(\theta) = {}^*\tilde{\mathfrak{Z}}(\theta) \cap {}^*\mathfrak{Z}$ was studied in §6 of [13]. Defining ${}^*\tilde{\mathfrak{Z}}(\theta)_{{}^*\check{d}} = {}^*\tilde{\mathfrak{Z}}(\theta) \cap {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}}$ induces on ${}^*\tilde{\mathfrak{Z}}(\theta)$ a filtered ring structure as well.

We want to measure growth and decay of polynomials in ${}^*\tilde{\mathfrak{Z}}(\theta)$. Classically, there are a number of ways one measures the size of a polynomial (c.f. the Appendix to §3 of [26]). We will be interested in two: the height and the Mahler measure. Scaling by degree makes them tropical equivalent in a sense to made precise below.

For a standard polynomial $f \in \mathbb{Z}[X]$ denote by $\mathfrak{h}(f)$ the **height**: the maximum of the absolute values of the coefficients of f. Note that the height is subadditive. For ${}^*f = {}^*\{f_i\} \in \tilde{\mathfrak{Z}}$ define

$$ \mathfrak{h}({}^*f) = {}^*\{\mathfrak{h}(f_i)\} \in {}^*\mathbb{Z}_{+}. $$

On the other hand: recall that if $f(X) = a\prod(X - \alpha_i)$ the **Mahler measure** is

$$ \mathfrak{m}(f) = |a| \prod \max(1, |\alpha_i|). $$

By definition, the Mahler measure is multiplicative. The Mahler measure of ${}^*f$ is defined in the customary way: $\mathfrak{m}({}^*f) = {}^*\{\mathfrak{m}(f_i)\} \in {}^*\mathbb{R}_{+}$. The relationship between the height and the Mahler measure is given by the double inequality (see page 113 of [26]):

$$ (\deg({}^*f) + 1)^{-1/2} \cdot \mathfrak{m}({}^*f) \leq \mathfrak{h}({}^*f) \leq 2^{\deg({}^*f)} \cdot \mathfrak{m}({}^*f). \tag{42} $$

It will be to our advantage to scale growth according to degree. That is, for each ${}^*d \in {}^*\mathbb{N}$ we will define a ${}^*d$-scaled notion of size for elements of ${}^*\tilde{\mathfrak{Z}}_{{}^*d}$. In particular, the size of a fixed polynomial ${}^*f$ will be, as such, a relative measure, relative to the group ${}^*\tilde{\mathfrak{Z}}_{{}^*d}$ in which it is being considered. For ${}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*d}$ define

$$ \mathfrak{h}_{{}^*d}({}^*f) := \mathfrak{h}({}^*f)^{1/{}^*d}, \quad \mathfrak{m}_{{}^*d}({}^*f) := \mathfrak{m}({}^*f)^{1/{}^*d}. $$

Notice that by (42)

$$ \langle \mathfrak{h}_{{}^*d}({}^*f) \rangle = \langle \mathfrak{m}_{{}^*d}({}^*f) \rangle. $$

Thus, measuring degree-normalized growth in ${}^*\mathbb{PR}$ permits us to use either the height or the Mahler measure according to the exigencies of the situation we find ourselves in, as the paragraphs which follow illustrate.

In fact, for reasons which will soon become clear, it is natural to begin the parametrization of growth and decay in the polynomial setting using the Frobenius tropical semi ring ${}^{*}\overline{\mathbb{PR}}_{\varepsilon}$ (introduced at the end of §1) rather than ${}^{*}\mathbb{PR}_{\varepsilon}$. For any ${}^{*}d \geq \deg({}^{*}f)$ define

$$\bar{\mu}_{{}^*d}({}^*f) := [\mathfrak{h}_{{}^*d}({}^*f)^{-1}] = [\mathfrak{m}_{{}^*d}({}^*f)^{-1}] \in {}^{*}\overline{\mathbb{PR}}_{\varepsilon},$$

where we recall that $[{}^*x]$ is the image in ${}^*\overline{\mathbb{PR}}$ of ${}^*x \in {}^*\mathbb{R}$. Note that if ${}^*d' \in {}^*\check{d}$ then $\bar{\mu}_{{}^*d'}({}^*f) = \bar{\mu}_{{}^*d}({}^*f)$: for this reason the ${}^*\check{d}$-normalized expression

$$\bar{\mu}_{{}^*\check{d}}({}^*f) = \bar{\mu}_{{}^*d}({}^*f)$$

is well-defined. In particular, when ${}^*d = d \in \mathbb{N}$ we have $\bar{\mu}_{\check{d}}({}^*f) = \bar{\mu}_{\check{1}}({}^*f) = [\mathfrak{h}({}^*f)^{-1}]$. Thus we may define

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}} = \left\{{}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}} \middle|\, \bar{\mu}_{{}^*\check{d}}({}^*f) > \bar{\mu}\right\} = \left\{{}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}} \middle|\, \mathfrak{h}_{{}^*d}({}^*f)\cdot\bar{\mu} = \mathfrak{m}_{{}^*d}({}^*f)\cdot\bar{\mu} \in {}^*\overline{\mathbb{PR}}_{\varepsilon}\right\}$$

(where in the last expression above ${}^*d \in {}^*\check{d}$). Due to the subadditivity of the height, we have, assuming, say, $\mathfrak{h}_{{}^*d}({}^*f) \geq \mathfrak{h}_{{}^*d}({}^*g)$:

$$\begin{aligned}\mathfrak{h}_{{}^*d}({}^*f + {}^*g)\cdot\bar{\mu} &\leq \big(\mathfrak{h}({}^*f) + \mathfrak{h}({}^*g)\big)^{1/{}^*d}\cdot\bar{\mu}\\ &\leq 2^{1/{}^*d}\,\mathfrak{h}_{{}^*d}({}^*f)\cdot\bar{\mu} \in {}^*\overline{\mathbb{PR}}_{\varepsilon}\end{aligned}$$

so that ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}}$ is a group. Moreover, by the multiplicativity of the Mahler measure, we also have:

$$\begin{aligned}\bar{\mu}_{{}^*\check{d}+{}^*\check{e}}({}^*f\cdot{}^*g) &= [(\mathfrak{m}({}^*f)\cdot\mathfrak{m}({}^*g))^{1/({}^*d+{}^*e)}]^{-1}\\ &\geq \bar{\mu}_{{}^*\check{d}}({}^*f)\cdot\bar{\mu}_{{}^*\check{e}}({}^*g)\end{aligned}\tag{43}$$

for ${}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*\check{d}}$ and ${}^*g \in {}^*\tilde{\mathfrak{Z}}_{{}^*\check{e}}$. In particular, ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}}$ is a ring (take ${}^*\check{d} = {}^*\check{e}$ in (43) and note that $2{}^*\check{d} = {}^*\check{d}$, $\bar{\mu}^2 = \bar{\mu}$),

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}}\cdot{}^*\tilde{\mathfrak{Z}}^{\bar{\mu}'}_{{}^*\check{e}} \subset {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}\cdot\bar{\mu}'}_{{}^*\check{d}+{}^*\check{e}}$$

and therefore ${}^*\tilde{\mathfrak{Z}} = \{{}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}}\}$ has the structure of a ring doubly filtered by rings.

Now given $\theta \in \mathbb{R}$ and ${}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)_{{}^*\check{d}}$ we define the normalized decay via

$$\bar{\nu}_{{}^*\check{d}}({}^*f) := [|{}^*f(\theta)|^{1/{}^*d}], \quad {}^*d \in {}^*\check{d}$$

and the subring

$${}^*\tilde{\mathfrak{Z}}_{\bar{\nu}}(\theta)_{{}^*\check{d}} = \{{}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)_{{}^*\check{d}} |\, \bar{\nu}_{{}^*\check{d}}({}^*f) \leq \bar{\nu}\}.$$

This filtration by rings has the expected product law:

$${}^*\tilde{\mathfrak{Z}}_{\bar{\nu}}(\theta)_{{}^*\check{d}}\cdot{}^*\tilde{\mathfrak{Z}}_{\bar{\nu}'}(\theta)_{{}^*\check{e}} \subset {}^*\tilde{\mathfrak{Z}}_{\bar{\nu}\cdot\bar{\nu}'}(\theta)_{{}^*\check{d}+{}^*\check{e}}.$$

Write as before

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\check{d}} = {}^*\tilde{\mathfrak{Z}}_{\bar{\nu}}(\theta)_{{}^*\check{d}} \cap {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{{}^*\check{d}}.$$

Finally, note that for all ${}^*\check{d}$ and $n \in \mathbb{N}$, the power map ${}^*f \mapsto {}^*f^n$ induces a multiplicative inclusion

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\check{d}} \subset {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{\check{d}}.\tag{44}$$

Theorem 35. *The triply indexed filtration*

$${}^*\tilde{\mathfrak{Z}}(\theta) = \{{}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\check{d}}\}$$

by subrings makes of ${}^*\tilde{\mathfrak{Z}}(\theta)$ *a ring triply filtered by rings. Moreover,*

$${}^*\tilde{\mathfrak{Z}}^{\bar{\nu}}_{{}^*\check{d}}\cdot{}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\check{d}} \subset {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}\cdot\bar{\nu}}(\theta)_{{}^*\check{d}}.\tag{45}$$

Proof. The proof of the first statement is contained in the paragraphs preceding the Theorem. As for the second statement, note that for all ${}^*g \in {}^*\tilde{\mathfrak{Z}}_{{}^*\breve{d}}$ with $|{}^*g(\theta)| \geq 1$, ${}^*d \in \breve{d}$,

$$\begin{aligned}[|{}^*g(\theta)|^{1/{}^*d}] &\leq [\theta^{\deg({}^*g)/{}^*d}(\deg({}^*g)\cdot \mathfrak{h}({}^*g))^{1/{}^*d}] \\ &= [\mathfrak{h}_{{}^*d}({}^*g)].\end{aligned}$$

If $|{}^*g(\theta)| \leq 1$ then it is trivial that $[|{}^*g(\theta)|^{1/{}^*d}] \leq \bar{1} \leq [\mathfrak{h}_{{}^*d}({}^*g)]$. Then for any ${}^*f \in {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\breve{d}}$ and ${}^*g \in {}^*\tilde{\mathfrak{Z}}^{\bar{\nu}}_{{}^*\breve{d}}$

$$\begin{aligned}\bar{\nu}_{{}^*\breve{d}}({}^*g\cdot{}^*f) &= [|{}^*g(\theta)\cdot{}^*f(\theta)|^{1/{}^*\breve{d}}] \\ &\leq [\mathfrak{h}_{{}^*\breve{d}}({}^*g)]\cdot\bar{\nu} \in {}^*\overline{\mathbb{PR}}_{\varepsilon}.\end{aligned}$$

□

We may thus speak of ${}^*\tilde{\mathfrak{Z}}(\theta)$ as being a **polynomial ideology** (the latter concept defined by equation (45) of Theorem 35). Since it is indexed by the Frobenius tropical semi-ring, we refer to it as the **Frobenius polynomial ideology**. We may view ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\breve{d}}$ as an ideology within the ring ${}^*\tilde{\mathfrak{Z}}_{{}^*\breve{d}}$. In particular, the Frobenius ideology can be regarded as providing a filtration of ${}^*\tilde{\mathfrak{Z}}(\theta)$ by ideologies, indexed by the degree class ${}^*\breve{d}$.

In order to bring the type of a Mahler class into play, we introduce as well the **ordinary polynomial ideology**

$${}^*\tilde{\mathfrak{Z}}(\theta) = \{{}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*d}\}$$

which is defined exactly as the Frobenius ideology, using ${}^*\mathbb{PR}_{\varepsilon}$ in place of ${}^*\overline{\mathbb{PR}}_{\varepsilon}$ to index degree normalized growth and decay. In particular, the ordinary ideology makes of ${}^*\tilde{\mathfrak{Z}}(\theta)$ a ring tri-filtered by groups, which satisfies the following analogue of Theorem 35:

Theorem 36. *For all* ${}^*d \leq {}^*e \in {}^*\mathbb{N}$ *and* $\mu, \nu \in {}^*\mathbb{PR}_{\varepsilon}$,

$${}^*\tilde{\mathfrak{Z}}^{\nu}_{{}^*d}\cdot{}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*e} \subset {}^*\tilde{\mathfrak{Z}}^{\mu\cdot\nu}(\theta)_{{}^*d+{}^*e}.$$

Proof. Fix ${}^*g \in {}^*\tilde{\mathfrak{Z}}^{\nu}_{{}^*d}$ and ${}^*f \in {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*e}$. By the same argument employed in the proof of Theorem 35, $\langle|{}^*g(\theta)|^{1/{}^*d}\rangle \leq \langle\mathfrak{h}_{{}^*d}({}^*g)\rangle$. Moreover,

$$\begin{aligned}\nu_{{}^*d+{}^*e}({}^*g\cdot{}^*f) &= \langle|{}^*g(\theta)\cdot{}^*f(\theta)|^{1/{}^*d+{}^*e}\rangle \\ &\leq \mathfrak{h}_{{}^*d}({}^*g)^{{}^*d/({}^*d+{}^*e)}\cdot\nu^{{}^*e/({}^*d+{}^*e)} \\ &\leq (\mathfrak{h}_{{}^*d}({}^*g)\cdot\nu)^{{}^*e/({}^*d+{}^*e)} \in {}^*\mathbb{PR}_{\varepsilon}.\end{aligned}$$

□

The ordinary ideology gives a refinement of the Frobenius ideology:

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu},\mu}_{\bar{\nu},\nu}(\theta)_{{}^*\breve{d},{}^*d} := {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*d} \cap {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\breve{d}}$$

whenever $\mu \in \bar{\mu}$, $\nu \in \bar{\nu}$ and ${}^*d \in {}^*\breve{d}$. Since $\bar{\mu}^2 = \bar{\mu}$, $\bar{\nu}^2 = \bar{\nu}$ and $2{}^*\breve{d} = {}^*\breve{d}$,

$${}^*\tilde{\mathfrak{Z}}^{\bar{\mu},\mu}_{\bar{\nu},\nu}(\theta)_{{}^*\breve{d},{}^*d}\cdot{}^*\tilde{\mathfrak{Z}}^{\bar{\mu},\mu'}_{\bar{\nu},\nu'}(\theta)_{{}^*\breve{d},{}^*d'} \subset {}^*\tilde{\mathfrak{Z}}^{\bar{\mu},\mu\cdot\mu'}_{\bar{\nu},\nu\cdot\nu'}(\theta)_{{}^*\breve{d},{}^*d+{}^*d'}.$$

Therefore, the ordinary ideology endows each ring ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\breve{d}}$ with the structure of a tri-filtered ring.

A ring homomorphism

$$\Phi : {}^*\tilde{\mathfrak{Z}}(\theta) \longrightarrow {}^*\tilde{\mathfrak{Z}}(\eta)$$

preserving the Frobenius ideology (and the ordinary ideology) will be called a(n) **(ordinary) Frobenius ideological homomorphism**, and if Φ is invertible, a(n) **(ordinary) Frobenius ideological isomorphism**. This latter notion of equivalence is too strong for our purposes so we weaken it in two essential ways.

1. First, we will need to allow that Φ be merely multiplicative (not necessarily preserving the additive structure), that is, we will consider the ideologies ${}^*\tilde{\mathfrak{Z}}(\theta)$ as simply semigroups, referring to them as **semigroup ideologies**, and use the terminology of the previous paragraph to qualify semigroup homomorphisms. It will be clear in this section and those that follow that in the polynomial theory, the multiplicative structure plays the role of the passive operation ("sum"); the role of active operation ("multiplication") will be played by the resultant product, see §12.
2. A pair $\Phi = (\Phi_1, \Phi_2)$ of injective Frobenius semigroup homomorphisms

   $$\Phi_1 : {}^*\tilde{\mathfrak{Z}}(\theta) \longrightarrow {}^*\tilde{\mathfrak{Z}}(\eta), \quad \Phi_2 : {}^*\tilde{\mathfrak{Z}}(\eta) \longrightarrow {}^*\tilde{\mathfrak{Z}}(\theta)$$

   for which there exists $m, n \in \mathbb{N}$ such that

   $$\Phi_1({}^*\tilde{\mathfrak{Z}}^{\mu^m}_{\nu^m}(\theta)_{*d}) \subset {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\eta)_{*d\cdot m}, \quad \Phi_2({}^*\tilde{\mathfrak{Z}}^{\mu^n}_{\nu^n}(\eta)_{*d}) \subset {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\eta)_{*d\cdot n}$$

   will be called an **ideological consensus**, and we denote the equivalence relation defined by an ideological consensus by

   $${}^*\tilde{\mathfrak{Z}}(\theta) \rightleftharpoons {}^*\tilde{\mathfrak{Z}}(\eta).$$

We now turn our attention to the search for a notion of equivalence of real numbers which is adapted to the dictates of polynomial diophantine approximation. We will insist that such a notion of equivalence implies an ideological consensus between corresponding ideologies. Since the equivalence appropriate to linear diophantine approximation is projective linear, it stands to reason that higher degree rational maps will provide a relevant notion of equivalence in the polynomial milieu.

Thus: let $\mathsf{Rat}_{\mathbb{Z}}$ be the set of rational maps that may be written in the "relatively prime form"

$$R(X) = \frac{p(X)}{q(X)}$$

where $p(X), q(X) \in \mathbb{Z}[X]$ are relatively prime and $q(X)$ is not the zero polynomial. We equip $\mathsf{Rat}_{\mathbb{Z}}$ with the operation $\circ$ of composition; its field operations will play no role in what follows.

Let us describe the relatively prime form of the composition $R \circ S$ of $R, S \in \mathsf{Rat}_{\mathbb{Z}}$. First, recall that the degree of R is defined $\deg(R) = \max(\deg p, \deg q)$, and that [3]

$$\deg(R \circ S) = \deg(R)\deg(S).$$

If we write $p(X) = \sum_{i=0}^{\deg p} p_i X^i$, $q(X) = \sum_{j=0}^{\deg q} q_j X^j$ and $S(X) = t(X)/u(X)$ with $t(X), u(X)$ relatively prime, then clearing denominators gives

$$(R \circ S)(X) = u(X)^{\deg q - \deg p} \cdot \frac{\sum_{i=0}^{\deg p} p_i t(X)^i u(X)^{\deg p - i}}{\sum_{j=0}^{\deg q} q_j t(X)^j u(X)^{\deg q - j}}. \tag{46}$$

Suppose that $\deg p \geq \deg q$ and $\deg t \geq \deg u$. Then the degree of the numerator of (46) is $\deg p \deg t$ and the degree of the denominator is

$$\deg q \deg t + (\deg p - \deg q) \deg u \leq \deg p \deg t.$$

This implies that the numerator and denominator of (46) are relatively prime: for otherwise, we would obtain $\deg(R \circ S) < \deg(R)\deg(S)$. The other three cases ($\deg p \leq \deg q$ and $\deg t \geq \deg u$, etc.) produce the same result, namely, that (46) is the relatively prime form of $\deg(R \circ S)$.

Write

$$\theta \geqslant \eta$$

if there exists $R \in \mathsf{Rat}_{\mathbb{Z}}$ such that $R(\theta) = \eta$. If $\eta \geqslant \theta$ as well, we will write

$$\theta \doteq \eta$$

and say that $\theta, \eta \in \mathbb{R}$ are **$\mathsf{Rat}_{\mathbb{Z}}$-equivalent**. Note that $\theta \doteq \eta \Leftrightarrow \mathbb{Q}(\theta) = \mathbb{Q}(\eta)$, and that $\theta \doteq \eta \Rightarrow \theta$ and η are algebraically dependent. The converse is false e.g. if $\mathbb{Q}(\theta, \eta)$ is not purely transcendental.

Given $R(X) = p(X)/q(X) \in \mathsf{Rat}_{\mathbb{Z}}$ and $f(X) = \sum_{i=0}^{d} a_i X^i \in \mathbb{Z}[X]$, define

$$(f \square R)(X) := \sum_{i=0}^{d} a_i p(X)^i q(X)^{n-i} = q(X)^d \cdot (f \circ R)(X). \tag{47}$$

This definition extends in the usual way to any ${}^*f \in {}^*\tilde{\mathfrak{Z}}$. Note that

$$\deg({}^*f \square R) = \deg({}^*f) \cdot \deg(R).$$

In particular,

$${}^*\tilde{\mathfrak{Z}}_{{}^*d} \square R \subset {}^*\tilde{\mathfrak{Z}}_{{}^*d \cdot \deg(R)} \quad \text{and} \quad {}^*\tilde{\mathfrak{Z}}_{{}^*\breve{d}} \square R \subset {}^*\tilde{\mathfrak{Z}}_{{}^*\breve{d}}$$

Proposition 30. *The map*

$$({}^*\tilde{\mathfrak{Z}}, \cdot) \times \mathsf{Rat}_{\mathbb{Z}} \longrightarrow ({}^*\tilde{\mathfrak{Z}}, \cdot)$$

described by (47) defines an action of $\mathsf{Rat}_{\mathbb{Z}}$ *by multiplicative monomorphisms. That is, for all* ${}^*f, {}^*g \in {}^*\tilde{\mathfrak{Z}}$ *and* $R, S \in \mathsf{Rat}_{\mathbb{Z}}$*,*

1. $({}^*f \cdot {}^*g) \square R = ({}^*f \square R) \cdot ({}^*g \square R)$.
2. $({}^*f \square R) \square S = {}^*f \square (R \circ S)$.

Proof. Item (1) is clear. It is enough to prove (2) for a standard polynomial f of degree d. Suppose that $R = p(X)/q(X)$, $S(X) = t(X)/u(X)$ are parametrized as before, and that $\deg R = \deg p$. Then

$$\begin{aligned}
(f \square R) \square S &= \big(q(X)^d \cdot (f \circ R)(X)\big) \square S \\
&= q(t(X)/u(X))^d \cdot (f \circ (R \circ S))(X) \cdot u(X)^{d \deg p} \\
&= \Big(u(X)^{\deg p - \deg q} \cdot \sum q_j t(X)^j u(X)^{\deg q - j}\Big)^d \cdot (f \circ (R \circ S))(X) \\
&= f \square (R \circ S)
\end{aligned}$$

where the last equality follows from the "relatively prime form" of $R \circ S$ given in (46). The case where $\deg R = \deg q$ produces the same result. $\square$

Note 10. If we restrict to $\mathrm{PGL}_2(\mathbb{Z}) \subset \mathsf{Rat}_{\mathbb{Z}}$ acting on ${}^*\mathfrak{Z}_1$ we essentially recover the action of $\mathrm{PGL}_2(\mathbb{Z})$ (by inverses of matrices) on projective classes of vectors in ${}^*\mathbb{Z}^2$.

We now consider the effect of the action of $\mathsf{Rat}_{\mathbb{Z}}$ on both the ordinary and the Frobenius ideological structure of ${}^{*}\tilde{\mathfrak{Z}}$. Given $R \in \mathsf{Rat}_{\mathbb{Z}}$ define $\mathfrak{h}(R) = \max(\mathfrak{h}(p), \mathfrak{h}(q))$. Fix ${}^{*}f \in {}^{*}\tilde{\mathfrak{Z}}_{{}^{*}d}^{\mu^{\deg(R)}}$. Then

$$\langle \mathfrak{h}({}^{*}f \square R)\rangle \leq \langle \mathfrak{h}({}^{*}f)\mathfrak{h}(R)^{{}^{*}d}\rangle.$$

Since ${}^{*}f \square R \in {}^{*}\tilde{\mathfrak{Z}}_{{}^{*}d\cdot\deg(R)}$, we will calculate its growth index with respect to normalization by the degree ${}^{*}d \cdot \deg(R)$:

$$\begin{aligned}\mu({}^{*}f \square R) &= \langle \mathfrak{h}({}^{*}f \square R)^{-1/({}^{*}d\cdot\deg(R))}\rangle \\ &\geq \mathfrak{h}(R)^{-1/\deg(R)} \cdot \mu \\ &= \mu.\end{aligned}$$

Therefore,

$${}^{*}\tilde{\mathfrak{Z}}_{{}^{*}d}^{\mu^{\deg(R)}} \square R \subset {}^{*}\tilde{\mathfrak{Z}}^{\mu}_{{}^{*}d\cdot\deg(R)}.$$

Now suppose that ${}^{*}f \in {}^{*}\tilde{\mathcal{Z}}_{\nu^{\deg(R)}}(\theta)_{{}^{*}d}$ and $R(\eta) = \theta$. Then we claim that ${}^{*}f \square R \in {}^{*}\tilde{\mathcal{Z}}_{\nu}(\eta)_{{}^{*}d\cdot\deg(R)}$. Indeed, calculating the decay $\nu({}^{*}f \square R)$ (with respect to η) we have

$$\begin{aligned}\nu({}^{*}f \square R) &= \langle |({}^{*}f \square R)(\eta)|^{1/{}^{*}d\cdot\deg(R)}\rangle \\ &= \langle |{}^{*}f(\theta)\cdot q(\theta)|^{1/{}^{*}d\cdot\deg(R)}\rangle \\ &\leq \nu.\end{aligned}$$

Therefore,

$${}^{*}\tilde{\mathfrak{Z}}_{\nu^{\deg(R)}}(\theta)_{{}^{*}d} \square R \subset {}^{*}\tilde{\mathfrak{Z}}_{\nu}(\eta)_{{}^{*}d\cdot\deg(R)}.$$

We have shown:

Theorem 37. *Suppose that* $\theta \preccurlyeq \eta$. *Then*

$${}^{*}\tilde{\mathfrak{Z}}^{\mu^{\deg(R)}}_{\nu^{\deg(R)}}(\theta)_{{}^{*}d} \square R \subset {}^{*}\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\eta)_{{}^{*}d\cdot\deg(R)} \quad \textit{and} \quad {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}\breve{d}} \square R \subset {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\eta)_{{}^{*}\breve{d}}.$$

In particular, if $\theta \doteq \eta$ *then the associated pair of rational maps* (R, S) *defines an ideological consensus* ${}^{*}\tilde{\mathfrak{Z}}(\theta) \rightleftharpoons {}^{*}\tilde{\mathfrak{Z}}(\eta)$.

11. POLYNOMIAL NONVANISHING SPECTRA AND MAHLER'S CLASSIFICATION

For each degree class ${}^{*}\breve{d}$, define the ${}^{*}\breve{\boldsymbol{d}}$ **Frobenius nonvanishing spectrum** by

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^{*}\breve{d}} = \left\{(\bar{\mu}, \bar{\nu}) \middle| \; {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}\breve{d}} \neq 0\right\}.$$

The Frobenius spectrum refines as follows: for each ${}^{*}d \in {}^{*}\breve{d}$, define

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^{*}d} = \left\{(\bar{\mu}, \bar{\nu}) \middle| \; {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^{*}d} \neq 0\right\},$$

then for ${}^{*}d' \in {}^{*}\breve{d}$, ${}^{*}d' > {}^{*}d$, we have an inclusion $\overline{\mathrm{Spec}}[X](\theta)_{{}^{*}d} \subset \overline{\mathrm{Spec}}[X](\theta)_{{}^{*}d'}$ and thus we obtain the filtration

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^{*}\breve{d}} = \bigcup_{{}^{*}d\in{}^{*}\breve{d}} \overline{\mathrm{Spec}}[X](\theta)_{{}^{*}d}.$$

Furthermore, if we formally add the "northeast perimeter"

$$(48) \qquad {}^{\star}\mathbb{PR}^{2}_{\leq 1} - {}^{\star}\mathbb{PR}^{2}_{\varepsilon} = (\{\bar{1}\} \times {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}) \cup ({}^{\star}\overline{\mathbb{PR}}_{\varepsilon} \times \{\bar{1}\}) \cup (\bar{1}, \bar{1})$$

to $\overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{d}}$, then for all ${}^*\breve{d} < {}^*\breve{e}$ there is a well-defined filtration preserving inclusion

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{d}} \hookrightarrow \overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{e}}, \quad (\bar{\mu},\bar{\nu}) \mapsto (\bar{\mu}^{{}^*\breve{d}/{}^*\breve{e}}, \bar{\nu}^{{}^*\breve{d}/{}^*\breve{e}}). \tag{49}$$

The ${}^*\breve{d}/{}^*\breve{e}$-powers are well-defined since ${}^*\breve{d}/{}^*\breve{e}$ is the class of an infinitesimal *c.f. Note* 1 of §1; if either of the coordinates of $(\bar{\mu},\bar{\nu})$ is a ${}^*\breve{d}/{}^*\breve{e}$-root of unity, the image of the pair will belong to the perimeter (48). The **Frobenius nonvanishing spectrum** is the collection

$$\overline{\mathrm{Spec}}[X](\theta) = \{\overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{d}}\}$$

viewed as a directed system of filtered sets, where the inclusions are the power laws (49).

Likewise we may define the **ordinary nonvanishing spectrum**

$$\mathrm{Spec}[X](\theta) = \{\mathrm{Spec}[X](\theta)_{{}^*d}\}, \quad \mathrm{Spec}[X](\theta)_{{}^*d} = \left\{(\mu,\nu) \middle|\, {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*d} \neq 0\right\}.$$

Note 11. Observe that $(\bar{\mu},\bar{\nu}) \in \overline{\mathrm{Spec}}[X](\theta)_{{}^*d}$ implies that for all $\mu \in \bar{\mu}, \nu \in \bar{\nu}$, $(\mu,\nu) \in \mathrm{Spec}[X](\theta)_{{}^*d}$. The converse may not be true however: if $0 \neq {}^*f \in {}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu}(\theta)_{{}^*d}$ with $\bar{\mu}_{{}^*d}({}^*f) = \bar{\mu}$ then ${}^*f \notin {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*d}$. So the ordinary nonvanishing spectrum carries finer information not contained in the Frobenius nonvanishing spectrum.

We say that $\overline{\mathrm{Spec}}[X](\theta)$ and $\overline{\mathrm{Spec}}[X](\eta)$ are **finite power law equivalent** and write

$$\overline{\mathrm{Spec}}[X](\theta) \overset{\circ}{=} \overline{\mathrm{Spec}}[X](\eta)$$

if there exist $m, n \in \mathbb{N}$ such that for all ${}^*d$,

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^*d} \subset \overline{\mathrm{Spec}}[X](\eta)_{{}^*d\cdot m} \quad \text{and} \quad \overline{\mathrm{Spec}}[X](\eta)_{{}^*d} \subset \overline{\mathrm{Spec}}[X](\theta)_{{}^*d\cdot n}.$$

Clearly $\overline{\mathrm{Spec}}[X](\theta) \overset{\circ}{=} \overline{\mathrm{Spec}}[X](\eta)$ implies that $\overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{d}} = \overline{\mathrm{Spec}}[X](\eta)_{{}^*\breve{d}}$ for every class ${}^*\breve{d}$ but the converse need not be true.

We have the following Corollary to Theorem 37:

Corollary 16. *Suppose that* $\theta \preccurlyeq \eta$. *Then*

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^*d} \subset \overline{\mathrm{Spec}}[X](\eta)_{{}^*d\cdot \deg(R)} \ \textit{and}\ \overline{\mathrm{Spec}}[X](\theta)_{{}^*\breve{d}} \subset \overline{\mathrm{Spec}}[X](\eta)_{{}^*\breve{d}}.$$

If $\theta \doteq \eta$ *then*

$$\overline{\mathrm{Spec}}[X](\theta) \overset{\circ}{=} \overline{\mathrm{Spec}}[X](\eta).$$

Since finite degrees $d < e$ define the same Frobenius class, we have:

$$\overline{\mathrm{Spec}}[X](\theta)_{d} \subset \overline{\mathrm{Spec}}[X](\theta)_{e}. \tag{50}$$

From this we may derive the following nonvanishing result:

Proposition 31. *For all* ${}^*d \in {}^*\mathbb{N}$, $\{(\bar{\mu},\bar{\nu}) |\, \bar{\mu} < \bar{\nu}\} \subset \overline{\mathrm{Spec}}[X](\theta)_{{}^*d}$.

Proof. We first show that $\overline{\mathrm{Spec}}[X](\theta)_1$ contains the image of $\mathrm{Spec}(\theta)$ in ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$. Indeed, given $\bar{\mu} < \bar{\nu}$, since ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ is a dense linear order (Proposition 5), there exists $\bar{\mu}'$ with $\bar{\mu} < \bar{\mu}' < \bar{\nu}$. Choose $\mu' \in \bar{\mu}', \nu \in \bar{\nu}$ and $0 \neq {}^*n \in {}^*\mathbb{Z}^{\mu'}_{\nu}(\theta)$. Then ${}^*f_{{}^*n}(X) = {}^*nX - {}^*n^{\perp}$ satisfies $\bar{\mu}({}^*f_{{}^*n}) = [\mu({}^*n)] \geq \bar{\mu}' > \bar{\mu}$ and $\bar{\nu}({}^*f_{{}^*n}) = [\nu({}^*n)] \leq \bar{\nu}$. By (50), this implies that $(\bar{\mu},\bar{\nu}) \in \overline{\mathrm{Spec}}[X](\theta)_d$ for $d \in \mathbb{N}$. For infinite degree ${}^*d$, the power law inclusion (49) maps the set of pairs $\bar{\mu} < \bar{\nu}$ in $\overline{\mathrm{Spec}}[X](\theta)_1$ of elements which are not $({}^*d)^{-1}$ roots of unity onto $\{(\bar{\mu},\bar{\nu}) |\, \bar{\mu} < \bar{\nu}\}$. $\square$

We examine the nonvanishing spectra in terms of Mahler's classification, whose definition we now recall. For each pair of positive integers H, d, consider the polynomial $f_{H,d}(X)$ which minimizes $|f(\theta)|$ amongst those $f(X)$ of degree at most d with $f(\theta)\neq 0$ and satisfying $\mathfrak{h}(f)\leq H$. Define

$$\mathfrak{e}(d,H)=\frac{\log|f_{H,d}(\theta)|}{\log H^{-d}},$$

so that

$$|f_{H,d}(\theta)|^{1/d}=H^{-\mathfrak{e}(d,H)}.$$

Consider the limits

$$\mathfrak{e}(d)=\lim_{H\to\infty}\sup\mathfrak{e}(d,H),\qquad \mathfrak{e}=\lim_{d\to\infty}\sup\mathfrak{e}(d).$$

Note that if $\theta\in\mathfrak{W}_{>1}$ then $\mathfrak{e}(1)=\kappa=$ exponent of θ (the supremum of $\lambda>1$ for which $\theta\in\mathfrak{W}_{\lambda}$).

For d fixed, let $\{g_i=f_{H_i,d}\}$ be a subsequence of $\{f_{H,d}\}_{H\geq 1}$ for which $\mathfrak{h}(f_{H_i,d})=H_i$ and $\mathfrak{e}_i(d):=\mathfrak{e}(d,H_i)$ converges to $\mathfrak{e}(d)$. The class

$$^*\widehat{g}_d\in{}^*\mathfrak{Z}(\theta)_d$$

corresponding to $\{g_i\}$ is called a **degree d best polynomial class**. When $d=1$, then $^*\widehat{g}_1$ is identified with a best denominator class $^*\widehat{q}$ via the isomorphism $^*\mathbb{Z}(\theta)\cong{}^*\mathfrak{Z}(\theta)_1$. Denote by $\widehat{\bar{\mu}}_d=\bar{\mu}(^*\widehat{g}_d)$ and $\widehat{\bar{\nu}}_d=\bar{\nu}(^*\widehat{g}_d)$: the associated **degree d best polynomial growth** and **decay classes**.

Similarly, let $\{g_i=f_{H_i,d_i}\}$ be a subsequence of $\{f_{H,d}\}_{H,d\geq 1}$ with $H_i,d_i\to\infty$ for which $\mathfrak{h}(f_{H_i,d_i})=H_i$, $\deg(f_{H_i,d_i})=d_i$ and $\mathfrak{e}_i:=\mathfrak{e}(d_i,H_i)$ converges to $\mathfrak{e}$. Then the class

$$^*\widehat{g}_{^*d}\in{}^*\tilde{\mathfrak{Z}}(\theta)_{^*d}$$

corresponding to $\{g_i\}$ is called a **degree $^*d$ best polynomial class**, with associated **degree $^*d$ best polynomial growth** $\widehat{\bar{\mu}}_{^*d}=\bar{\mu}_{^*d}(^*g)$ and **decay** $\widehat{\bar{\nu}}_{^*d}=\bar{\nu}_{^*d}(^*g)$. The infinite degree $^*d={}^*\{d_i\}$ is called the corresponding **best degree**.

Note 12. When θ is algebraic, a best polynomial class is only best amongst those polynomials which do not have θ as a root. This contrasts with the process of producing linear best approximations for $\theta\in\mathbb{Q}$, which terminates in the root of the defining linear polynomial.

Proposition 32. *Suppose that* θ *is not algebraic of degree* $\leq d$ *(that* θ *is transcendental). Let* $(\widehat{\bar{\mu}}_d,\widehat{\bar{\nu}}_d)$ *be a finite degree d best growth-decay pair (let* $(\widehat{\bar{\mu}}_{^*d},\widehat{\bar{\nu}}_{^*d})$ *be an infinite degree best growth-decay pair). For all* $\bar{\mu}\geq\widehat{\bar{\mu}}_d$ *and* $\bar{\nu}<\widehat{\bar{\nu}}_d$ *(for all* $\bar{\mu}\geq\widehat{\bar{\mu}}_{^*d}$ *and* $\bar{\nu}<\widehat{\bar{\nu}}_{^*d}$*),*

$$^*\mathfrak{Z}^{\bar{\mu}}_{\bar{\nu}}(\theta)_d=0\qquad\left(^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{^*d}=0\right).$$

In particular,

$$\overline{\mathrm{Spec}}[X](\theta)_d\subsetneq{}^*\mathbb{PR}^2_{\varepsilon}\quad\left(\overline{\mathrm{Spec}}[X](\theta)_{^*d}\subsetneq{}^*\mathbb{PR}^2_{\varepsilon}\right).$$

Proof. We begin with the finite degree d statements. Suppose otherwise: let $\bar{\mu}\geq\widehat{\bar{\mu}}_d$ and $\bar{\nu}<\widehat{\bar{\nu}}_d$ yet $0\neq{}^*f\in{}^*\mathfrak{Z}^{\bar{\mu}}_{\bar{\nu}}(\theta)_d$. Then (since the height function is positive), $\mathfrak{h}(^*f)\cdot\widehat{\bar{\mu}}_d\leq\mathfrak{h}(^*f)\cdot\bar{\mu}\in{}^*\mathbb{PR}_{\varepsilon}$ implies that $\mathfrak{h}(^*f)<\mathfrak{h}(^*\widehat{g}_d)$. However, by definition of $^*\widehat{g}_d$, we must have $\bar{\nu}(^*f)\geq\widehat{\bar{\nu}}_d>\bar{\nu}$ (since by hypothesis $^*f(\theta)\neq 0$), contradiction. This argument applies *mutadis mutandis* to the infinite degree case. □

Thus, we may attach to any degree ${}^*d$ best polynomial class ${}^*\widehat{g}_{{}^*d}$ a horizontal strip in the $(\bar{\mu},\bar{\nu})$ plane:

$$V({}^*\widehat{g}_{{}^*d}) = \left\{(\bar{\mu},\bar{\nu})\middle|\ \bar{\mu}\geq\widehat{\bar{\mu}}_d \text{ and } \bar{\nu}<\widehat{\bar{\nu}}_d\right\}$$

parametrizing a region where ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*d}$ vanishes. The extent to which $V({}^*\widehat{g}_{{}^*d})$ persists in ${}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\theta)_{{}^*\breve{d}}$ is a function of the Mahler type of θ, a point which will be discussed in detail at the end of this section.

The Frobenius polynomial spectrum $\overline{\mathrm{Spec}}[X](\theta)$ does for the real algebraic numbers what the ordinary linear spectrum $\mathrm{Spec}(\theta)$ does for the rationals:

Corollary 17. *The following statements are equivalent.*

i. $\theta\in\mathbb{R}$ *is algebraic.*
ii. $\overline{\mathrm{Spec}}[X](\theta)_d = {}^{\star}\mathbb{PR}^2_{\varepsilon}$ *for some* $d\in\mathbb{N}$.
iii. $\overline{\mathrm{Spec}}[X](\theta)_{{}^*d} = {}^{\star}\mathbb{PR}^2_{\varepsilon}$ *for every* ${}^*d\in{}^*\mathbb{N}-\mathbb{N}$.

Proof. i. ⇒ ii., iii. If there exists $f(X)\in\mathbb{Z}[X]$ (a standard polynomial) with $f(\theta) = 0$ then for all ${}^*d\geq\deg(f)$, $\mathfrak{h}_{{}^*d}(f)\cdot\mu\in{}^{\star}\mathbb{PR}_{\varepsilon}$ for all $\mu\in{}^{\star}\mathbb{PR}_{\varepsilon}$, and $\nu_{{}^*d}(f) = -\infty<\nu$ for all $\nu\in{}^{\star}\mathbb{PR}_{\varepsilon}$. Then for such ${}^*d$, $\overline{\mathrm{Spec}}[X](\theta)_{{}^*d} = {}^{\star}\mathbb{PR}^2_{\varepsilon}$. ii., iii. ⇒ i. Follows from Proposition 32. □

Denote by $\mathfrak{d}$ the smallest degree d for which $\mathfrak{e}(d)$ is infinite; if $\mathfrak{e}(d)$ is finite for all d write $\mathfrak{d}=\infty$. Recall that Mahler's classification [2] consists of the identification of the following non-empty, mutually exclusive classes of numbers whose union is $\mathbb{R}$: θ is an

- ***A*-number** if $\mathfrak{e}=0$ and $\mathfrak{d}=\infty$. The A-numbers are exactly the real algebraic numbers.
- ***S*-number** if $0<\mathfrak{e}<\infty$ and $\mathfrak{d}=\infty$. Lebesgue-almost all real numbers are S-numbers.
- ***T*-number** if $\mathfrak{e}=\infty$ and $\mathfrak{d}=\infty$.
- ***U*-number** if $\mathfrak{e}=\infty$ and $\mathfrak{d}<\infty$.

Note that if θ and η are algebraically dependent, they are of the same Mahler class [5]: in particular, the relation $\theta\doteq\eta$ preserves Mahler class.

We now interpret Mahler's classification in terms of the images of pairs of best polynomial growth and best polynomial decay in the Frobenius growth-decay semiring ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$.

Theorem 38. θ *is an A-number ⇔ for every infinite degree polynomial best growth decay pair* $(\widehat{\bar{\mu}}_{{}^*d},\widehat{\bar{\nu}}_{{}^*d})$,

$$\widehat{\bar{\mu}}_{{}^*d}<(\widehat{\bar{\nu}}_{{}^*d})^{1/{}^*d}.$$

Proof. θ is an A-number ⇔ for every infinite degree best polynomial class ${}^*\widehat{g}_{{}^*d}$,

$$\left|{}^*\widehat{g}(\theta)\right|^{1/{}^*d} = \left(\mathfrak{h}({}^*\widehat{g})^{-1}\right)^{{}^*\mathfrak{e}}$$

where ${}^*\mathfrak{e} = {}^*\{\mathfrak{e}_i\}$ is infinitesimal. Then $\left(\mathfrak{h}({}^*\widehat{g})^{-1}\right)^{{}^*\mathfrak{e}} > \left(\mathfrak{h}({}^*\widehat{g})^{-1}\right)^{e}$ for any $e\in\mathbb{R}_+$, and the result follows on passing to ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ classes of degree ${}^*d$ normalized growth and decay. □

Theorem 39. θ *is an S-number ⇔ for every infinite degree polynomial best growth decay pair* $(\widehat{\bar{\mu}}_{{}^*d},\widehat{\bar{\nu}}_{{}^*d})$,

$$\widehat{\bar{\mu}}_{{}^*d}=(\widehat{\bar{\nu}}_{{}^*d})^{1/{}^*d}.$$

In this case, the associated vanishing strip $V({}^*\widehat{g}_{{}^*d})$ *persists in* $\overline{\mathrm{Spec}}[X](\theta)_{{}^*\check{d}}$*:*

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^*\check{d}} \cap V({}^*\widehat{g}_{{}^*d}) = \varnothing.$$

In particular,

$$\overline{\mathrm{Spec}}[X](\theta)_{{}^*\check{d}} \neq {}^{\star}\overline{\mathbb{PR}}^2_{\varepsilon}.$$

Proof. θ is an S-number $\Leftrightarrow$ for every infinite degree best polynomial class ${}^*\widehat{g}$ of degree ${}^*d$

$$|{}^*\widehat{g}(\theta)|^{1/{}^*d} = (\mathfrak{h}({}^*\widehat{g})^{-1})^{{}^*\mathfrak{e}}$$

where ${}^*\mathfrak{e} \simeq \mathfrak{e} \in \mathbb{R}_+$. Then the first statement follows upon passage to ${}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ classes of degree ${}^*d$ normalized growth and decay. Now suppose that ${}^*d < {}^*e$ both belong to the class ${}^*\check{d}$ i.e. ${}^*e/{}^*d$ is bounded. Suppose further that there exists a best growth decay pair $(\hat{\hat{\mu}}_{{}^*e}, \hat{\hat{\nu}}_{{}^*e}) \in V({}^*\widehat{g}_{{}^*d})$. Then $\hat{\hat{\mu}}_{{}^*e} \geq \hat{\hat{\mu}}_{{}^*d}$ and $\hat{\hat{\nu}}_{{}^*e} < \hat{\hat{\nu}}_{{}^*d}$, however:

$$(\hat{\hat{\nu}}_{{}^*d})^{1/{}^*d} = \hat{\hat{\mu}}_{{}^*d} \leq \hat{\hat{\mu}}_{{}^*e} = (\hat{\hat{\nu}}_{{}^*e})^{1/{}^*e}$$

that is

$$(\hat{\hat{\nu}}_{{}^*d})^{{}^*e/{}^*d} = \hat{\hat{\nu}}_{{}^*d} \leq \hat{\hat{\nu}}_{{}^*e},$$

contradiction. □

Theorem 40. θ *is a* T*-number* $\Leftrightarrow$ *for every*

- *infinite degree best growth decay pair* $(\hat{\hat{\mu}}_{{}^*d}, \hat{\hat{\nu}}_{{}^*d})$,
$$\hat{\hat{\mu}}_{{}^*d} > (\hat{\hat{\nu}}_{{}^*d})^{1/{}^*d}.$$
- *finite degree best growth decay pair* $(\hat{\hat{\mu}}_d, \hat{\hat{\nu}}_d)$,
$$\hat{\hat{\mu}}_d = \hat{\hat{\nu}}_d.$$

In the latter case, the associated vanishing strip $V({}^*\widehat{g}_d)$ *persists in* $\overline{\mathrm{Spec}}[X](\theta)_{\check{1}} = \overline{\mathrm{Spec}}[X](\theta)_{\check{d}}$*:*

$$\overline{\mathrm{Spec}}[X](\theta)_{\check{1}} \cap V({}^*\widehat{g}_d) = \varnothing.$$

In particular,

$$\overline{\mathrm{Spec}}[X](\theta)_{\check{1}} \neq {}^{\star}\overline{\mathbb{PR}}^2_{\varepsilon}.$$

Proof. θ is a T-number $\Leftrightarrow$ $\mathfrak{e}(d) < \infty$ for all d, the set of which is unbounded. Let ${}^*\widehat{g}_{{}^*d}$ be an infinite degree ${}^*d$ polynomial best class defined by the sequence $\{g_i = f_{H_i,d_i}\}$. Let ${}^*d = {}^*\{d_i\}$ and ${}^*\mathfrak{e} = {}^*\{\mathfrak{e}(H_i, d_i)\}$, the latter infinite by hypothesis. Then we have $|{}^*\widehat{g}_{{}^*d}(\theta)|^{1/{}^*d} = (\mathfrak{h}({}^*\widehat{g}_{{}^*d})^{-1})^{{}^*\mathfrak{e}}$ which implies after passing to Frobenius classes that

$$\hat{\hat{\mu}}_{{}^*d} = (\hat{\hat{\nu}}_{{}^*d})^{1/({}^*d{}^*\mathfrak{e})} > (\hat{\hat{\nu}}_{{}^*d})^{1/{}^*d}.$$

If ${}^*\widehat{g}_d$ is a finite degree d polynomial best class, then since $0 \leq \mathfrak{e}(d) < \infty$, we have

$$\hat{\hat{\mu}}_d \leq \hat{\hat{\nu}}_d.$$

But if the inequality were strict, it would contradict Proposition 31. The statement concerning vanishing strips, is argued exactly as in Theorem 39. □

We leave the proof of the last Theorem in this series to the reader.

Theorem 41. θ *is a* U*-number* $\Leftrightarrow$ *for every*

- *infinite degree best growth decay pair* $(\hat{\hat{\mu}}_{{}^*d}, \hat{\hat{\nu}}_{{}^*d})$,
$$\hat{\hat{\mu}}_{{}^*d} > (\hat{\hat{\nu}}_{{}^*d})^{1/{}^*d}.$$

- *There exists* $\mathfrak{d} \in \mathbb{N}$ *such that for every finite degree* $d \geq \mathfrak{d}$ *polynomial best growth decay pair* $(\widehat{\widehat{\mu}}_d, \widehat{\widehat{\nu}}_d)$*,*

$$\widehat{\widehat{\mu}}_d > \widehat{\widehat{\nu}}_d.$$

In Figure 3 we have displayed the sheets of the filtered Frobenius nonvanishing spectra for each of the four Mahler classes, (in the A-number portrait we have removed spectral elements corresponding to groups all of whose elements are multiples of the minimal polynomial).

Recall [5] that each of the transcendental Mahler classes can be further partitioned according to the **Mahler type**. To observe the type, we must work with the ordinary nonvanishing spectra.

For S-numbers, the type $\mathfrak{t}$ is equal to the exponent $\mathfrak{e} \in [1, \infty)$.

Theorem 42. *Let* θ *be an* S*-number of type* $\mathfrak{t} \in [1, \infty)$*. Then for every infinite degree* ${}^*d$ *best polynomial class* ${}^*\widehat{g}_{{}^*d}$*, the associated ordinary growth-decay pair satisfies*

$$\widehat{\mu}_{{}^*d} = (\widehat{\nu}_{{}^*d})^{1/({}^*\mathfrak{t}\,{}^*d)} > (\widehat{\nu}_{{}^*d})^{1/(\mathfrak{l}\,{}^*d)}$$

where ${}^*\mathfrak{t} = {}^*\{\mathfrak{e}(d_i)\}$ *and where* $\mathfrak{l} < \mathfrak{t}$*.*

Proof. Clear from the definition of best polynomial class and the type $\mathfrak{t}$. □

For T numbers, one writes for each finite degree d

$$\mathfrak{e}(d) = d^{\mathfrak{t}_d - 1}$$

and then the type is defined

$$\mathfrak{t} = \limsup \mathfrak{t}_d \in [1, \infty].$$

Theorem 43. *Let* θ *be a* T*-number of type* $\mathfrak{t} \in [1, \infty]$*. Then for every infinite degree* ${}^*d$ *best polynomial class* ${}^*\widehat{g}_{{}^*d}$*, the associated ordinary growth-decay pair satisfies*

$$\widehat{\mu}_{{}^*d} = (\widehat{\nu}_{{}^*d})^{1/{}^*d^{{}^*\mathfrak{t}}} > (\widehat{\nu}_{{}^*d})^{1/{}^*d^{\mathfrak{l}}}$$

where ${}^*\mathfrak{t} = {}^*\{\mathfrak{t}_{d_i}\}$ *and where* $\mathfrak{l} < \mathfrak{t}$*.*

Proof. From the definitions:

$$|{}^*\widehat{g}_{{}^*d}(\theta)|^{1/{}^*d} = (\mathfrak{h}({}^*\widehat{g}_{{}^*d})^{-1})^{{}^*d^{{}^*\mathfrak{t}-1}} = (\mathfrak{h}({}^*\widehat{g}_{{}^*d})^{-1/{}^*d})^{{}^*d^{{}^*\mathfrak{t}}}.$$

Taking ${}^*d^{{}^*\mathfrak{t}}$-roots of both sides and passing to ${}^{\star}\mathbb{PR}_{\varepsilon}$ classes gives the result, since ${}^*\mathfrak{t} > \mathfrak{l}$. □

Finally, the type of a U number is the first integer $\mathfrak{t}$ for which $\mathfrak{e}(\mathfrak{t}) = \infty$. We leave the proof of the following to the reader:

Theorem 44. *Let* θ *be an* U*-number of type* $\mathfrak{t} \in \mathbb{N}$*. Then for every* ${}^*\mathfrak{t}$ *and* ${}^*d \geq \mathfrak{t}$*, there exists a best polynomial class* ${}^*\widehat{g}_{{}^*d}$ *such that the associated ordinary growth-decay satisfies*

$$\widehat{\mu}_{{}^*d} > (\widehat{\nu}_{{}^*d})^{1/{}^*d^{{}^*\mathfrak{t}}}.$$

Thus, we can say that the S, T and U numbers are those for which the decay is related to growth linearly, polynomially and exponentially, respectively, as a function of degree.



Figure 3. Frobenius Nonvanishing Spectra According to Mahler Class

12. Resultant Arithmetic

In this section we present polynomial diophantine approximation as a natural nonlinear extension of classical diophantine approximation (approximation by linear polynomials). We are faced first with the problem of finding the right notions of sum and product in the polynomial ring $\mathbb{Z}[X]$. By "right", we mean that

I. They should be compatible with fractional sum and product in $\mathbb{Q}$ i.e. the map $a/b \mapsto bX - a$ should be a monomorphism modulo multiplication by non zero integer constants. As a consequence, this will ensure that the map,
$$^*\mathbb{Z}(\theta) \to {}^*\mathfrak{Z}(\theta)_1 \subset {}^*\mathfrak{Z}(\theta), \quad {}^*n \mapsto {}^*nX - {}^*n^\perp,$$
will respect fractional sum and product of numerator denominator pairs of diophantine approximations.

II. If α, β are algebraic and $f(\alpha) = 0$, $g(\beta) = 0$ then $\alpha + \beta$ should be a root of the sum of f and g, and $\alpha\eta$ should be a root of the product of f and g.

Regrettably, neither the Cauchy nor the Dirichlet products satisfy the above criteria.

We begin by defining the sought after product, which we call the **resultant product**. In what follows, denote by
$$\mathfrak{Z} := \{f \in \mathbb{Z}[X] |\ \deg f \geq 1\} \cup \{1\},$$
a monoid with respect to the Cauchy product. Consider elements $f, g \in \mathfrak{Z}$:
$$f(X) = a_m X^m + \cdots + a_0 = a_m \prod_{i=1}^{m} (X - \alpha_i), \quad g(X) = b_n X^n + \cdots + b_0 = b_n \prod_{j=1}^{n} (X - \beta_j).$$
If $\deg(f), \deg(g) \geq 1$, their resultant product is defined
$$(f \otimes g)(X) := a_m^n b_n^m \prod (X - \alpha_i \beta_j);$$
otherwise, if either f or g is 1, it is defined to be 1. Note that $\deg(f \otimes g) = mn = \deg(f)\deg(g)$. The resultant product is clearly commutative and associative, and the polynomial $1_{\otimes}(X) := X - 1$ acts as the identity. The proof of the Theorem which follows was suggested to us by Gregor Weingart.

Theorem 45. *If $f, g \in \mathfrak{Z}$ then $f \otimes g \in \mathfrak{Z}$.*

Proof. We assume that $f, g \neq 1$. Consider the homogenization of f with respect to the new variable Y:
$$f_X(Y) = Y^m f(X/Y) = \sum_{i=0}^{m} \big(a_{m-i} X^{m-i}\big) Y^i,$$
viewed as a polynomial in $\big(\mathbb{Z}[X]\big)[Y]$ i.e. in the single variable Y. Then
$$(f \otimes g)(X) = (-1)^{mn} a_0^n b_n^m \prod \big((X/\alpha_i) - \beta_j\big) = (-1)^{mn} \mathrm{res}\big(f_X(Y), g(Y)\big),$$
where $\mathrm{res}(\cdot, \cdot)$ denotes the resultant. But the resultant is the determinant of a corresponding Sylvester matrix [1], whose non zero entries consist of the coefficients of $f_X(Y)$ and $g(Y)$. Thus $f \otimes g \in \mathfrak{Z}$ as claimed.

□

The **resultant sum** of $f, g \in \mathfrak{Z}$, is defined
$$(f \oplus g)(X) := a_m^n b_n^m \prod \big(X - (\alpha_i + \beta_j)\big)$$

when $f, g \neq 1$; otherwise it is defined to be 1. The resultant sum defines an element of $\mathfrak{Z}$ by an argument similar to that found in Theorem 45. As in the case of the resultant product, $\deg(f \oplus g) = \deg(f)\deg(g)$. The identity element for $\oplus$ is the polynomial $1_{\oplus}(X) := X$. There is also a **resultant difference**

$$f \ominus g := a_m^n b_n^m \prod \left(X - (\alpha_i - \beta_j)\right)$$

which may be used to formulate the counterpart of Theorem 11.

Proposition 33. *Both the resultant product and the resultant sum distribute over the Cauchy product:*

$$f \otimes (g \cdot h) = (f \otimes g) \cdot (f \otimes h), \quad f \oplus (g \cdot h) = (f \oplus g) \cdot (f \oplus h).$$

In particular, $\mathfrak{Z}$ has the structure of a double semiring (a semiring with respect to each of $\oplus, \otimes$ separately).

Proof. Consider the polynomials $f(X) = a_m \prod_{i=1}^{m}(X - \alpha_i)$, $g(X) = b_n \prod_{j=1}^{n}(X - \beta_j)$ and $h(X) = c_p \prod_{k=1}^{p}(X - \gamma_k)$. Then

$$\begin{aligned} f \otimes (g \cdot h) &= \left\{a_m \prod_{i=1}^{m}(X - \alpha_i)\right\} \otimes \left\{b_n \prod_{j=1}^{n}(X - \beta_j) \cdot c_p \prod_{k=1}^{p}(X - \gamma_k)\right\} \\ &= a_m^{n+p}(b_n c_p)^m \prod (X - \alpha_i\beta_j) \cdot \prod (X - \alpha_i\gamma_k) \\ \\ &= (f \otimes g) \cdot (f \otimes h). \end{aligned}$$

An identical argument shows that the resultant sum distributes over the Cauchy product. ☐

For the simple reason of degree, the resultant product does not distribute over the resultant sum:

$$\begin{aligned} \deg(f \otimes (g \oplus h)) &= \deg(f)\deg(g)\deg(h) \\ &\leq \deg(f)^2 \deg(g)\deg(h) \\ &= \deg((f \otimes g) \oplus (f \otimes h)). \end{aligned}$$

Nevertheless, the subset of linear polynomials $\mathfrak{Z}_1$ is closed with respect to $\oplus$, $\otimes$, and one may verify that elements of $\mathfrak{Z}_1$, $f \otimes (g \oplus h) = (f \otimes g) \oplus (f \otimes h)$. Thus $\mathfrak{Z}_1$ itself is a semiring with respect to the operations $\oplus, \otimes$.

Let

$$\check{\mathfrak{Z}} := \mathfrak{Z}/\sim$$

where $f \sim f' \Leftrightarrow$ there exists $0 \neq n \in \mathbb{Z}$ such that $f = nf'$. Then the operations $\oplus, \otimes$ respect $\sim$ and define operations in $\check{\mathfrak{Z}}$ making the latter a double semiring as well. Denote by $\check{\mathfrak{Z}}_1$ the classes of linear polynomials, which is by the above remarks a semiring with respect to $\oplus, \otimes$.

Proposition 34. *Consider the semiring $\tilde{\mathbb{Q}} = \{(a, b) \in \mathbb{Z}^2 \mid b \neq 0\}$ equipped with the usual fractional laws of addition and multiplication. The bijective map*

$$\tilde{\mathbb{Q}} \hookrightarrow \mathfrak{Z}_1, \quad a/b \mapsto bX - a$$

induces an isomorphism of fields $\mathbb{Q} \cong \check{\mathfrak{Z}}_1$ in which $\times, +, -$ are taken to $\otimes, \oplus, \ominus$.

Proof. The map $(a,b)\mapsto bX-a$ clearly identifies isomorphically $\tilde{\mathbb{Q}}=\{(a,b)|\, b\neq 0\}$ with $\mathfrak{Z}_1$. Quotienting by the multiplicative action of $\mathbb{Z}-\{0\}$ on either side gives the result. ☐

Proposition 35. *The map*

$$\mathfrak{Z}_1\times\mathfrak{Z}\longrightarrow\mathfrak{Z},\quad (l(X),f(X))\mapsto l(X)\otimes f(X)$$

defines an action of $\mathbb{Q}\cong\check{\mathfrak{Z}}_1$ *by* $\oplus$*-isomorphisms of* $\check{\mathfrak{Z}}$.

Proof. The action is clearly well-defined, by isomorphisms since the elements of $\check{\mathfrak{Z}}_1$ are $\otimes$-invertible. ☐

Thus $(\check{\mathfrak{Z}},\oplus,\otimes)$ is a $\mathbb{Q}$-*semialgebra*: a semiring equipped with an action by $\mathbb{Q}$ commuting with $\otimes$.

Note 13. We may regard $\check{\mathfrak{Z}}$ as a generalized field extension of $\mathbb{Q}$. Since $\otimes$ does not distribute over $\oplus$, it is closely related to the notion of a *nonlinear number field*: an extension of $\mathbb{Q}$ defined using a projectivization of the field algebra of $\mathbb{Q}$ with the operations of Cauchy and Dirichlet products. See [15].

The (classes of) monic polynomials in $\mathfrak{Z}$ play the role of integers, the **resultant integers**, which are evidently closed w.r.t. $\oplus$, $\otimes$ and Cauchy product. We denote them $\mathfrak{O}$, and note that $\mathbb{Z}$ is identified via the isomorphism of Proposition 34 with $\check{\mathfrak{O}}_1:=\mathfrak{O}_1/\sim$.

The resultant sum, difference and product extend in the obvious way to ${}^*\tilde{\mathfrak{Z}}$ e.g. ${}^*f\oplus{}^*g={}^*\{f_i\oplus g_i\}$. The next result shows that the normalized growth is supermultiplicative with respect to resultant sums, differences and products.

Proposition 36. *Let* ${}^*f\in{}^*\tilde{\mathfrak{Z}}_{{}^*d}, {}^*g\in{}^*\tilde{\mathfrak{Z}}_{{}^*e}$. *Then*

$$\mu_{{}^*d}({}^*f)\cdot\mu_{{}^*e}({}^*g)\ \leq\ \mu_{{}^*d\cdot{}^*e}({}^*f\oplus{}^*g),\ \mu_{{}^*d\cdot{}^*e}({}^*f\ominus{}^*g),\ \mu_{{}^*d\cdot{}^*e}({}^*f\otimes{}^*g).$$

Proof. We begin with $\otimes$ and prove $\mathfrak{m}_{{}^*d\cdot{}^*e}({}^*f\otimes{}^*g)\leq\mathfrak{m}_{{}^*d}({}^*f)\cdot\mathfrak{m}_{{}^*e}({}^*g)$. It suffices to show that for the non normalized Mahler measure,

$$\mathfrak{m}({}^*f\otimes{}^*g)\leq\mathfrak{m}({}^*f)^{\deg({}^*g)}\cdot\mathfrak{m}({}^*g)^{\deg({}^*f)},$$

and moreover, it is enough to prove this for standard polynomials f,g of degree m, n respectively:

$$\begin{aligned}\mathfrak{m}(f\otimes g)&=a_m^nb_n^m\prod\max(1,|\alpha_i\beta_j|)\\&\leq a_m^n\prod_{i=1}^m\left(\max(1,|\alpha_i|)^n\big(b_n\prod_{j=1}^n\max(1,|\beta_j|)\big)\right)\\&=\mathfrak{m}(f)^n\cdot\mathfrak{m}(g)^m.\end{aligned}$$

As for $\oplus$:

$$\begin{aligned}
\mathfrak{m}(f \oplus g) &= a_m^n b_n^m \prod \max(1, |\alpha_i + \beta_j|) \\
&\leq a_m^n \prod_{i=1}^{m} \left(b_n \prod_{j=1}^{n} (\max(1,|\alpha_i|) + \max(1,|\beta_j|)) \right) \\
&= a_m^n \prod_{i=1}^{m} \left(\max(1,|\alpha_i|)^n \left(b_n \prod_{j=1}^{n} \left(1 + \max(1,|\beta_j|) \cdot \max(1,|\alpha_i|)^{-1} \right) \right) \right) \\
&\leq a_m^n \prod_{i=1}^{m} \left(\max(1,|\alpha_i|)^n \left(b_n \prod_{j=1}^{n} (1 + \max(1,|\beta_j|)) \right) \right) \\
&\leq a_m^n \prod_{i=1}^{m} \left(\max(1,|\alpha_i|)^n \left(b_n \prod_{j=1}^{n} 2\max(1,|\beta_j|) \right) \right) \\
&= 2^{mn} \mathfrak{m}(f)^n \cdot \mathfrak{m}(g)^m.
\end{aligned}$$

Applying the above calculation to ${}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*d}, {}^*g \in {}^*\tilde{\mathfrak{Z}}_{{}^*e}$ gives

$$\mathfrak{m}_{{}^*d{}^*e}({}^*f \oplus {}^*g) \leq 2\mathfrak{m}_{{}^*d}({}^*f) \cdot \mathfrak{m}_{{}^*e}({}^*g).$$

The inequality for $\oplus$ follows immediately on passage to growth classes. The corresponding $\ominus$ inequality is proved in much the same way. □

13. Polynomial Ideological Arithmetic I: Finite Degree

Given ${}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)$ and ${}^*g \in {}^*\tilde{\mathfrak{Z}}(\eta)$, in this section we look for conditions which ensure that

$${}^*f \otimes {}^*g \in {}^*\tilde{\mathfrak{Z}}(\theta\eta), \quad {}^*f \oplus {}^*g \in {}^*\tilde{\mathfrak{Z}}[X](\theta + \eta), \quad {}^*f \ominus {}^*g \in {}^*\tilde{\mathfrak{Z}}[X](\theta - \eta).$$

In other words, we seek to formulate polynomial ideological composition laws that are compatible with the linear ideological arithmetic viz. Theorem 11.

It is worth recording the following "ideal-theoretic" arithmetic for elements of $\bar{\mathbb{Q}} \cap \mathbb{R}$. Given $\alpha \in \bar{\mathbb{Q}} \cap \mathbb{R}$, let $\llparenthesis\alpha\rrparenthesis$ denote the Cauchy semigroup of $({}^*\tilde{\mathfrak{Z}}, \cdot)$ generated by the minimal polynomial $m_\alpha(X)$ of α: that is,

$$\llparenthesis\alpha\rrparenthesis := \{ {}^*f \in {}^*\tilde{\mathfrak{Z}} |\ {}^*f(\alpha) = 0 \}.$$

Note that $\llparenthesis\alpha\rrparenthesis \subsetneqq {}^*\tilde{\mathfrak{Z}}(\alpha)$.

Proposition 37. *For all* $\alpha, \beta \in \bar{\mathbb{Q}} \cap \mathbb{R}$,

$$\llparenthesis\alpha\rrparenthesis \otimes \llparenthesis\beta\rrparenthesis \subset \llparenthesis\alpha\beta\rrparenthesis, \quad \llparenthesis\alpha\rrparenthesis \oplus \llparenthesis\beta\rrparenthesis \subset \llparenthesis\alpha + \beta\rrparenthesis, \quad \llparenthesis\alpha\rrparenthesis \ominus \llparenthesis\beta\rrparenthesis \subset \llparenthesis\alpha - \beta\rrparenthesis.$$

Proof. Trivial. □

Consider the Frobenius growth ideology on $\llparenthesis\alpha\rrparenthesis$

$$\llparenthesis\alpha\rrparenthesis^{\bar{\mu}}_{{}^*d} = \llparenthesis\alpha\rrparenthesis \cap {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}(\alpha)_{{}^*d}.$$

There is no interesting decay filtration on $\llparenthesis\alpha\rrparenthesis$ since for all ${}^*f \in \llparenthesis\alpha\rrparenthesis$, $\nu({}^*f) = -\infty$.

Proposition 38. *Let* $\alpha \in \bar{\mathbb{Q}} \cap \mathbb{R}$. *If* $\bar{\mu} \geq \bar{\nu}^{1/{}^*d}$ *then*

$$(51) \qquad {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\alpha)_{{}^*d} = \llparenthesis\alpha\rrparenthesis^{\bar{\mu}}_{{}^*d}.$$

Proof. If there were ${}^*f \in {}^*\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}}(\alpha)_{{}^*d} - (\!|\alpha|\!)^{\bar{\mu}}_{{}^*d}$ a best class then

$$\bar{\mu}_{{}^*d}({}^*f) > \bar{\mu} \geq \bar{\nu}^{1/{}^*d} \geq \bar{\nu}_{{}^*d}({}^*f)^{1/{}^*d}$$

which contradicts the result of Theorem 38. □

We will call the triple $(\bar{\mu}, \bar{\nu}, {}^*d)$ ***A*-exact** if (51) holds: otherwise we call it ***A*-approximative**.

We will now describe $\diamondtimes$-ideological arithmetic for arbitrary real numbers, a discussion which will comprise the rest of this section and the next. All of the statements proved here have counterparts for $\diamondplus, \diamondminus$ whose formulations and proofs can be produced, *mutatis mutandis.* In this section, we restrict ourselves to finite degree polynomials i.e. elements of ${}^*\mathfrak{Z}$. The infinite degree case presents additional complications which will be deferred to §14. We develop first the ordinary ideological arithmetic, from which the Frobenius ideological arithmetic follows as an immediate corollary.

Let $\theta \in \mathbb{R}$ and suppose that ${}^*f \in {}^*\mathfrak{Z}(\theta)_d$. Denote by $\mathfrak{r}({}^*f)$ the set of $\deg {}^*f \leq d$ roots (repeated according to multiplicity). Then there exist ${}^*\alpha_{i_k} \in \mathfrak{r}({}^*f)$, $k = 1, \dots l$, with

$$\theta \simeq {}^*\alpha_{i_k}$$

for all $k = 1, \dots l \leq \deg {}^*f$. If $l = 1$ we will say that ${}^*f \in {}^*\mathfrak{Z}(\theta)$ is a **simple** polynomial diophantine approximation of θ. Write ${}^*\delta_{i_k} = \theta - {}^*\alpha_{i_k} \in {}^*\mathbb{C}_{\varepsilon}$, $k = 1, \dots l$, and define

$$\zeta_d({}^*f) := \langle \prod_{k=1}^{l} {}^*\delta_{i_k} \rangle^{1/d}.$$

Lemma 4. *Let* ${}^*f \in {}^*\mathfrak{Z}(\theta)_d$. *Then*

$$\nu_d({}^*f) = \mathfrak{m}_d({}^*f) \cdot \zeta_d({}^*f).$$

Proof. It is enough prove the claim when $\deg({}^*f) = d$. Assume that ${}^*f$ is simple and write ${}^*f(X) = {}^*a_d \prod_{i=1}^{d}(X - {}^*\alpha_i)$. Suppose that $\theta \simeq {}^*\alpha = {}^*\alpha_1$. Then

$$\nu_d({}^*f) = \langle |{}^*a_d \prod_{i=1}^{d} (\theta - {}^*\alpha_i)| \rangle^{1/d} = \left(|{}^*a_d| \prod_{i=2}^{d} |\theta - {}^*\alpha_i| \right)^{1/d} \cdot \zeta_d({}^*f). \tag{52}$$

The only factors on the right hand side of (52) which act nontrivially by multiplication on $\zeta_d({}^*f) \in {}^*\mathbb{PR}_{\varepsilon}$ are $|{}^*a_d|^{1/d}$ (if the latter is infinite) and those of the form $|\theta - {}^*\alpha_i|^{1/d}$ where ${}^*\alpha_i \in {}^*\mathbb{C} - {}^*\mathbb{C}_{\mathrm{fin}}$, in which case $|\theta - {}^*\alpha_i|^{1/d} \cdot \zeta_d({}^*f) = |{}^*\alpha_i|^{1/d} \cdot \zeta_d({}^*f)$. It follows immediately that

$$\left(|{}^*a_d| \prod_{i=2}^{d} |\theta - {}^*\alpha_i| \right)^{1/d} \cdot \zeta_d({}^*f) = \mathfrak{m}_d({}^*f) \cdot \zeta_d({}^*f),$$

where we note that

- For $\theta \neq 0$, $|{}^*\alpha|^{1/d} \in {}^*\mathbb{R}_{\mathrm{fin}} - {}^*\mathbb{R}_{\epsilon}$ and so acts trivially on $\zeta_d({}^*f)$).
- For $\theta = 0$, $|{}^*\alpha|^{1/d} \in {}^*\mathbb{R}_{\epsilon}$ does not occur in $\mathfrak{m}_d({}^*f)$.

The argument in the non simple case is identical. □

Lemma 5. *Let* $\theta, \eta \in \mathbb{R}$ *and let* ${}^*f \in {}^*\mathfrak{Z}(\theta)_d$, ${}^*g \in {}^*\mathfrak{Z}(\eta)_e$. *Then*

(53)

$$\nu_{de}({}^*f \diamondtimes {}^*g) \leq \left(\mathfrak{m}_d({}^*f)^{1-e^{-1}} \mathfrak{m}_e({}^*g) \cdot \nu_d({}^*f)^{e^{-1}} \right) + \left(\mathfrak{m}_d({}^*f) \mathfrak{m}_e({}^*g)^{1-d^{-1}} \cdot \nu_e({}^*g)^{d^{-1}} \right).$$

Proof. We will show, equivalently, that the unnormalized decay

$$\nu({}^*f \diamondtimes {}^*g) = \nu_{de}({}^*f \diamondtimes {}^*g)^{de}$$

satisfies

$$\nu({}^*f \diamondtimes {}^*g) \leq \big(\mathfrak{m}({}^*f)^{e-1}\mathfrak{m}({}^*g)^{d} \cdot \nu({}^*f)\big) + \big(\mathfrak{m}({}^*f)^{e}\mathfrak{m}({}^*g)^{d-1} \cdot \nu({}^*g)\big).$$

In what follows we will denote $\zeta({}^*f) = \zeta_d({}^*f)^d$, $\zeta({}^*g) = \zeta_e({}^*g)^e$. Without loss of generality we may assume that $\deg({}^*f) = d$ and $\deg({}^*g) = e$. Assume first that ${}^*f$, ${}^*g$ are simple and $\not\equiv 0$. Then there exist unique roots, say ${}^*\alpha = {}^*\alpha_1$ and ${}^*\beta = {}^*\beta_1$, so that $\theta = {}^*\alpha + {}^*\delta$, $\eta = {}^*\beta + {}^*\varepsilon$. By Lemma 4, $\nu({}^*f) = \mathfrak{m}({}^*f) \cdot \zeta({}^*f)$, $\nu({}^*g) = \mathfrak{m}({}^*g) \cdot \zeta({}^*g)$. Now

$$|({}^*f \diamondtimes {}^*g)(\theta\eta)| = a_d^e b_e^d \left(\prod_{i=2}^{d} \Big(\prod_{j=1}^{e} |\theta\eta - {}^*\alpha_i {}^*\beta_j| \Big) \right) \times \prod_{j=1}^{e} (\theta\eta - {}^*\alpha {}^*\beta_j). \tag{54}$$

The last factor in (54) can be written

$$\prod_{j=1}^{e} |\theta\eta - {}^*\alpha {}^*\beta_j| = |{}^*\alpha {}^*\varepsilon + {}^*\beta {}^*\delta + {}^*\delta {}^*\varepsilon| \cdot \prod_{j=2}^{e} |\theta\eta - {}^*\alpha {}^*\beta_j|.$$

Projecting $|{}^*\alpha {}^*\varepsilon + {}^*\beta {}^*\delta + {}^*\delta {}^*\varepsilon|$ to ${}^*\mathbb{P}\mathbb{R}_{\varepsilon}$ gives

$$\langle |{}^*\alpha {}^*\varepsilon + {}^*\beta {}^*\delta + {}^*\delta {}^*\varepsilon| \rangle \leq \zeta({}^*f) + \zeta({}^*g)$$

and therefore

$$\begin{aligned} \Big\langle \prod_{j=1}^{e} |\theta\eta - {}^*\alpha {}^*\beta_j| \Big\rangle &\leq \left(\prod_{j=2}^{e} |\theta\eta - {}^*\alpha {}^*\beta_j| \right) \cdot \big(\zeta({}^*f) + \zeta({}^*g) \big) \\ &= \left(|{}^*\alpha|^{e-1} \prod_{j=2}^{e} \big| \big(1 + ({}^*\delta/{}^*\alpha)\big)\eta - {}^*\beta_j \big| \right) \cdot \big(\zeta({}^*f) + \zeta({}^*g) \big) \\ &= \left(|{}^*\alpha|^{e-1} \prod_{j=2}^{e} \big| \eta - {}^*\beta_j \big| \right) \cdot \big(\zeta({}^*f) + \zeta({}^*g) \big) \\ &\leq |{}^*\alpha|^{e} \cdot \mathfrak{m}({}^*g) \cdot \big(\zeta({}^*f) + \zeta({}^*g) \big). \end{aligned}$$

In the above, we were able to insert an extra factor $|{}^*\alpha|$, and replace $|\eta - {}^*\beta_j|$ by $|{}^*\beta_j|$, since both $|{}^*\alpha|$ and $|\eta|$ belong to $({}^*\mathbb{R}_{\mathrm{fin}})^{\times}_{+}$ and hence act trivially by multiplication in ${}^*\mathbb{P}\mathbb{R}_{\varepsilon}$. The remaining factors $\prod_{j=1}^{e} |\theta\eta - {}^*\alpha_i {}^*\beta_j|$, $i = 2, \ldots d$, appearing in (54) can be treated similarly, as follows:

$\boxed{\textit{Case 1}}$ ${}^*\alpha_i$ is bounded, non infinitesimal. Write $|\theta\eta - {}^*\alpha_i {}^*\beta_j| = |{}^*\alpha_i||(\theta/{}^*\alpha_i)\eta - {}^*\beta_j|$. We may replace $|(\theta/{}^*\alpha_i)\eta - {}^*\beta_j|$ by $|{}^*\beta_j|$ unless the latter is infinitesimal. If $|{}^*\beta_j|$ is infinitesimal, $\mathfrak{m}({}^*g)$ does not include the factor $|{}^*\beta_j|$, and moreover, $|(\theta/{}^*\alpha)\eta - {}^*\beta_j|$ is bounded and so may be omitted anyway. Thus the factor $\prod_{j=1}^{e} |\theta\eta - {}^*\alpha_i {}^*\beta_j|$ may be replaced by $|{}^*\alpha_i|^e \mathfrak{m}({}^*g) \leq \max(|{}^*\alpha_i|, 1)^e \mathfrak{m}({}^*g)$.

$\boxed{\textit{Case 2}}$ ${}^*\alpha_i$ is infinitesimal. There is no need to factor out $|{}^*\alpha_i|$ since it will not appear in $\mathfrak{m}({}^*f)$. The factor $|\theta\eta - {}^*\alpha_i {}^*\beta_j|$ will either be

- infinite: in which case ${}^*\beta_j$ is infinite, and we can replace $|\theta\eta - {}^*\alpha_i {}^*\beta_j|$ by $|{}^*\beta_j|$, with the possible cost of an inequality $\leq$.
- bounded: in which case ${}^*\beta_j$ may be either finite or infinite. We can in either case replace $|\theta\eta - {}^*\alpha_i {}^*\beta_j|$ by $|{}^*\beta_j|$, with the possible cost of an $\leq$ inequality in the event that ${}^*\beta_j$ is infinite.

In this case, it follows then that the factor $\prod_{j=1}^{e} |\theta\eta - {}^*\alpha_i {}^*\beta_j|$ may be replaced by $\mathfrak{m}({}^*g) = \max(|{}^*\alpha_i|, 1)^e \mathfrak{m}({}^*g)$, with the possible cost of the inequality $\leq$.

$\boxed{\textit{Case } 3}$ ${}^*\alpha_i$ is infinite. Factor out $|{}^*\alpha_i|$ as in $\boxed{\textit{Case } 1}$. Since $(\theta/{}^*\alpha)\eta$ is infinitesimal, we can replace $|(\theta/{}^*\alpha)\eta - {}^*\beta_j|$ by $|{}^*\beta_j|$, except when the latter is infinitesimal, but then $|{}^*\beta_j|$ does not appear in $\mathfrak{m}({}^*g)$ and we can omit it, again with a possible cost of an $\leq$ inequality. Once again, the associated factor can be replaced by $\max(|{}^*\alpha_i|, 1)^e \mathfrak{m}({}^*g)$.

From the analysis provided above, we conclude

$$\begin{aligned} \nu({}^*f \diamondtimes {}^*g) &\leq \mathfrak{m}({}^*f)^e \mathfrak{m}({}^*g)^d \cdot \big(\zeta({}^*f) + \zeta({}^*g)\big) \\ &= \big(\mathfrak{m}({}^*f)^{e-1} \mathfrak{m}({}^*g)^d \cdot \nu({}^*f)\big) + \big(\mathfrak{m}({}^*f)^e \mathfrak{m}({}^*g)^{d-1} \cdot \nu({}^*g)\big) \end{aligned}$$

as claimed. Now consider the case of nonsimple polynomial diophantine approximations ${}^*f$, ${}^*g$ in which ${}^*\alpha_{i_1} \simeq \theta, \dots, {}^*\alpha_{i_r} \simeq \theta$ and ${}^*\beta_{j_1} \simeq \eta, \dots {}^*\beta_{j_s} \simeq \eta$. Arguing as we did in the simple case we obtain the inequality

$$\nu({}^*f \diamondtimes {}^*g) \leq \mathfrak{m}({}^*f)^e \cdot \mathfrak{m}({}^*g)^d \prod_{k=1}^{r} \prod_{l=1}^{s} \langle |{}^*\delta_{i_k} + {}^*\varepsilon_{j_l}| \rangle$$

where $\theta = {}^*\alpha_{i_k} + {}^*\delta_{i_k}$, $\eta = {}^*\beta_{j_l} + {}^*\varepsilon_{j_l}$. But

$$\begin{aligned} \prod_{k=1}^{r} \prod_{l=1}^{s} \langle |{}^*\delta_{i_k} + {}^*\varepsilon_{j_l}| \rangle &\leq \prod_{k=1}^{r} \prod_{l=1}^{s} \big(\langle |{}^*\delta_{i_k}| \rangle + \langle |{}^*\varepsilon_{j_l}| \rangle \big) \\ &= \sum \langle |{}^*\delta_{i_1}^{u_1} \cdots {}^*\delta_{i_r}^{u_s} \cdot {}^*\varepsilon_{j_1}^{v_1} \cdots {}^*\varepsilon_{j_s}^{v_s}| \rangle \end{aligned} \tag{55}$$

where the sum is over exponents $(u_1, \dots u_s; v_1, \dots, v_s)$ whose sum is rs and for which $0 \leq u_1, \dots, u_r \leq s$ and $0 \leq v_1, \dots, v_s \leq r$. But each summand in (55) contains a factor of either $\langle \prod_{k=1}^{r} |{}^*\delta_{i_k}| \rangle$ or $\langle \prod_{l=1}^{s} |{}^*\varepsilon_{j_l}| \rangle$, and hence is bounded by one or the other. It follows that

$$\prod_{k=1}^{r} \prod_{l=1}^{s} \langle |{}^*\delta_{i_k} + {}^*\varepsilon_{j_l}| \rangle \leq \langle \prod_{k=1}^{r} |{}^*\delta_{i_k}| \rangle + \langle \prod_{l=1}^{s} |{}^*\varepsilon_{j_l}| \rangle = \zeta({}^*f) + \zeta({}^*g),$$

from which we derive the desired conclusion in this case. ☐

Theorem 46 (Ordinary Ideological Arithmetic–Finite Degree)**.** *There is a well-defined Cauchy bilinear map given by the* $\diamondtimes$*-product*

$${}^*\mathfrak{Z}^{\mu}_{\mu^{e-1}\cdot\nu^{e}}(\theta)_d \times {}^*\mathfrak{Z}^{\nu}_{\mu^{d}\cdot\nu^{d-1}}(\eta)_e \xrightarrow{\diamondtimes} {}^*\mathfrak{Z}^{\mu\cdot\nu}(\theta\eta)_{de}$$

which agrees with the linear ideological product when $d = e = 1$*.*

Proof. This is a fairly straightforward consequence of Lemma 5. If ${}^*f \in {}^*\mathfrak{Z}^{\mu}_{\mu^{e-1}\cdot\nu^{e}}(\theta)_d$, ${}^*g \in {}^*\mathfrak{Z}^{\nu}_{\mu^{d}\cdot\nu^{d-1}}(\eta)_e$ then $\nu_{de}({}^*f \diamondtimes {}^*g) \in {}^*\mathbb{PR}_{\varepsilon}$ since $\nu_d({}^*f) \leq \mu^{e-1} \cdot \nu^e$ and $\nu_e({}^*g) \leq \mu^d \cdot \nu^{d-1}$ imply that the right hand side of (53) is infinitesimal. Moreover, by Proposition 36,

$$\mu \cdot \nu < \mu_{de}({}^*f \diamondtimes {}^*g).$$

☐

Corollary 18 (Frobenius Ideological Arithmetic–Finite Degree)**.** *Let* $d, e \in \mathbb{N}$ *and suppose that* $\bar{\mu} \geq \bar{\nu}$*. Then*

(1) *If $d, e \neq 1$, there is a well-defined Cauchy bilinear pairing*

$$^*\mathfrak{Z}^{\bar{\mu}}_{\bar{\nu}}(\theta)_d \times {}^*\mathfrak{Z}^{\bar{\nu}}_{\bar{\nu}}(\eta)_e \xrightarrow{\diamondplus} {}^*\mathfrak{Z}^{\bar{\nu}}(\theta\eta)_{de}.$$

(2) *If $d = 1$ and $e \neq 1$ there is a well-defined Cauchy bilinear pairing*

$$^*\mathfrak{Z}^{\bar{\mu}}_{\bar{\nu}}(\theta)_1 \times {}^*\mathfrak{Z}^{\bar{\nu}}_{\bar{\mu}}(\eta)_e \xrightarrow{\diamondplus} {}^*\mathfrak{Z}^{\bar{\nu}}(\theta\eta)_e$$

and a similar pairing in the case $d \neq 1$ and $e = 1$.

(3) *If $d = 1 = e$, there is a well-defined Cauchy bilinear pairing*

$$^*\mathfrak{Z}^{\bar{\mu}}_{\bar{\nu}}(\theta)_1 \times {}^*\mathfrak{Z}^{\bar{\nu}}_{\bar{\mu}}(\eta)_1 \xrightarrow{\diamondplus} {}^*\mathfrak{Z}^{\bar{\nu}}(\theta\eta)_1$$

which is just the Frobenius image of the linear (multiplicative) ideological product.

Proof. This is immediate from Theorem 46, where we use Proposition 6 to conclude that $\bar{\mu} \cdot \bar{\nu} = \bar{\nu}$. □

14. Polynomial Growth-Decay Arithmetic II: Infinite Degree

We now consider growth-decay arithmetic for infinite degree polynomial approximations, which takes place within $^*\tilde{\mathfrak{Z}}$. Given $^*f = {}^*\{f_i\} \in {}^*\tilde{\mathfrak{Z}}$, by a **root** of $^*f$ we mean $^*\alpha = {}^*\{\alpha_i\} \in {}^*\bar{\mathbb{Q}}$ such that $^*\{f_i(\alpha_i)\} = 0$ in $^*\bar{\mathbb{Q}}$. The set of roots of $^*f$ (in which roots are repeated according to multiplicity) can be identified with the ultraproduct

$$\mathbb{r}(^*f) := \prod_{\mathfrak{u}} \mathbb{r}(f_i)$$

where $\mathfrak{u}$ is the ultrafilter used throughout this paper to define the ultrapowers $^*\mathbb{Z}, {}^*\mathbb{R}$, etc. In particular, if $^*f \in {}^*\mathfrak{Z}$, $|\mathbb{r}(^*f)| < \infty$; otherwise, for $^*f \in {}^*\tilde{\mathfrak{Z}} - {}^*\mathfrak{Z}$, $|\mathbb{r}(^*f)| > \aleph_0$. The infinite degree case is complicated by the existence of diophantine approximations $^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)$ having infinitely many roots for which $\theta \not\simeq {}^*\alpha$.

Consider $\theta \neq 0$ and $f = a_d \prod(X - \alpha_i) \in \mathbb{Z}[X]$ standard. Define the **θ-Mahler measure** and the **θ-disk measure** by

$$\mathfrak{m}(f)_\theta := a_d \prod \max(|\theta - \alpha_i|, 1), \quad \mathfrak{z}(f)_\theta := a_d \prod_{|\theta-\alpha_i|\leq 1} |\theta - \alpha_i| = |f(\theta)|/\mathfrak{m}_\theta(f).$$

Notice that $\mathfrak{m}(f)_\theta = \mathfrak{m}(f_\theta)$ and $\mathfrak{z}(f)_\theta = \mathfrak{z}(f_\theta)$, where $f_\theta(X) := f(X - \theta)$. These definitions extend in the usual way to $^*f \in {}^*\tilde{\mathfrak{Z}}$. If $^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)_{^*d}$, the analogue of Lemma 4

$$\nu_{^*d}(^*f) = \mathfrak{m}_{^*d}(^*f)_\theta \cdot \zeta_{^*d}(^*f)_\theta$$

(where $\zeta(^*f)_\theta := \langle \mathfrak{z}(f)_\theta \rangle$ and the sub index $^*d$ indicates the degree normalized versions) follows by definition.

Given $f \in \mathbb{Z}[X]$, a θ-**disk root** is a root α of f for which $|\theta - \alpha| < 1$. We may extend this definition to $^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)$, declaring that a root $^*\alpha$ of $^*f$ is a θ-disk root if $|\theta - {}^*\alpha| < 1$. Let $\Delta_\theta(^*f)$ denote the set of θ-disk roots of $^*f$; note that we may identify $\Delta_\theta(^*f)$ as an ultraproduct:

$$\Delta_\theta(^*f) = \prod_{\mathfrak{u}} \Delta_\theta(f_i).$$

It follows from the definitions that

$$\mathfrak{z}(^*f)_\theta = \prod_{\Delta_\theta(^*f)} |\theta - {}^*\alpha| := {}^*\Big\{ \prod_{\alpha \in \Delta_\theta(f_i)} |\theta - \alpha| \Big\}.$$

We will call a root ${}^*\alpha$ of ${}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)$ which satisfies $\theta \simeq {}^*\alpha$ a **θ-root**, the set of which is denoted

$$\mathbb{r}_\theta({}^*f) \subset \Delta_\theta({}^*f).$$

Note that $\mathbb{r}_\theta({}^*f)$ is not in general an ultraproduct: rather, it may be expressed as a union of ultraproducts

$$\mathbb{r}_\theta({}^*f) = \bigcup_{{}^*\varepsilon} \prod_{\mathfrak{u}} \mathbb{r}_{\varepsilon_i}(f_i)$$

where ${}^*\varepsilon = {}^*\{\varepsilon_i\} \in ({}^*\mathbb{R}_\varepsilon)_+$ and $\mathbb{r}_{\varepsilon_i}(f_i)$ is the set of roots of f_i with absolute value less than ε_i.

We distinguish two classes of polynomial diophantine approximations of possibly infinite degree. We say that ${}^*f = {}^*\{f_i\} \in {}^*\tilde{\mathfrak{Z}}(\theta)$ is

- **pure** if $\Delta_\theta({}^*f) - \mathbb{r}_\theta({}^*f)$ is finite or equivalently,

$$\mathbb{r}_\theta({}^*f) = \prod_{\mathfrak{u}} \mathbb{r}_{\varepsilon_i}(f_i) \tag{56}$$

  for some ${}^*\varepsilon \in ({}^*\mathbb{R}_\varepsilon)_+$. In this case, the contributions from $\Delta_\theta({}^*f) - \mathbb{r}_\theta({}^*f)$ to $\zeta_\theta({}^*f)$ can be omitted. The set of pure approximations defines a Cauchy sub semigroup:

$$^*\mathfrak{Z}(\theta) \subset {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{p}} \subset {}^*\tilde{\mathfrak{Z}}(\theta).$$

- **mixed** if $\Delta_\theta({}^*f) - \mathbb{r}_\theta({}^*f)$ is infinite. Here the contributions to $\zeta_\theta(f)$ coming from $\Delta_\theta({}^*f) - \mathbb{r}_\theta({}^*f)$ cannot be ignored. The set of mixed approximations defines a Cauchy sub semigroup

$$^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{m}} \subset {}^*\tilde{\mathfrak{Z}}(\theta).$$

Notice that

$$^*\tilde{\mathfrak{Z}}(\theta) = {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{p}} \sqcup {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{m}}$$

as Cauchy semigroups and ${}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{p}} \cdot {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{m}} \subset {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{m}}$.

In order to state the main theorems regarding infinite degree ideological arithmetic, we need to introduce the following notation:

$$^*\tilde{\mathfrak{Z}}^{\mu}_{\nu^{*e}}(\theta)_{*d} := \{{}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)_{*d} | \ \mu_{*d}({}^*f) > \mu \in {}^*\mathbb{P}\mathbb{R}_\varepsilon, \nu_{*d}({}^*f)^{1/{}^*e} \leq \nu\}$$

where ${}^*e \in {}^*\mathbb{Z}$. The exponential expression $\nu^{{}^*e}$ is not necessarily defined since ${}^*e$ may be infinite, which is why ${}^*\tilde{\mathfrak{Z}}^{\mu}_{\nu^{*e}}(\theta)_{*d}$ required definition.

Theorem 47 (Ordinary Polynomial Ideological Arithmetic–Pure)**.** *Let* ${}^*f \in {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{p}}_{*d}$, ${}^*g \in {}^*\tilde{\mathfrak{Z}}(\eta)^{\mathtt{p}}_{*e}$. *Then*

$$\nu_{*d^*e}({}^*f \diamondtimes {}^*g) \leq \left(\mathfrak{m}_{*d}({}^*f)^{1-{}^*e^{-1}} \mathfrak{m}_{*e}({}^*g) \cdot \nu_{*d}({}^*f)^{{}^*e^{-1}}\right) + \left(\mathfrak{m}_{*d}({}^*f)\mathfrak{m}_{*e}({}^*g)^{1-{}^*d^{-1}} \cdot \nu_{*e}({}^*g)^{{}^*d^{-1}}\right). \tag{57}$$

In particular, there is a well-defined map

$$^*\tilde{\mathfrak{Z}}^{\mu}_{(\mu^{1-{}^*e^{-1}}\cdot\nu)^{*e}}(\theta)^{\mathtt{p}}_{*d} \times {}^*\tilde{\mathfrak{Z}}^{\nu}_{(\mu\cdot\nu^{1-{}^*d^{-1}})^{*d}}(\eta)^{\mathtt{p}}_{*e} \xrightarrow{\diamondtimes} {}^*\tilde{\mathfrak{Z}}^{\mu\cdot\nu}(\theta\eta)_{*d^*e}.$$

Proof. The proof involves the adaptation of the proof of Lemma 5 to the infinite degree case. Let ${}^*f = {}^*\{f_i\} \in {}^*\tilde{\mathfrak{Z}}(\theta)^{\mathtt{p}}_{*d}$, ${}^*g = {}^*\{g_i\} \in {}^*\tilde{\mathfrak{Z}}(\eta)^{\mathtt{p}}_{*e}$. First, by purity, only θ-roots, η-roots are required to define $\zeta_{*d}({}^*f)_\theta$, $\zeta_{*e}({}^*g)_\eta$: that is,

$$\zeta_{*d}({}^*f)_\theta = \langle {}^*\{ \prod_{\alpha_{j_i} \in \mathbb{r}_{x_i}(f_i)} |\theta - \alpha_{j_i}| \} \rangle = \langle {}^*\{ \prod_{\alpha_{j_i} \in \mathbb{r}_{x_i}(f_i)} |\delta_{j_i}| \} \rangle$$

where ${}^*x = {}^*\{x_i\} \in {}^*\mathbb{R}_\varepsilon$ and we have used the fact that $\mathbb{r}_\theta({}^*f)$ is an ultraproduct, see (56)). Similarly,

$$\zeta_{{}^*e}({}^*g)_\eta = \langle {}^*\{ \prod_{\beta_{k_i}\in \mathbb{r}_{y_i}(g_i)} |\eta - \beta_{k_i}| \}\rangle = \langle {}^*\{ \prod_{\beta_{k_i}\in \mathbb{r}_{y_i}(g_i)} |\varepsilon_{k_i}| \}\rangle .$$

Now $|({}^*f \diamondtimes {}^*g)(\theta\eta)|^{1/{}^*d{}^*e}$ is the class of the sequence

$$|a_{d_i}|^{1/d_i} |b_{e_i}|^{1/e_i} \left(\prod_{j_i=1}^{d_i} \prod_{k_i=1}^{e_i} |\theta\eta - \alpha_{j_i}\beta_{k_i}| \right)^{1/d_i e_i} =$$

$$P_i^{1/d_i e_i} \times |a_{d_i}|^{1/d_i} |b_{e_i}|^{1/e_i} \left(\prod_{\alpha_{j_i}\in \mathbb{r}_{x_i}(f_i)} \prod_{\beta_{k_i}\in \mathbb{r}_{y_i}(g_i)} |\alpha_{j_i}\varepsilon_{k_i} + \beta_{k_i}\delta_{j_i} + \delta_{j_i}\varepsilon_{k_i}| \right)^{1/d_i e_i} \tag{58}$$

where P_i is the product of the $|\theta\eta - \alpha_{j_i}\beta_{k_i}|$ where either $\alpha_{k_i} \notin \mathbb{r}_{x_i}(f_i)$ or $\beta_{k_i} \notin \mathbb{r}_{y_i}(g_i)$. Upon passage to ${}^{\star}\mathbb{PR}_\varepsilon$, the expression in the parenthesis in (58) is bounded by either $\langle {}^*\{d_i e_i \prod |\delta_{j_i}|\}\rangle$ or $\langle {}^*\{d_i e_i \prod |\varepsilon_{k_i}|\}\rangle$. In the limit we obtain

$$\nu_{{}^*d{}^*e}({}^*f \diamondtimes {}^*g) \leq {}^*P |{}^*a|^{1/{}^*d} |{}^*b|^{1/{}^*e} \cdot (\zeta_{{}^*d}({}^*f)_\theta^{1/{}^*e} + \zeta_{{}^*e}({}^*g)_\eta^{1/{}^*d}).$$

We are left with analyzing the remaining factor ${}^*P|{}^*a|^{1/{}^*d}|{}^*b|^{1/{}^*e}$, and to this end, we return to a representative sequence

$$|a_{d_i}|^{1/d_i} |b_{e_i}|^{1/e_i} \left(\prod |\theta\eta - \alpha_{j_i}\beta_{k_i}| \right)^{1/d_i e_i} . \tag{59}$$

The first remark is that we can insert a factor of the form

$$\begin{aligned} \left(\prod_{\alpha_{j_i}\in \mathbb{r}_{x_i}(f_i)} \prod_{\beta_{k_i}\in \mathbb{r}_{y_i}(g_i)} |\alpha_{j_i}\beta_{k_i}| \right)^{1/d_i e_i} &\leq |\theta\eta| \left(\prod_{\alpha_{j_i}\in \mathbb{r}_{x_i}(f_i)} \prod_{\beta_{k_i}\in \mathbb{r}_{y_i}(g_i)} \big|1 + |\delta_{j_i}\theta^{-1}|\big| \big|1 + |\varepsilon_{k_i}\eta^{-1}|\big| \right)^{1/d_i e_i} \\ &\longrightarrow 4|\theta\eta|. \end{aligned}$$

We can do this, since we will be multiplying the result with $(\zeta_{{}^*d}({}^*f)_\theta^{1/{}^*e} + \zeta_{{}^*e}({}^*g)_\eta^{1/{}^*d}) \in {}^{\star}\mathbb{PR}_\varepsilon$, and thus we are free to modify (59) by any uniformly bounded factor without changing the bound we have so far on $\nu_{{}^*d{}^*e}({}^*f \diamondtimes {}^*g)$. Now we claim that for each factor $|\theta\eta - \alpha_{j_i}\beta_{k_i}|$ in (59),

$$|\theta\eta - \alpha_{j_i}\beta_{k_i}| = |\alpha_{j_i}\beta_{k_i}||(\theta\eta)/(\alpha_{j_i}\beta_{k_i}) - 1| \leq C \max(1, |\alpha_{j_i}|) \cdot \max(1, |\beta_{k_i}|) \tag{60}$$

for some constant $C > 0$ independent of i and depending only on θ, η. If both $|\alpha_{j_i}|, |\beta_{k_i}| \geq 1$, this is clear since $|(\theta\eta)/(\alpha_{j_i}\beta_{k_i}) - 1|$ has a uniform upper bound for such roots. This is true more generally whenever $|\alpha_{j_i}\beta_{k_i}| \geq 1$. If $|\alpha_{j_i}\beta_{k_i}| < 1$, then $|\theta\eta - \alpha_{j_i}\beta_{k_i}|$ is uniformly bounded by some constant C, which *a fortiori* satisfies (60). Therefore we may replace each factor of $|\theta\eta - \alpha_{j_i}\beta_{k_i}|$ occurring in P_i with $C \max(1, |\alpha_{j_i}|) \cdot \max(1, |\beta_{k_i}|)$, at the cost of an inequality. It follows that

$$\begin{aligned} \nu_{{}^*d{}^*e}({}^*f \diamondtimes {}^*g) \leq& \mathfrak{m}_{{}^*d}({}^*f)\mathfrak{m}_{{}^*e}({}^*g) \cdot (\zeta_{{}^*d}({}^*f)_\theta^{{}^*e^{-1}} + \zeta_{{}^*e}({}^*g)_\eta^{{}^*d^{-1}}) \\ =& \mathfrak{m}_{{}^*d}({}^*f)\mathfrak{m}_{{}^*e}({}^*g)\mathfrak{m}_{{}^*d}({}^*f)_\theta^{-1/{}^*e^{-1}} \cdot \nu({}^*f)^{{}^*e^{-1}} + \\ & \mathfrak{m}_{{}^*d}({}^*f)\mathfrak{m}_{{}^*e}({}^*g)\mathfrak{m}_{{}^*e}({}^*g)_\eta^{-{}^*d^{-1}} \cdot \nu({}^*g)^{{}^*d^{-1}} \end{aligned}$$

What remains is to show is that we may replace the θ and η (normalized) Mahler measures by the usual ones. This follows from an argument similar to that used to produce the inequality (60). □

We now extend the ideological product to include the mixed classes. For $\rho \in (0,1]$ let

$$\Delta^{\rho}_{\theta}({}^{*}f) = \{{}^{*}\alpha \in \Delta_{\theta}({}^{*}f) \mid |\theta - {}^{*}\alpha| < \rho\} \subset \Delta_{\theta}({}^{*}f).$$

Using $\Delta^{\rho}_{\theta}({}^{*}f)$ we define $\mathfrak{z}^{\rho}_{{}^{*}d}(f)_{\theta}$ and $\zeta^{\rho}_{{}^{*}d}({}^{*}f)_{\theta}$.

Lemma 6. *For all* $\rho \in (0,1]$, $\zeta^{\rho}_{{}^{*}d}({}^{*}f)_{\theta} = \zeta_{{}^{*}d}({}^{*}f)_{\theta}$.

Proof. By Wirsing's estimate [27] – in the formulation given in Corollary A.1 in Appendix A.1 of [5] – we have:

$$2^{-({}^{*}d+1)/{}^{*}d} \cdot ({}^{*}d+1)^{-1/(2{}^{*}d)} \cdot \max(1,|\theta|)^{-1} \cdot \left(\frac{|{}^{*}f(\theta)|}{\mathfrak{h}({}^{*}f)}\right)^{1/{}^{*}d} \leq \mathfrak{z}^{\rho}_{{}^{*}d}({}^{*}f)_{\theta}$$

and

$$\mathfrak{z}^{\rho}_{{}^{*}d}({}^{*}f)_{\theta} \leq 2^{({}^{*}d+1)/{}^{*}d} \cdot \binom{{}^{*}d}{[{}^{*}d/2]}^{1/{}^{*}d} \cdot \rho^{-1} \max(1,|\theta|) \cdot \left(\frac{|{}^{*}f(\theta)|}{\mathfrak{h}({}^{*}f)}\right)^{1/{}^{*}d}.$$

Taking $\langle\cdot\rangle$ of either bound gives immediately

$$\zeta^{\rho}_{{}^{*}d}({}^{*}f)_{\theta} = \zeta_{{}^{*}d}({}^{*}f)_{\theta} = \nu_{{}^{*}d}({}^{*}f) \cdot \mu_{{}^{*}d}({}^{*}f).$$

□

Theorem 48 (Ordinary Polynomial Ideological Arithmetic–General)**.** *Let* ${}^{*}f \in {}^{*}\tilde{\mathfrak{Z}}(\theta)_{{}^{*}d}, {}^{*}g \in {}^{*}\tilde{\mathfrak{Z}}(\eta)_{{}^{*}e}$. *Then*

$$\nu_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g) \leq \left(\mathfrak{m}_{{}^{*}d}({}^{*}f)^{1-{}^{*}e^{-1}} \mathfrak{m}_{{}^{*}e}({}^{*}g) \cdot \nu_{{}^{*}d}({}^{*}f)^{{}^{*}e^{-1}}\right) + \left(\mathfrak{m}_{{}^{*}d}({}^{*}f)\mathfrak{m}({}^{*}g)^{1-{}^{*}d^{-1}} \cdot \nu_{{}^{*}e}({}^{*}g)^{{}^{*}d^{-1}}\right).$$

In particular, there is a well-defined map

$${}^{*}\tilde{\mathfrak{Z}}^{\mu}_{(\mu^{1-{}^{*}e^{-1}} \cdot \nu)^{{}^{*}e}}(\theta)_{{}^{*}d} \times {}^{*}\tilde{\mathfrak{Z}}^{\nu}_{(\mu \cdot \nu^{1-{}^{*}d^{-1}})^{{}^{*}d}}(\eta)_{{}^{*}e} \xrightarrow{\diamondtimes} {}^{*}\tilde{\mathfrak{Z}}^{\mu \cdot \nu}(\theta\eta)_{{}^{*}d{}^{*}e}.$$

Proof. Choose, $\rho, \sigma \in (0,1)$ such that

$$\Delta^{\rho}_{\theta}({}^{*}f) \cdot \Delta^{\sigma}_{\eta}({}^{*}g) \subset \Delta_{\theta\eta}({}^{*}f \diamondtimes {}^{*}g).$$

Now $\nu_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g) = \mathfrak{m}_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g)_{\theta\eta} \cdot \zeta_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g)_{\theta\eta}$. By an argument similar to that found in the proof of Theorem 47 by Lemma 6, we may bound

$$\zeta_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g)_{\theta\eta} \leq \zeta^{\rho}_{{}^{*}d}({}^{*}f)_{\theta} + \zeta^{\sigma}_{{}^{*}e}({}^{*}g)_{\eta} = \zeta_{{}^{*}d}({}^{*}f)_{\theta} + \zeta_{{}^{*}e}({}^{*}g)_{\eta} :$$

since $\zeta_{{}^{*}d{}^{*}e}({}^{*}f \diamondtimes {}^{*}g)_{\theta\eta}$ is calculated using elements of $\Delta_{\theta\eta}({}^{*}f \diamondtimes {}^{*}g) \supset \Delta^{\rho}_{\theta}({}^{*}f) \cdot \Delta^{\sigma}_{\eta}({}^{*}g)$, and being a product, omitting the elements of $\Delta_{\theta\eta}({}^{*}f \diamondtimes {}^{*}g)$ not belonging to $\Delta_{\theta}({}^{*}f) \cdot \Delta_{\eta}({}^{*}g)$ makes the result larger. The rest of the proof follows that of Theorem 47. □

Passing to Frobenius classes gives:

Theorem 49 (Frobenius Polynomial Ideological Arithmetic)**.** *Let* ${}^{*}d, {}^{*}e \in {}^{*}\mathbb{N}$ *and* $\bar{\mu} \geq \bar{\nu}$. *There is a well-defined map*

$${}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}^{{}^{*}e}}(\theta)_{{}^{*}d} \times {}^{*}\tilde{\mathfrak{Z}}^{\bar{\nu}}_{\bar{\nu}^{{}^{*}d}}(\eta)_{{}^{*}e} \xrightarrow{\diamondtimes} {}^{*}\tilde{\mathfrak{Z}}^{\bar{\nu}}(\theta\eta)_{{}^{*}d{}^{*}e}.$$

If either ${}^{*}d$ *or* ${}^{*}e$ *is 1, we may adjust the indices in the manner of item (2) of Corollary 18.*

Let $\mu \geq \nu$ resp. $\bar{\mu} \geq \bar{\nu}$ and define the relations

$$\theta\, {}^{*d}_{\mu}\!\otimes{}^{*e}_{\nu}\, \eta, \quad \theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \eta$$

whenever a product of the type defined in Theorem 48 resp. Theorem 49 is nontrivial. Note that it is not necessarily the case that $\theta\, {}^{*d}_{\mu}\!\otimes{}^{*e}_{\nu}\, \eta \Rightarrow \theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \eta$ e.g. recall Note 11. By Corollary 16 we have:

Corollary 19. *Suppose that* $\gamma \geqslant \theta$ *and* $\xi \geqslant \eta$ *by* $R, S \in \mathsf{Rat}_{\mathbb{Z}}$*, respectively. Then*

$$\theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \eta \quad \Longrightarrow \quad \gamma\, {}^{*d\cdot\deg(R)}_{\bar{\mu}}\!\otimes{}^{*e\cdot\deg(R)}_{\bar{\nu}}\, \xi.$$

There are nontrivial polynomial ideological products in degree 1: this just corresponds to the case of the linear ideological product. In what follows, we will show that there exist nontrivial products for (infinite degree) polynomial classes. The following Proposition shows that A-numbers are always composable with other reals, provided the relevant indices are A-exact (the latter term was defined at the beginning of §13).

Theorem 50. *Let* α *be an* A*-number and* $\theta \in \mathbb{R}$.

- *Suppose that* ${}^{*}\breve{d} > {}^{*}\breve{e}$ *and* ${}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}^{*e}}(\theta)_{*d}$ *is nonempty. Then* $\theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \alpha$.
- *Suppose the pair* $(\bar{\mu}, \bar{\nu}^{*e})$ *is* A*-exact and* ${}^{*}\tilde{\mathfrak{Z}}^{\bar{\nu}}_{\bar{\nu}^{*d}}(\theta)_{*e}$ *is not empty. Then* $\alpha\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \theta$.

Proof. If ${}^{*}\breve{d} > {}^{*}\breve{e}$ then $\bar{\nu} > (\bar{\nu}^{*d})^{1/{}^{*}e}$, so the triple $(\bar{\mu}, \bar{\nu}^{*d}, {}^{*}e)$ is A-exact and $\theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \alpha$. The second bulleted item is trivial. ☐

The next result is negative: it asserts that A-numbers on the A-approximative indices, as well as S-numbers, may not be composed with one another.

Theorem 51. *Let* θ *be either an* A*-number or an* S*-number and let* $\eta \in \mathbb{R}$*. Suppose that* $\bar{\mu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ *is not an* ${}^{*}e$*th root of unity and that if* θ *is an* A*-number then the triple* $(\bar{\mu}, \bar{\nu}^{*e}, {}^{*}d)$ *is* A*-approximative. If* $\theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \eta$ *then*

$$\bar{\mu}^{1/{}^{*}e} < \bar{\nu}^{1/{}^{*}d}.$$

In particular, $\theta\, {}^{*d}_{\bar{\mu}}\!\otimes{}^{*e}_{\bar{\nu}}\, \eta$ *does* **not** *hold when*

1. ${}^{*}d = d < \infty$ *or* ${}^{*}\breve{d} < {}^{*}\breve{e}$.
2. η *is either an* A*-number for which the triple* $(\bar{\nu}, \bar{\nu}^{*d}, {}^{*}e)$ *is* A*-approximative, or an* S*-number.*

Proof. Let ${}^{*}f \in {}^{*}\tilde{\mathfrak{Z}}^{\bar{\mu}}_{\bar{\nu}^{*e}}(\theta)_{*d}$; if θ is an A-number then we assume further that ${}^{*}f$ does not belong to the semigroup $(\!|\theta|\!)$ generated by the minimal polynomial of θ (see the paragraph preceding Proposition 37). By the classification of the Frobenius nonvanishing spectra,

$$\bar{\mu}_{*d}({}^{*}f) \leq \bar{\nu}_{*d}({}^{*}f)^{1/{}^{*}d},$$

where the inequality is strict if θ is an A-number. Since $\bar{\mu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ is not an ${}^{*}e$th root of 1, $\bar{\mu}^{1/{}^{*}e} < \bar{\mu}_{*d}({}^{*}f)^{1/{}^{*}e}$. Moreover, by the hypotheses on ${}^{*}f$,

$$\bar{\mu}^{1/{}^{*}e} < \bar{\mu}_{*d}({}^{*}f)^{1/{}^{*}e} \leq \bar{\nu}_{*d}({}^{*}f)^{1/{}^{*}d{}^{*}e} \leq \bar{\nu}^{1/{}^{*}d}. \tag{61}$$

(The second inequality is not strict to allow for the possibility that $\bar{\mu}_{*d}({}^{*}f)$ is a ${}^{*}e$th root of unity.) As a consequence,

$$\bar{\mu} < \bar{\mu}^{1/{}^{*}e} < \bar{\nu}^{1/{}^{*}d} \leq \bar{\mu}^{1/{}^{*}d}. \tag{62}$$

If ${}^*d = d \in \mathbb{N}$ then $\bar{\mu}^{1/d} = \bar{\mu}$ and (62) yields $\bar{\mu} < \bar{\mu}$, contradiction. If ${}^*\breve{d} < {}^*\breve{e}$, then $\bar{\mu}^{{}^*d/{}^*e}$ is defined since ${}^*d/{}^*e$ is infinitesimal and

$$\bar{\mu} < \bar{\mu}^{{}^*d/{}^*e} \leq \bar{\mu}^{1/{}^*e} < 1.$$

But (62) implies that $\bar{\mu}^{{}^*d/{}^*e} < \bar{\mu}$, contradiction. Finally, suppose that η is either an A-number for which the triple $(\bar{\nu}, \bar{\nu}^{{}^*d}, {}^*e)$ is not A-exact, or an S-number. By (61) we have that

$$\bar{\mu}^{1/{}^*e} < \bar{\nu}^{1/{}^*d}. \tag{63}$$

The analogue of (61) for the triple $(\bar{\nu}, \bar{\nu}^{{}^*d}, {}^*e)$ gives $\bar{\nu}^{1/{}^*d} \leq \bar{\nu}^{1/{}^*e} \leq \bar{\mu}^{1/{}^*e}$, implying with (63) that $\bar{\mu}^{1/{}^*e} < \bar{\mu}^{1/{}^*e}$, again, contradiction. □

In what follows consider the special case $\bar{\mu} = \bar{\nu}$.

Lemma 7. *Let* η *be trascendental. Then for every infinite best degree* ${}^*e \in {}^*\mathbb{N} - \mathbb{N}$*, there exists* $\bar{\nu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ *such that for all* ${}^*d$ *with* ${}^*\breve{d} < {}^*\breve{e}$,

$${}^*\tilde{\mathfrak{Z}}^{\bar{\nu}}_{\bar{\nu}^{{}^*d}}(\eta)_{{}^*e} \neq \{1\}.$$

If θ *is either a* T*-number or a* U*-number then we may take* ${}^*\breve{d} = {}^*\breve{e}$ *as well. If* θ *is a* U*-number then we may take* ${}^*\breve{d} = {}^*\breve{e} = \breve{1}$.

Proof. Let ${}^*\widehat{g}$ be a best class for θ of infinite degree ${}^*e$. Then $\bar{\mu}_{{}^*e}({}^*\widehat{g}) \geq \bar{\nu}_{{}^*e}({}^*\widehat{g})^{1/{}^*e}$ and since ${}^*e$ is infinite, $\bar{\nu}_{{}^*e}({}^*\widehat{g})^{1/{}^*d} < \bar{\mu}_{{}^*e}({}^*\widehat{g})$ for all ${}^*d$ with ${}^*\breve{d} < {}^*\breve{e}$. Then the choice

$$\bar{\nu} \in [\bar{\nu}_{{}^*e}({}^*\widehat{g})^{1/{}^*d}, \bar{\mu}_{{}^*e}({}^*\widehat{g}))$$

will work. If θ is either a T-number or a U-number and ${}^*\breve{d} = {}^*\breve{e}$ then $\bar{\mu}_{{}^*d}({}^*\widehat{g}) > \bar{\nu}_{{}^*d}({}^*\widehat{g})^{1/{}^*d}$ and we may select $\bar{\nu}$ in the nontrivial interval $[\bar{\nu}_{{}^*d}({}^*\widehat{g})^{1/{}^*d}, \bar{\mu}_{{}^*d}({}^*\widehat{g}))$. If θ is a U-number then we use a degree d best class, $d \geq \mathfrak{t} =$ the type of η, and we may select $\bar{\nu}$ in the nontrivial interval $[\bar{\nu}_{\breve{1}}({}^*\widehat{g}), \bar{\mu}_{\breve{1}}({}^*\widehat{g}))$. □

Theorem 52. *Let* θ *be a* U*-number and let* η *be either a* T*-number or a* U*-number. Then there exists* ${}^*d \in {}^*\mathbb{N} - \mathbb{N}$ *and* $\bar{\mu} \in {}^{\star}\overline{\mathbb{PR}}_{\varepsilon}$ *such that* $\theta \; {}^{{}^*d}_{\bar{\mu}}\!\otimes^{{}^*d}_{\bar{\mu}} \eta$. *If both* θ *and* η *are* U*-numbers than we may choose* ${}^*d = d$ *finite.*

Proof. Choose ${}^*d = {}^*e$ and ${}^*\widehat{g}$ with regard to η as in Lemma 7. Since θ is a U-number, for d sufficiently large, the exponent $\mathfrak{e}(d, H)$ tends to ∞ as $H \to \infty$. Now let ${}^*H := \mathfrak{h}({}^*\widehat{g})$; thus $[{}^*H^{-1/{}^*d}] = \bar{\mu}_{{}^*d}({}^*\widehat{g})$. Let ${}^*f$ minimize $|{}^*f(\theta)|$ amongst ${}^*f \in {}^*\tilde{\mathfrak{Z}}_{{}^*d}$ having $\mathfrak{h}({}^*f) \leq {}^*H$. Let ${}^*\mathfrak{e} := \mathfrak{e}({}^*d, {}^*H)$; then

$$\bar{\mu}_{{}^*d}({}^*f) \geq \bar{\mu}_{{}^*d}({}^*\widehat{g}) = \bar{\nu}_{{}^*d}({}^*f)^{1/({}^*d \cdot {}^*\mathfrak{e})}$$

(the last equality follows from the defining equation $|{}^*f(\theta)|^{1/{}^*d} = {}^*H^{-{}^*\mathfrak{e}}$ of ${}^*\mathfrak{e}$). If ${}^*\mathfrak{e}$ is infinite, we are done, for then

$$\bar{\nu}({}^*f)^{1/{}^*d} < \bar{\mu}_{{}^*d}({}^*\widehat{g})$$

so taking

$$\bar{\mu} \in [\bar{\nu}({}^*f)^{1/{}^*d} + \bar{\nu}({}^*g)^{1/{}^*d}, \bar{\mu}_{{}^*d}({}^*\widehat{g}))$$

will give the relation $\theta \; {}^{{}^*d}_{\bar{\mu}}\!\otimes^{{}^*d}_{\bar{\mu}} \eta$. If not, we choose another best class ${}^*\widehat{g}$ having the same degree for which the associated height sequence ${}^*H$ produces an infinite ${}^*\mathfrak{e}$: since θ is a U-number, such a sequence can always be produced. The argument is identical if η is also a U-number and the degree is chosen finite and large enough. □

Corollary 20. *There exist nontrivial ordinary and Frobenius polynomial ideological products, of both infinite degree and finite degree* > 1.

Proof. By Thereom 52 there exist non trival Frobenius products of both finite and infinite degree. But $\theta \; {}^{*}{}^{d}_{\bar{\mu}} \diamondtimes {}^{*}{}^{d}_{\bar{\mu}} \; \eta$ implies $\theta \; {}^{*}{}^{d}_{\mu} \diamondtimes {}^{*}{}^{d}_{\mu} \; \eta$ for all $\mu \in \bar{\mu}$. ☐

15. EPILOGUE: IDEOLOGICAL CLASSES

In this section we define the set of ideological classes and begin the process of merging the latter with the classical ideal class group. The discussion here is not definitive and is intended primarily to provide structural harmony in these closing pages, by explaining the appearance of the product/sum/difference range of ideological arithmetic in terms of the classical product of ideals. A more evolved version of the ideas in this section will appear in the third installment in this series of papers [14], in the context of Galois and Class Field Theory.

Let $K/\mathbb{Q}$ be of finite degree, $\mathbb{K}$ the associated Minkowski space. For $\boldsymbol{z} \in \mathbb{K}$ define the **decoupled diophantine approximation group** by

$$ {}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z}) := {}^{*}\mathcal{O}(\boldsymbol{z}) + {}^{*}\mathcal{O}(\boldsymbol{z})^{\perp} \subset {}^{*}\mathcal{O}. $$

Note that if $\boldsymbol{z} \in \mathbb{K}$ is invertible then

$$ {}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z}) = {}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z}) + {}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z}^{-1}) \subset {}^{*}\mathcal{O}(\boldsymbol{z}, \boldsymbol{z}^{-1})^{\perp}. $$

One can regard ${}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z})$ as the outcome of "decoupling" (making independent) numerator denominator pairs associated to $\boldsymbol{z}$. Note that ${}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}(\boldsymbol{z})$ has a natural ideological structure when $\boldsymbol{z}$ is invertible, defined by the subgroups

$$ {}^{*}{\ulcorner\!\!\llcorner}\mathcal{O}{\urcorner\!\!\lrcorner}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) = \{ {}^{*}\alpha + {}^{*}\beta^{\perp} | {}^{*}\alpha \in {}^{*}\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}), {}^{*}\beta^{\perp} \in {}^{*}\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}^{-1}) \}. \tag{64} $$

In the Theorem which follows we will need Kronecker's criterion ([6], pg. 53) for the existence of inhomogeneous simultaneous approximations, recast in the language of matrix diophantine approximations. Consider an $r \times s$ real matrix Θ and a vector $\boldsymbol{w} = (w_1, \dots, w_r)^T \in \mathbb{R}^r$. Kronecker's criterion declares that the following statements are equivalent:

K1. There exists ${}^{*}\boldsymbol{n}^{\perp} \in {}^{*}\mathbb{Z}^r$, ${}^{*}\boldsymbol{n} \in {}^{*}\mathbb{Z}^s$ such that

$$ \Theta {}^{*}\boldsymbol{n} - {}^{*}\boldsymbol{n}^{\perp} \simeq \boldsymbol{w}. $$

K2. If $\boldsymbol{m} = (m_1, \dots, m_r) \in \mathbb{Z}^r$ satisfies $\boldsymbol{m} \cdot \Theta \in \mathbb{Z}^s$ then $\boldsymbol{m} \cdot \boldsymbol{w} \in \mathbb{Z}$.

Theorem 53. *For* $\theta \in \mathbb{R} - \mathbb{Q}$, ${}^{*}\mathbb{Z}(\theta, \theta^{-1})^{\perp} = {}^{*}\mathbb{Z}$. *Moreover,* ${}^{*}{\ulcorner\!\!\llcorner}\mathbb{Z}{\urcorner\!\!\lrcorner}(\theta) \subsetneqq {}^{*}\mathbb{Z} \Leftrightarrow \theta$ *is a quadratic irrationality.*

Proof. That ${}^{*}\mathbb{Z}(\theta, \theta^{-1})^{\perp} = {}^{*}\mathbb{Z}$ is a consequence of Kronecker's criterion. Indeed, for $\Theta = (\theta, \theta^{-1})$ there is no $m \in \mathbb{Z}$ with $m \cdot \Theta \in \mathbb{Z}^2$, so condition K2 is vacuously true: hence for any $w \in \mathbb{Z}$ we may find ${}^{*}\boldsymbol{n} \in {}^{*}\mathbb{Z}^2$, ${}^{*}\boldsymbol{n}^{\perp} \in {}^{*}\mathbb{Z}$ satisfying K1. Therefore, by a diagonal argument, for any ${}^{*}w = {}^{*}\{w_i\} \in {}^{*}\mathbb{Z}$ we may solve K1 with ${}^{*}w$ in place of w. Suppose now that θ is a quadratic irrationality. In this case we will show that ${}^{*}{\ulcorner\!\!\llcorner}\mathbb{Z}{\urcorner\!\!\lrcorner}(\theta) \cap \mathbb{Z} = \{0\}$: in other words, we will show that for all $0 \neq N \in \mathbb{Z}$, the equation

$$ {}^{*}m + {}^{*}n^{\perp} = N, \quad {}^{*}m \in {}^{*}\mathbb{Z}(\theta), {}^{*}n^{\perp} \in {}^{*}\mathbb{Z}(\theta^{-1}) \tag{65} $$

has no solution. To this end we consider the 2×2 diagonal matrix $\Theta = \mathrm{diag}(\theta, \theta^{-1})$. If there were a solution ${}^{*}n^{\perp} = N - {}^{*}m$ to (65) then the vectors ${}^{*}\boldsymbol{m} = ({}^{*}m, {}^{*}m)$, ${}^{*}\boldsymbol{m}^{\perp} = ({}^{*}m^{\perp}, -{}^{*}n)$ would provide a solution to the inhomogeneous equation

$$ \Theta {}^{*}\boldsymbol{m} - {}^{*}\boldsymbol{m}^{\perp} \simeq \boldsymbol{w} = (0, N\theta^{-1})^T. $$

On the other hand, θ is quadratic $\Leftrightarrow$ there exists a vector $\boldsymbol{p} = (p_1, p_2) \in \mathbb{Z}^2$ with $\boldsymbol{p}\cdot\Theta \in \mathbb{Z}^2$. But such a vector could never satisfy $\boldsymbol{p}\cdot\boldsymbol{w} = p_2 N\theta^{-1} \in \mathbb{Z}$. Therefore there can be no solutions to K1 for the choice $\boldsymbol{w} = (0, N\theta^{-1})^T$. If θ is not quadratic then condition K2 is vacuously satisfied since there exist no vectors $\boldsymbol{p} = (p_1, p_2) \in \mathbb{Z}^2$ with $\boldsymbol{p}\cdot\Theta \in \mathbb{Z}^2$. Thus $\mathbb{Z} \subset {}^{*}\ulcorner\mathbb{Z}\urcorner(\theta)$. By a diagonal argument we can then solve $\Theta^{*}\boldsymbol{n} - {}^{*}\boldsymbol{n}^{\perp} \simeq {}^{*}w$ for any ${}^{*}w \in {}^{*}\mathbb{Z}$. □

Note 14. Theorem 53 *does not* imply that for $\theta \in \mathbb{R} - \mathbb{Q}$ not a quadratic irrationality, ${}^{*}\ulcorner\mathbb{Z}\urcorner(\theta) = {}^{*}\mathbb{Z}$ as ideologies i.e. it is not the case that ${}^{*}\ulcorner\mathbb{Z}\urcorner^{\mu}_{\nu}(\theta) = {}^{*}\mathbb{Z}^{\mu}$ for all μ, ν. Indeed, if $(\mu, \nu) \notin \mathrm{Spec}(\theta)$ then ${}^{*}\ulcorner\mathbb{Z}\urcorner^{\mu}_{\nu}(\theta) = 0$.

We return to the finite extension $K/\mathbb{Q}$. For any $x, y \in \mathcal{O}$, the ultrapower ${}^{*}(x, y)$ of the ideal (x, y) is the ${}^{*}\mathcal{O}$-module generated by x, y. Let $\gamma = \alpha/\beta \in K$, $\alpha, \beta \in \mathcal{O}$. Then

$$ {}^{*}K \supset {}^{*}(\gamma, 1) \supset {}^{*}\ulcorner\mathcal{O}\urcorner(\gamma) \supset {}^{*}(\alpha, \beta) = {}^{*}\mathcal{O}(\alpha, \beta)^{\perp}. \tag{66} $$

Moreover, note that for $\alpha \in \mathcal{O}$,

$$ {}^{*}\ulcorner\mathcal{O}\urcorner(\alpha) = {}^{*}\mathcal{O}, $$

so that the construction ${}^{*}\ulcorner\mathcal{O}\urcorner(\cdot)$ already identifies the ultrapowers of classical principal ideals with the identity ideal ${}^{*}(1) = {}^{*}\mathcal{O}$.

Given $\boldsymbol{z}, \boldsymbol{z}' \in \mathbb{K}$ their corresponding decoupled groups are called **class equivalent** if there exists $\gamma \in K$ such that for all $\boldsymbol{\mu}, \boldsymbol{\nu} \in {}^{*}\mathbb{PK}_{\varepsilon}$,

$$ \gamma \cdot {}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z}) = {}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z}'). \tag{67} $$

This happens, for example, if $\boldsymbol{z}' = \gamma\boldsymbol{z}$. The associated equivalence class is denoted

$$ {}^{*}[\mathcal{O}](\boldsymbol{z}) := \{\gamma \cdot {}^{*}\ulcorner\mathcal{O}\urcorner(\boldsymbol{z}) |\, \gamma \in K\}. $$

By (67), ${}^{*}[\mathcal{O}](\boldsymbol{z})$ inherits the ideological filtration (64) and is called the **ideology class** of ${}^{*}\mathcal{O}(\boldsymbol{z})$. For $\mathfrak{a} = (\alpha, \beta) \subset \mathcal{O}$ a classical ideal we denote

$$ {}^{*}[\mathfrak{a}] := {}^{*}[\mathcal{O}](\alpha/\beta). $$

Let $\mathcal{C}\ell(\mathbb{K})$ the set of ideology classes associated to $\boldsymbol{z} \in \mathbb{K}$ and let $\mathcal{C}\ell(K)$ denote the usual ideal class group. Then there is a canonical inclusion

$$ \mathcal{C}\ell(K) \subset \mathcal{C}\ell(\mathbb{K}), \quad [\mathfrak{a}] \mapsto {}^{*}[\mathfrak{a}]. $$

Recall [10] that ideals $\mathfrak{a} = (\alpha, \beta), \mathfrak{a}' = (\alpha', \beta') \subset \mathcal{O}$ define the same class $[\mathfrak{a}] = [\mathfrak{a}']$ $\Leftrightarrow$ there exists $A \in \mathrm{PGL}_2(\mathcal{O})$ with $A(\gamma) = \gamma'$ where $\gamma = \alpha/\beta$ and $\gamma' = \alpha'/\beta'$. That is, there is a bijection

$$ \mathcal{C}\ell(K) \longleftrightarrow \mathrm{PGL}_2(\mathcal{O})\backslash K. $$

In particular, this bijection "explains" that the 2-generator property of ideals in the Dedekind domain $\mathcal{O}$ is the ideal-theoretic translation of the fact that (the $\mathrm{PGL}_2(\mathcal{O})$-class of) any element $\gamma \in K$ is determined by an (unordered) integral numerator denominator pair.

Theorem 54. *If* $\boldsymbol{z} \Bumpeq_K \boldsymbol{z}'$ *then* ${}^{*}[\mathcal{O}](\boldsymbol{z}) = {}^{*}[\mathcal{O}](\boldsymbol{z}')$ *as ideology classes.*

Proof. If $\boldsymbol{z}' = A(\boldsymbol{z})$ where $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then

$$ {}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z}') = \left\{(c^{*}\alpha^{\perp} + d^{*}\alpha) + (a^{*}\beta^{\perp} + b^{*}\beta) |\, {}^{*}\alpha, {}^{*}\beta \in \mathcal{O}^{\mu}_{\nu}(\boldsymbol{z})\right\} \subset {}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z})^{\mu} z. $$

By symmetry, we have ${}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z}) = {}^{*}\ulcorner\mathcal{O}\urcorner^{\mu}_{\nu}(\boldsymbol{z}')$. □

We conjecture that if ${}^*[\mathcal{O}](\boldsymbol{z}) = {}^*[\mathcal{O}](\boldsymbol{z}')$ as ideology classes then $\boldsymbol{z} \approx_K \boldsymbol{z}'$ as well. If true, then

$$\mathcal{C}\ell(\mathbb{K}) \longleftrightarrow \mathrm{PGL}_2(\mathcal{O})\backslash\mathbb{K}.$$

When $K = \mathbb{Q}$, this would imply that the ideology class set is in bijection with the moduli space of quantum tori (compactified at the cusp).

We would now like to migrate the ideological arithmetic as described in §§5, 8 to a counterpart arithmetic defined on $\mathcal{C}\ell(\mathbb{K})$. To define the latter, it is imperative that we understand multiplication in the ideal class group $\mathcal{C}\ell(K) \leftrightarrow \mathrm{PGL}_2(\mathcal{O})\backslash K$ in terms of the arithmetic of K. Such a description could not involve the fractional product in K alone since for $[\gamma] \in \mathrm{PGL}_2(\mathcal{O})\backslash K$ nontrivial $[\gamma] = [\gamma^{-1}]$ so in general $[\gamma]\cdot[\gamma^{-1}] \neq [1]$. The missing ingredients in this endeavor are the fractional sum and difference.

Given (nonprincipal) ideals $\mathfrak{a}_1 = (\alpha_1, \beta_1)$, $\mathfrak{a}_2 = (\alpha_2, \beta_2) \subset \mathcal{O}$ the product is

$$\mathfrak{a}_1 \cdot \mathfrak{a}_2 = (\alpha_1\alpha_2, \alpha_1\beta_2, \alpha_2\beta_1, \beta_1\beta_2) = (\gamma_1, \gamma_2).$$

It is not a routine matter to describe an explicit relation between the minimal (2-generator) generating set $\{\gamma_1, \gamma_2\}$ of $\mathfrak{a}\cdot\mathfrak{b}$ and the *a priori* generator set $\{\alpha_1\alpha_2, \alpha_1\beta_2, \alpha_2\beta_1, \beta_1\beta_2\}$. Nevertheless, the *a priori* generating set is generated by the numerators and the denominators of the product, sum and difference of $\gamma_1 = \alpha_1/\beta_1$ with $\gamma_2 = \alpha_2/\beta_2$. That is, in the notation of ideology classes:

Proposition 39. *Let $\mathfrak{a}_1, \mathfrak{a}_2$ be as in the previous paragraph. Then*

$$ {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1\gamma_2) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1 + \gamma_2) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1 - \gamma_2) \in {}^*[\mathfrak{a}_1 \cdot \mathfrak{a}_2].$$

Proof. Recall from (66) that ${}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_i) \in {}^*[\mathfrak{a}_i]$, $i = 1, 2$. On the other hand,

$$\begin{aligned} {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1\gamma_2) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1 + \gamma_2) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\gamma_1 - \gamma_2) &= {}^*(\alpha_1\alpha_2, \beta_1\beta_2, \alpha_1\beta_2 + \alpha_2\beta_1, \alpha_1\beta_2 - \alpha_2\beta_1) \\ &= {}^*(\alpha_1\alpha_2, \beta_1\beta_2, \alpha_1\beta_2, \alpha_2\beta_1) \\ &\in {}^*[\mathfrak{a}_1 \cdot \mathfrak{a}_2]. \end{aligned}$$

□

Proposition 39 lays bare the significance of the image of the ideological product (29) belonging to the triple intersection ${}^*\mathcal{O}(\boldsymbol{zw}) \cap {}^*\mathcal{O}(\boldsymbol{z}+\boldsymbol{w}) \cap {}^*\mathcal{O}(\boldsymbol{z}-\boldsymbol{w})$, and suggests that a corresponding product of decoupled diophantine approximations groups be contemplated.

Define the **⊛-product**

$$ {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}) \circledast {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{w}) := {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{zw}) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}+\boldsymbol{w}) + {}^*\ulcorner\!\llcorner\mathcal{O}\lrcorner\!\urcorner(\boldsymbol{z}-\boldsymbol{w})$$

which inherits an ideology from each of the three summands on the right hand side. Taking K-projective classes defines the ⊛-product on ideological classes. By Proposition 39 the ⊛-product coincides with the ideal class product upon restriction to $\mathcal{C}\ell(K)$.

There is no reason to believe in general that ${}^*[\mathcal{O}](\boldsymbol{z}) \circledast {}^*[\mathcal{O}](\boldsymbol{w}) = {}^*[\mathcal{O}](\boldsymbol{x})$ for some $\boldsymbol{x} \in \mathbb{K}$. Nevertheless, there are canonical inclusions from the domains and ranges of the ideological product into each of the three defining ideologies of the

$\circledast$-product:

$$\begin{tikzcd}
{}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \arrow[d, hook] & \times & {}^*\mathcal{O}^{\boldsymbol{\nu}}_{\boldsymbol{\mu}}(\boldsymbol{w}) \arrow[r, "\cdot"] \arrow[d, hook] & {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{zw}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}+\boldsymbol{w}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}-\boldsymbol{w}) \arrow[d, hook] \\
\Big({}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) , & & {}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{w}) \Big) \arrow[r, mapsto] & {}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \circledast {}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{w})
\end{tikzcd}$$

where the first two vertical arrows are given by ${}^*\alpha \mapsto {}^*\alpha + {}^*\alpha^{\perp}$, ${}^*\beta \mapsto {}^*\beta + {}^*\beta^{\perp}$ and the last arrow is defined:

$$ {}^*\xi \mapsto {}^*\xi + {}^*\xi^{\perp \boldsymbol{zw}} + {}^*\xi^{\perp \boldsymbol{z}+\boldsymbol{w}} + {}^*\xi^{\perp \boldsymbol{z}-\boldsymbol{w}}. $$

The bottom horizontal arrow is a pairing of elements and not of sets, so the diagram is not intended to be commutative in any familiar sense.

How might one interpret the ideology class ${}^*[\mathcal{O}](\boldsymbol{z}) \circledast {}^*[\mathcal{O}](\boldsymbol{w})$? As one varies $\boldsymbol{\mu}, \boldsymbol{\nu}$, the images of the ideological product

$$ {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \cdot {}^*\mathcal{O}^{\boldsymbol{\nu}}_{\boldsymbol{\mu}}(\boldsymbol{w}) \subset {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{zw}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}+\boldsymbol{w}) \cap {}^*\mathcal{O}^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}-\boldsymbol{w}) $$

generate a subgroup which inherits the ambient ideological structure. It seems unlikely however that this group is equal to ${}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{z}) \circledast {}^*\ulcorner\mathcal{O}\lrcorner^{\boldsymbol{\mu}}_{\boldsymbol{\nu}}(\boldsymbol{w})$. What seems more plausible is that it will be necessary to expand the domain of definition of the $\circledast$-product from (classes of) decoupled diophantine approximation ideologies, indexed by ($\mathrm{PGL}_2(\mathcal{O})$-classes of) points in $\mathbb{K}$ to a space in which the points of $\mathbb{K}$ are discrete. This is already suggested in the construction of the j-invariant for quantum tori defined in [7],[8], which is defined using a universal j-invariant with expanded domain the Stone space of ultrafilters on the set of hyperfinite subsets of ${}^*\mathbb{Z}^2$.

REFERENCES

[1] Akritas, A. G., Sylvester's Forgotten Form of the Resultant. *Fib. Quart.* **31** (1993), 325–332.

[2] Baker, Alan, *Transcendental Number Theory.* Second edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1990.

[3] Beardon, Alan, *Iteration of Rational Functions.* Graduate Texts in Mathematics **132**, Springer-Verlag, New York, 1991.

[4] Bernik, V.I. & Dodson, M.M., *Metric Diophantine Approximation on Manifolds.* Cambridge Tracts in Mathematics **137**, Cambridge University Press, Cambridge, UK, 1999.

[5] Bugeaud, Yann, *Approximation by Algebraic Numbers.* Cambridge Tracts in Mathematics **160**, Cambridge University Press, Cambridge, 2004.

[6] Cassels, J.W.S., *An Introduction to Diophantine Approximation.* Cambridge Tracts in Mathematics and Mathematical Physics **45**. Cambridge University Press, New York, 1957.

[7] Castaño Bernard, C. & Gendron, T.M., Modular invariant of quantum tori I: definitions nonstandard and standard. arXiv:0909.0143

[8] Castaño Bernard, C. & Gendron, T.M., Modular invariant of quantum tori II: the golden mean. arXiv:1204.2540.

[9] Erdös, P., Representations of real numbers as sums and products of Liouville numbers. *Michigan Math. J.* **9**, Issue 1 (1962), 59-60.

[10] Freitag, Eberhard., *Hilbert Modular Forms.* Springer-Verlag, New York, NY, 1990.

[11] Gendron, T.M., The algebraic theory of the fundamental germ. *Bull. Braz. Math. Soc. (N.S.)* **37** (2006), no. 1, 49–87.

[12] Gendron, T.M., The geometric theory of the fundamental germ. Nagoya Math. J. **190** (2008), 1–34.

[13] Gendron, T.M., Real algebraic number theory I. Diophantine approximation groups, Kronecker foliations and independence.

[14] Gendron, T.M., Real algebraic number theory III. Transcendental Galois Theory. *in preparation.*

[15] Gendron, T.M. & Verjovsky, A., Geometric Galois theory, nonlinear number fields and a Galois group interpretation of the idele class group. Internat. J. Math. **16** (2005), no. 6, 567–593. *Errata* Internat. J. Math. **22** (2011), no. 2, 307–309.

[16] Goodman, Frederick M.; de la Harpe, Pierre & Jones, Vaughan F.R., *Coxeter Graphs and Towers of Algebras.* MSRI Publications **14**, Springer-Verlag, New York, NY, 1989.

[17] Hall, Marshall, On the sum and product of continued fractions. *Ann. of Math.* **48** (4), (1947), 966–993.

[18] Horn, Roger A. & Johnson, Charles R., *Topics in Matrix Analysis.* Cambridge University Press, 1994.

[19] Lang, Serge, *Introduction to Diophantine Approximations.* Second edition. Springer-Verlag, New York, 1995.

[20] Niven, Ivan; Zuckerman, Herbert S. & Montgomery, Hugh L., *An Introduction to the Theory of Numbers.*, 5th edition, John Wiley & Sons, Inc., New York, 1991.

[21] Queffélec, Martine, An introduction to Littlewood's conjecture. *Séminaires & Congrès* **20** (2009), 129–152.

[22] Schmidt, Wolfgang M., *Diophantine Approximation.* Lecture Notes in Mathematics **785**. Springer, Berlin, 1980.

[23] Schweiger, Fritz, *Multidimensional Continued Fractions.* Oxford Science Publications. Oxford University Press, Oxford, UK, 2000.

[24] Shalitt, J.O., Simple continued fractions for some irrational numbers, II, *J. Num. Theory* **14** (1982), 228–231.

[25] Waldschmidt, Michel, Report on some recent advances in Diophantine approximation. www.math.jussieu.fr/ miw/articles/pdf/miwLangMemorialVolume.pdf.

[26] Waldschmidt, Michel, *Diophantine Approximation on Linear Algebraic Groups: Transcendence Properties of the Exponential Function in Several Variables.* Grundlehren der mathematischen Wissenschaften **326**, Springer-Verlag, 2000.

[27] Wirsing, Eduard., Approximation mit algebraischen Zahlen beschränkten Grades. *J. reine ang. Math.* **206**, 67–77.

[28] Zariski, Oscar & Samuel, Pierre, *Commutative Algebra, Volume II.* The University Series in Higher Mathematics. D. Van Nostrand Co., Inc., Princeton, N. J.-Toronto-London-New York, 1960.

[29] Zeckendorf, E., Représentation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas. *Bull. Soc. R. Sci. Liège* **41** (1972), 179–182.

Instituto de Matemáticas – Unidad Cuernavaca, Universidad Nacional Autonoma de México, Av. Universidad S/N, C.P. 62210 Cuernavaca, Morelos, MÉXICO

E-mail address: `tim@matcuer.unam.mx`